

UTMS 2013–6


June 11, 2013

**Structure of
the positive radial solutions for
the supercritical Neumann problem**

$$\varepsilon^2 \Delta u - u + u^p = 0 \text{ in a ball}$$

by

Yasuhito Miyamoto



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

**STRUCTURE OF
THE POSITIVE RADIAL SOLUTIONS FOR
THE SUPERCRITICAL NEUMANN PROBLEM
 $\varepsilon^2 \Delta u - u + u^p = 0$ IN A BALL**

YASUHITO MIYAMOTO

ABSTRACT. We are interested in the structure of the positive radial solutions of the supercritical Neumann problem in a unit ball

$$\begin{cases} \varepsilon^2 (U_{rr} + \frac{N-1}{r} U_r) - U + U^p = 0, & 0 < r < 1, \\ U_r(1) = 0, \\ U > 0, \end{cases} \quad 0 < r < 1,$$

where N is the spatial dimension and $p > p_S := (N+2)/(N-2)$, $N \geq 3$. We show that there exists a sequence $\{\varepsilon_n^*\}_{n=1}^\infty$ ($\varepsilon_1^* > \varepsilon_2^* > \dots \rightarrow 0$) such that this problem has infinitely many singular solutions $\{(\varepsilon_n^*, U_n^*)\}_{n=1}^\infty \subset \mathbb{R} \times (C^2(0,1) \cap C^0(0,1])$ and that the nonconstant regular solutions consist of infinitely many smooth curves in the $(\varepsilon, U(0))$ -plane. It is shown that each curve blows up at ε_n^* and if $p_S < p < p_{JL}$, then each curve has infinitely many turning points around ε_n^* . Here,

$$p_{JL} := \begin{cases} 1 + \frac{4}{N-4-2\sqrt{N-1}} & (N \geq 11), \\ \infty & (2 \leq N \leq 10). \end{cases}$$

In particular, the problem has infinitely many solutions if $\varepsilon \in \{\varepsilon_n^*\}_{n=1}^\infty$. We also show that there exists $\bar{\varepsilon} > 0$ such that the problem has no nonconstant regular solution if $\varepsilon > \bar{\varepsilon}$. The main technical tool is the intersection number between the regular and singular solutions.

CONTENTS

1. Introduction and main results	2
2. Preliminaries	11
2.1. Regular and singular solutions of the limit equation	11
2.2. Nonexistence of the entire solution	13
3. Local and global parametrization results	15

2000 *Mathematics Subject Classification*. Primary 35J25, 25B32; Secondary 34C23, 34B16.

This work was partially supported by the Japan Society for the Promotion of Science, Grant-in-Aid for Young Scientists (B) (Subject No. 24740100) and by Keio Gijyuku Academic Development Funds.

3.1.	Local parametrization result	15
3.2.	Local bifurcation from the trivial branch	16
3.3.	Extension to $1 < \gamma < \infty$	16
3.4.	Extension to $0 < \gamma < 1$	19
4.	Entire singular solution	20
4.1.	Existence of the singular solution near $r = 0$	20
4.2.	Positivity of the entire singular solution	26
4.3.	Nonexistence of the entire singular solution	27
4.4.	Existence of critical points of $u^*(s)$	29
4.5.	Proof of Theorem A	30
5.	Convergence to the singular solution as $\gamma \rightarrow \infty$	30
5.1.	Convergence to u^*	30
5.2.	Convergence to λ_n^*	35
6.	Oscillation of $\lambda_n(\gamma)$	36
7.	Boundary concentrating solution	38
7.1.	One-dimensional problem	38
7.2.	Notation	40
7.3.	A priori estimates	41
7.4.	Invertibility of \mathcal{L}	42
7.5.	Contraction mapping	44
7.6.	Nondegeneracy	46
7.7.	Asymptotic behavior of $\lambda_1(\gamma)$ ($\gamma \downarrow 0$)	47
8.	Proof of Theorem B	48
	References	49

1. INTRODUCTION AND MAIN RESULTS

Let $\Omega \subset \mathbb{R}^N$, $N \geq 3$ be a bounded domain with smooth boundary. We are concerned with the elliptic Neumann problem

$$(1.1) \quad \begin{cases} \varepsilon^2 \Delta U - U + U^p = 0, & \text{in } \Omega, \\ \partial_\nu U = 0, & \text{on } \partial\Omega, \\ U > 0, & \text{in } \Omega, \end{cases}$$

where $p > 1$ and $\varepsilon \in \mathbb{R}_+ := \{x; x > 0\}$. The problem (1.1) arises in physical and biological models. In particular, (1.1) appears in the stationary problem of the Keller-Segel model for chemotaxis aggregation [19] and the shadow system of the Gierer-Meinhardt model for

biological pattern formations [12]. For these two decades (1.1) has attracted considerable attention and solutions with various shapes have been found. See [29, 31] for single-peak solutions, [14] for multi-peak solutions, and [21] for boundary concentrating solutions. Many papers, including [14, 21, 27, 29, 31], study the subcritical case $1 < p < p_S$, where

$$p_S := \begin{cases} \frac{N+2}{N-2}, & (N \geq 3), \\ \infty, & (N = 1, 2). \end{cases}$$

In this case the compact embedding $H^1(\Omega) \hookrightarrow L^{p+1}(\Omega)$ is applicable, hence a variational method works well. In this paper we are interested in the solution structure of (1.1) in the supercritical case $p > p_S$. Specifically, we consider the case where the domain Ω is the unit ball $B := \{x \in \mathbb{R}^N; \|x\| < 1\}$, and study the bifurcation diagram $\{(\lambda, U)\}$ of the positive radial solutions. The problem (1.1) can be reduced to the ODE

$$(1.2) \quad \begin{cases} \varepsilon^2 (U_{rr} + \frac{N-1}{r}U_r) - U + U^p = 0, & 0 < r < 1, \\ U_r(1) = 0, \\ U > 0, & 0 \leq r \leq 1, \end{cases}$$

Throughout the present paper we define $f(U) := -U + U^p$ and $\lambda := 1/\varepsilon^2$. Then $\lambda \in \mathbb{R}_+$ and λ diverges as $\varepsilon \downarrow 0$. Since we study the bifurcation diagram of (1.2), it is convenient to transform (1.2) into

$$(1.3) \quad \begin{cases} U_{rr} + \frac{N-1}{r}U_r + \lambda f(U) = 0, & 0 < r < 1, \\ U_r(1) = 0, \\ U > 0, & 0 \leq r \leq 1. \end{cases}$$

We need some notations. We call the constant solutions $\{(\lambda, 1)\}$ the *trivial branch* which is denoted by \mathcal{C}_0 . Let Δ_N denote the Neumann Laplacian, and let $\{\mu_n\}_{n=0}^\infty$ be the eigenvalues of $-\Delta_N$ on B in the space of radial functions. Since each μ_n is simple, $0 = \mu_0 < \mu_1 < \mu_2 < \dots$. Let $\bar{\lambda}_n := \mu_n/(p-1)$. By Crandall-Rabinowitz bifurcation theorem we easily see that $(\bar{\lambda}_n, 1)$ ($n = 1, 2, \dots$) is a bifurcation point from which a curve of nontrivial solutions emanates. We denote the closure of the curve by \mathcal{C}_n . We will see in Proposition 3.1 of the present paper that each curve \mathcal{C}_n , which we call the branch, can be locally parametrized by $\gamma := U_n(0)$. Then each curve \mathcal{C}_n can be described as $\{(\lambda_n(\gamma), U_n(r, \gamma))\}$ ($\gamma := U_n(0, \gamma)$). We define

$$\mathcal{C}_n^+ := \mathcal{C}_n \cap \{\gamma > 1\}, \quad \mathcal{C}_n^- := \mathcal{C}_n \cap \{\gamma < 1\}.$$

By $\mathcal{Z}_I[v(\cdot)]$ we denote the number of the zeros of the function $v(\cdot)$ in the interval $I \subset \mathbb{R}$, i.e.,

$$\mathcal{Z}_I[v(\cdot)] := \#\{x \in I; v(x) = 0\}.$$

Let $(\lambda_n, U_n) \in \mathcal{C}_n$. Every zero of $U_n(r) - 1$ ($0 \leq r \leq 1$) is simple, because of the uniqueness of the solution of the ODE (1.3). If (λ_n, U_n) is near $(\bar{\lambda}_n, 1)$, then $U_n(r) - 1$ is close to the n -th eigenfunction of the corresponding eigenvalue problem. By Sturm-Liouville theory, $\mathcal{Z}_{[0,1]}[U_n(\cdot, \gamma) - 1] = n$ provided that (λ_n, U_n) is near $(\bar{\lambda}_n, 1)$. Since each zero is simple and $U_n(r) - 1$ continuously changes along each \mathcal{C}_n , $\mathcal{Z}_{[0,1]}[U_n(\cdot, \gamma) - 1]$ is preserved along each \mathcal{C}_n . Therefore, $\mathcal{Z}_{[0,1]}[U_n(\cdot, \gamma) - 1] = n$ along \mathcal{C}_n .

Let us recall known results about (1.3). It was shown in [18] that in the subcritical case $1 < p < p_S$ the set of the regular solutions of (1.3) is bounded in L^∞ , i.e.,

$$(\gamma^* :=) \sup\{\|U\|_\infty; U \text{ is a solution of (1.3)}\} < \infty.$$

and that $\lambda_1(\gamma) \rightarrow \infty$ as $\gamma \uparrow \gamma^*$. Since $\partial_\gamma \lambda_1(\gamma)|_{\gamma=1} < 0$ ([24]), the branch \mathcal{C}_1 has at least one turning point. Moreover, if $\gamma (< \gamma^*)$ is close to γ^* , then $U_n(\gamma)$ is nondegenerate, hence $\partial_\gamma \lambda_1(\gamma) > 0$ ([31]). In the critical case $p = p_S$ the solution structure depends on the spatial dimension N . For $N \in \{4, 5, 6\}$, (1.3) admits a (nonconstant) radially decreasing solution for $0 < \lambda < \bar{\lambda}_1$ ([1]), which implies that \mathcal{C}_1 is unbounded in γ . For $N \geq 7$, there is $\underline{\lambda} > 0$ such that (1.3) has no nonconstant solution for $0 < \lambda < \underline{\lambda}$ ([1, 2, 3]). When $N = 3$, the structure depends on the radius of the ball. See [1] for a partial result. In the supercritical case $p > p_S$, there are few results about (1.1). When $p = p_S + \varepsilon$ for small $\varepsilon > 0$, [9] constructed a bubble tower solution of (1.1). See [26, 20, 11] for other results. A brief history of this problem is written in [11].

On the other hand, the branch of the positive solutions of the critical or supercritical Dirichlet problem

$$(1.4) \quad \begin{cases} U_{rr} + \frac{N-1}{r}U_r + \lambda g(U) = 0, & 0 < r < 1, \\ U(1) = 0, \\ U > 0, & 0 \leq r < 1, \end{cases}$$

was studied by [4, 7, 15, 17, 23]. In [17] the case $g(U) = (1 + U)^p$ was studied. The structure depends on p and N . Let

$$(1.5) \quad p_{JL} := \begin{cases} 1 + \frac{4}{N-4-2\sqrt{N-1}}, & (N \geq 11), \\ \infty, & (2 \leq N \leq 10). \end{cases}$$

When $p_S < p < p_{JL}$, Joseph-Lundgren[17] have shown that the branch which emanates from $(0, 0)$ has infinitely many turning points around $\lambda = \lambda^* := \theta(N - 2 - \theta)$ and blows up at $\lambda = \lambda^*$, where

$$(1.6) \quad \theta := \frac{2}{p-1}.$$

Moreover, there is a singular solution $U^* = r^{-\theta} - 1 \in H_0^1(B)$ for $\lambda = \lambda^*$. When $p \geq p_{JL}$, the branch exists for $0 < \lambda < \lambda^*$, does not have a turning point, and blows up at $\lambda = \lambda^*$. Moreover the singular solution (λ^*, U^*) exists. In particular, (1.3) has a unique regular solution for $0 < \lambda < \lambda^*$.

In the study of the bifurcation diagram singular solutions play an important role. First we study the problem

$$(1.7) \quad \begin{cases} u_{ss} + \frac{N-1}{s}u_s + f(u) = 0, & s > 0, \\ u(0) = \gamma, \quad u_s(0) = 0. \end{cases}$$

Theorem 1.1. *Suppose that $p > p_S$. There is a singular positive solution $u^*(s)$ of*

$$\begin{cases} u_{ss} + \frac{N-1}{s}u_s + f(u) = 0, & 0 < s < \infty, \\ u(s) \rightarrow \infty & (s \downarrow 0). \end{cases}$$

$u^*(s)$ oscillates around 1 hence it has infinitely many critical points. Moreover, $u(s, \gamma) \rightarrow u^*(s)$ in $C_{loc}^0(0, \infty)$ as $\gamma \rightarrow \infty$. Here $u(s, \gamma)$ is the solution of (1.7).

When $p > p_{JL}$, this theorem was already proved by [5, Theorem 1.1]. Our method is different from [5] and it can prove the case $p > p_S$.

Using the singular solution u^* , we prove the existence of singular solutions of (1.3). The first main result is the following:

Theorem A. *Suppose that $p > p_S$. The problem (1.3) has infinitely many singular solutions $(\lambda_n^*, U_n^*(r)) \in \mathbb{R}_+ \times (C^2(0, 1) \cap C^0(0, 1])$ ($n = 1, 2, \dots$ and $\lambda_1^* < \lambda_2^* < \dots \rightarrow \infty$) such that the following assertions hold:*

(i) $U_n^*(r)$ satisfies

$$(1.8) \quad U_n^*(r) = A(p, N)(\sqrt{\lambda_n^* r})^{-\theta}(1 + o(1)) \quad \text{as } (r \downarrow 0),$$

where

$$(1.9) \quad A(p, N) := \{\theta(N - 2 - \theta)\}^{\frac{1}{p-1}}.$$

(ii) $\mathcal{Z}_{(0,1]}[U_n^*(\cdot) - 1] = n$.

(iii) $U_n^*(r) > 0$ ($0 < r \leq 1$).

Moreover, the singular solution (λ_n^*, U_n^*) is unique, i.e., if $(\tilde{\lambda}_n^*, \tilde{U}_n^*)$ is a singular solution such that (i) and (ii) hold, then $(\tilde{\lambda}_n^*, \tilde{U}_n^*) = (\lambda_n^*, U_n^*)$.

The second main result of the paper is the following:

Theorem B. *Suppose that $p > p_S$. Let λ_n^* ($n = 1, 2, \dots$) be as in Theorem A. Let \mathcal{S} denote the set of the regular solutions of (1.3). Then*

$$(1.10) \quad \mathcal{S} = \mathcal{C}_0 \cup \bigcup_{n=1}^{\infty} (\mathcal{C}_n^+ \cup \mathcal{C}_n^-).$$

Each \mathcal{C}_n ($n = 1, 2, \dots$) can be parametrized by $\gamma = U_n(0)$, hence \mathcal{C}_n can be described as $\{(\lambda_n(\gamma), U_n(r, \gamma))\}$. Moreover, $\lambda_n(\gamma) \in C^1(0, \infty)$ and the following assertions hold:

(i) For each $n \geq 1$, $\lambda_n(1) = \bar{\lambda}_n$.

(ii) For each $n \geq 1$, $\lambda_n(\gamma) \rightarrow \lambda_n^*$ ($\gamma \rightarrow \infty$).

(iii) If $p_S < p < p_{JL}$, then for each $n \geq 1$, $\lambda_n(\gamma)$ oscillates around λ_n^* infinitely many times as $\gamma \rightarrow \infty$.

(iv) For each $n \geq 1$, $\lambda_n(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$).

(v) If $\gamma > 0$ is small, then $U_1(r, \gamma)$ is nondegenerate, and it becomes a boundary concentrating solution as $\gamma \downarrow 0$.

(vi) For each $\gamma \in \mathbb{R}_+$, $\lambda_1(\gamma) < \lambda_2(\gamma) < \dots$.

See Figure 1 for the case $p_S < p < p_{JL}$.

(iv), (v), and (vi) of Theorem B hold for every $p > 1$.

It follows from Theorem B (iii) that each branch \mathcal{C}_n^+ has infinitely many turning points around λ_n^* if $p_S < p < p_{JL}$.

Remark 1.2. *If $3 \leq N \leq 10$, then $p_{JL} = \infty$. Hence, the supercritical equation (1.3) always has infinitely many solutions for $\lambda = \lambda_n^*$ ($n = 1, 2, \dots$) and each \mathcal{C}_n^+ has infinitely many turning points.*




FIGURE 1. The schematic picture of the bifurcation diagram of (1.3) in the case $p_S < p < p_{JL}$.

Corollary 1.3. *Suppose that $p > p_S$. There exists $\underline{\lambda} > 0$ such that if $0 < \lambda < \underline{\lambda}$, then (1.3) has no nonconstant solution. Moreover, there is $\bar{\lambda} > 0$ such that the radially decreasing solution does not exist for $\lambda > \bar{\lambda}$.*

Corollary 1.4. *Suppose that $p_S < p < p_{JL}$. There exists $\underline{\lambda} > 0$ such that (1.3) has a nonconstant solution if $\lambda \geq \underline{\lambda}$ and that (1.3) has no nonconstant solution if $0 < \lambda < \underline{\lambda}$.*

Let us explain technical details. In [17] they used a special change of variables in the study of (1.4) with $g(U) = (1 + U)^p$. The equation (1.4) can be transformed to a first order autonomous system. See (2.5) in the present paper. They used a phase plane analysis to obtain the bifurcation diagram. However, for a general nonlinearity g , we cannot expect such a change of variables. In [23] the author studied the bifurcation diagram of the solutions of (1.4) when $g(U) = U^p + h(U) (> 0)$ and $h(U)$ is a lower order term. Let $u(s) := U(r)$ and $s := \sqrt{\lambda}r$. The equation (1.4) becomes

$$(1.11) \quad \begin{cases} u_{ss} + \frac{N-1}{s}u_s + g(u) = 0, & 0 < s < \sqrt{\lambda}, \\ u(\sqrt{\lambda}) = 0, \\ u > 0, & 0 \leq s < \sqrt{\lambda}. \end{cases}$$

The equation (1.11) has a singular solution $(\lambda^*, u^*(s))$. Let $(\lambda, u(s, \gamma))$ denote the regular solution of (1.11) such that $u(0, \gamma) = \gamma$ and $u_s(0, \gamma) =$

0. It was shown that λ is a C^1 -function of γ . He used the intersection number between $u^*(s)$ and $u(s, \gamma)$ which can be written as

$$(1.12) \quad \mathcal{Z}_{I_\gamma}[u^*(\cdot) - u(\cdot, \gamma)],$$

where $I_\gamma := (0, \min\{\sqrt{\lambda^*}, \sqrt{\lambda(\gamma)}\}]$. He showed by a scaling argument that if $p_S < p < p_{JL}$, then

$$(1.13) \quad \mathcal{Z}_{I_\gamma}[u^*(\cdot) - u(\cdot, \gamma)] \rightarrow \infty \quad (\gamma \rightarrow \infty).$$

Because of the uniqueness of the solution of the ODE (1.11), each zero of $u^*(\cdot) - u(\cdot, \gamma)$ is simple. Hence, each zero depends continuously on γ . The number of the zeros in I_γ is preserved if another zero does not come from the boundary of I_γ . Since $u^*(0) - u(0, \gamma) = \infty$, a zero cannot enter I_γ from $s = 0$. Because of (1.13), a zero enters I_γ from $s = \min\{\sqrt{\lambda^*}, \sqrt{\lambda(\gamma)}\}$ infinitely many times as $\gamma \rightarrow \infty$. Therefore, $\lambda(\gamma)$ oscillates around λ^* infinitely many times which indicates that the branch has infinitely many turning points. This method cannot be directly applied to the Neumann problem, since in the Neumann case $u(\sqrt{\lambda(\gamma)}, \gamma)$ is not necessarily equal to $u^*(\sqrt{\lambda^*})$ and the oscillation of $\lambda(\gamma)$ around λ^* is not trivial. The shape of the singular solution u^* of the Neumann problem becomes important to study the behavior of each zero.

In our problem (1.3) we see that \mathcal{C}_n is locally parametrized by $\gamma = U_n(0)$ (Proposition 3.1). We can write the solution as $(\lambda_n(\gamma), U_n(r, \gamma))$.

Section 3 is devoted to the study of the fundamental property of $\lambda_n(\gamma)$. In Lemma 3.3 we show that for each $\gamma^* > 1$, $\lambda_n(\gamma)$ does not diverge as $\gamma \uparrow \gamma^*$. Therefore, combining this result and the local parametrization of \mathcal{C}_n , we show in Lemma 3.4 that the domain of $\lambda_n(\gamma)$ can be extended to $\gamma > 1$. It is perhaps interesting to note that the nondivergence of $\lambda_n(\gamma)$ is proved by the nonexistence of the entire positive solution of $\Delta u - u + u^p = 0$ ($p \geq p_S$) which is proved by the Pohozaev identity. We also show that $\lambda_n(\gamma)$ can be extended to $0 < \lambda < 1$. Hence, $\lambda_n(\gamma)$ is defined in $\gamma \in \mathbb{R}_+$.

In Section 4 we study the singular solution of (1.3) and prove Theorem A. Let $u(s) := U(r)$ and $s := \sqrt{\lambda}r$. The equation (1.3) is transformed to the problem

$$(1.14) \quad \begin{cases} u_{ss} + \frac{N-1}{s}u_s + f(u) = 0, & 0 < s < \sqrt{\lambda}, \\ u_s(\sqrt{\lambda}) = 0, \\ u > 0, \end{cases} \quad 0 \leq s \leq \sqrt{\lambda}.$$

In Lemma 4.1 we construct the singular solution $u^*(s)$ of the equation in (1.14) near $s = 0$ and show that $u^*(s) = As^{-\theta}(1 + o(1))$ ($s \downarrow 0$). In Lemma 4.4 we show that the domain of $u^*(s)$ can be extended to $0 < s < \infty$, that $u^*(s)$ satisfies the equation in (1.14), and that $u^*(s) > 0$ for $s > 0$. In Lemma 4.8 we show that $u^*(s)$ oscillates around 1 infinitely many times as $s \rightarrow \infty$ and that $u^*(s)$ has the set of the critical points $\{s_n^*\}_{n=1}^\infty$ of u^* such that $0 < s_1^* < s_2^* < \dots \rightarrow \infty$ and

$$\begin{cases} s_n^* \text{ is a local minimum point of } u^* \text{ and } u^*(s_n^*) < 1 \text{ if } n \in \{1, 3, 5, \dots\}, \\ s_n^* \text{ is a local maximum point of } u^* \text{ and } u^*(s_n^*) > 1 \text{ if } n \in \{2, 4, 6, \dots\}. \end{cases}$$

We set $\lambda_n^* := (s_n^*)^2$ and $U_n^*(r) := u^*(s)$ ($s = \sqrt{\lambda_n^*}r$). Then, (λ_n^*, U_n^*) is a singular solution stated in Theorem A.

Let $(\lambda_n(\gamma), u(s, \gamma))$ denote the solution of (1.14) such that $u(0, \gamma) = \gamma$ and $u_s(0, \gamma) = 0$. In Section 5 we show that $\lambda_n(\gamma) \rightarrow \lambda_n^*$ as $\gamma \rightarrow \infty$ and that $u(s, \gamma)$ converges to $u^*(s)$ in an appropriate sense. In [22] the authors proved a similar convergence result for the Dirichlet problem

$$\begin{cases} U_{rr} + \frac{N-1}{r}U_r + \lambda U + U^p = 0, & 0 < r < 1, \\ U(1) = 0, \\ U > 0, \end{cases} \quad 0 \leq r < 1.$$

when $p > p_S$. In Theorem 5.1 we show that $u(s, \gamma) \rightarrow u^*(s)$, following arguments in the proof of [22, Theorem A].

In Section 6 we show that $\lambda_n(\gamma)$ oscillates around λ_n^* if $p_S < p < p_{JL}$. Let $\rho := \gamma^{\frac{p-1}{2}}s$. We define $\tilde{u}(\rho, \gamma) := u(s, \gamma)/\gamma$ and $\tilde{u}^*(\rho) := u^*(s)/\gamma$. We use the intersection number between \tilde{u} and \tilde{u}^* . The function $\tilde{u}(\rho, \gamma)$ satisfies

$$(1.15) \quad \begin{cases} \tilde{u}_{\rho\rho} + \frac{N-1}{\rho}\tilde{u}_\rho + \tilde{u}^p - \frac{1}{\gamma^{p-1}}\tilde{u} = 0, & 0 < \rho < \infty, \\ \tilde{u}(0) = 1, \quad \tilde{u}_\rho(0) = 0. \end{cases}$$

Let $\bar{u}(\rho, \gamma)$ be the regular solution of

$$(1.16) \quad \begin{cases} \bar{u}_{\rho\rho} + \frac{N-1}{\rho}\bar{u}_\rho + \bar{u}^p = 0, & 0 < \rho < \infty, \\ \bar{u}(0) = \gamma, \quad \bar{u}_\rho(0) = 0. \end{cases}$$

We show that as $\gamma \rightarrow \infty$,

$$\tilde{u}(\rho, \gamma) \rightarrow \bar{u}(\rho, 1) \quad \text{in} \quad C_{loc}^2(0, \infty) \cap C_{loc}^0[0, \infty)$$

and

$$\tilde{u}^*(\rho) \rightarrow \bar{u}^*(\rho) \quad \text{in} \quad C_{loc}^0(0, \infty),$$

where $\bar{u}^*(\rho)$ is given by (2.2) which is a singular solution of the equation in (1.16). In Section 2 we recall the fact that $\mathcal{Z}_{(0, \infty)}[\bar{u}^*(\cdot) - \bar{u}(\cdot, 1)] = \infty$. Hence, for each $\delta > 0$,

$$(1.17) \quad \mathcal{Z}_{(0, \delta)}[u^*(\cdot) - u(\cdot, \gamma)] \rightarrow \infty \quad (\gamma \rightarrow \infty),$$

since $s \in (0, \delta)$ is corresponding to $\rho \in (0, \delta\gamma^{\frac{N-1}{2}})$ and $\delta\gamma^{\frac{N-1}{2}} \rightarrow \infty$ ($\gamma \rightarrow \infty$). Since each zero of $u^*(\cdot) - u(\cdot, \gamma)$ is simple, each zero depends continuously on γ . The divergence (1.17) tells us that a zero which is simple enters the interval $(0, \sqrt{\lambda_n^*}]$ from $s = \sqrt{\lambda_n^*}$ infinitely many times. Therefore, there exists a sequence of large numbers $\{\gamma_j\}_{j=1}^\infty$ ($\gamma_1 < \gamma_2 < \dots \rightarrow \infty$) such that $u_s(\sqrt{\lambda_n^*}, \gamma_j) < 0$ for $j \in \{1, 3, 5, \dots\}$ and $u_s(\sqrt{\lambda_n^*}, \gamma_j) > 0$ for $j \in \{2, 4, 6, \dots\}$. In Theorem 6.1, using the convergence $u(s, \gamma) \rightarrow u^*(s)$, we show that if $n \in \{1, 3, 5, \dots\}$ (resp. $n \in \{2, 4, 6, \dots\}$)

$$(1.18) \quad \lambda_n(\gamma_j) \begin{cases} > \lambda_n^*, & (j \in \{1, 3, 5, \dots\}), \\ < \lambda_n^*, & (j \in \{2, 4, 6, \dots\}), \end{cases}$$

$$\left(\text{resp. } \lambda_n(\gamma_j) \begin{cases} < \lambda_n^*, & (j \in \{1, 3, 5, \dots\}), \\ > \lambda_n^*, & (j \in \{2, 4, 6, \dots\}). \end{cases} \right)$$

which implies that $\lambda_n(\gamma)$ oscillates around λ_n^* infinitely many times as $\gamma \rightarrow \infty$.

In Section 7 we construct a smooth branch of boundary concentrating solutions, using a standard blow-up argument with the contraction mapping theorem. We also show that this branch is in \mathcal{C}_1 and that $\lambda_1(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$).

The paper consists of 8 sections. In Section 2 we recall known result about the intersection number and the nonexistence of the entire solution. In Section 3 we collect fundamental properties of $\lambda_n(\gamma)$. In Section 4 we prove Theorem A. In Section 5 we prove the convergence

to (λ_n^*, U_n^*) . In Section 6 we prove the oscillation of $\lambda_n(\gamma)$. In Section 7 we prove (iv) and (v) of Theorem B. In Section 8 we prove the other assertions of Theorem B and Corollaries 1.3 and 1.4.

2. PRELIMINARIES

2.1. Regular and singular solutions of the limit equation. Let us consider the radial solution $\bar{u}(\rho)$ of the elliptic equation on \mathbb{R}^N

$$(2.1) \quad \begin{cases} \Delta \bar{u} + \bar{u}^p = 0 & \text{in } \mathbb{R}^N, \\ \bar{u} > 0 & \text{in } \mathbb{R}^N. \end{cases}$$

It is well known that

$$(2.2) \quad \bar{u}^*(\rho) := A\rho^{-\theta}$$

is a singular solution of (2.1), where θ is defined in (1.6) and $A := A(p, N)$ is defined in (1.9). Next, we consider the regular radial solution $\bar{u}(\rho)$. The solution $\bar{u}(\rho)$ satisfies (1.16). We define $\bar{v}(\rho) := A^{-1}\rho^\theta \bar{u}(\rho)$. The function $\bar{v}(\rho)$ satisfies

$$\begin{cases} \bar{v}_{\rho\rho} + \frac{1}{\rho}(N-1-2\theta)\bar{v}_\rho + \frac{A^{p-1}}{\rho^2}(\bar{v}^p - \bar{v}) = 0, & 0 < \rho < \infty, \\ r^{-\theta}\bar{v}(\rho) \rightarrow \frac{\gamma}{A} \quad (\rho \downarrow 0), \\ \bar{v} > 0, & 0 \leq \rho < \infty. \end{cases}$$

In order to make the equation autonomous we change variables to $t := A^{\frac{p-1}{2}} \log \rho$ and $y(t) := \bar{v}(\rho)$. The function $y(t)$ satisfies

$$(2.3) \quad \begin{cases} y'' + \alpha y' - y + y^p = 0, & -\infty < t < \infty, \\ e^{-\theta mt} y(t) \rightarrow \frac{\gamma}{A} \quad (t \rightarrow -\infty), \\ y > 0, & -\infty < t < \infty, \end{cases}$$

where the prime stands for the derivative,

$$(2.4) \quad m := A^{-\frac{p-1}{2}}, \quad \text{and} \quad \alpha := m(N-2-2\theta).$$

The equation (2.3) is transformed into the first order autonomous system

$$(2.5) \quad \begin{cases} y' = z \\ z' = -\alpha z + y - y^p. \end{cases}$$

This system has two equilibrium points $(0, 0)$ and $(1, 0)$. By simple calculation we see that $(0, 0)$ is a saddle point. The eigenvalues of the linearized operator at $(0, 0)$ are the roots of $\Lambda^2 + \alpha\Lambda - 1 = 0$. The eigenvalues are $m\theta (> 0)$ and $-m(N-2-\theta) (< 0)$. On the other

hand, the eigenvalues of the linearized operator at $(1, 0)$ are the roots of $\Lambda^2 + \alpha\Lambda + p - 1 = 0$. Therefore, $(1, 0)$ is a spiral point if and only if

$$(2.6) \quad \alpha^2 - 4(p - 1) < 0, \text{ i.e., } (N - 2 - 2\theta)^2 - 8(N - 2 - \theta) < 0.$$

Then,

$$(2.7) \quad (N - 4 - 2\sqrt{N - 1})/2 < \theta < (N - 4 + 2\sqrt{N - 1})/2.$$

We easily see that $1 + 4/(N - 4 + 2\sqrt{N - 1}) < p_S (< p)$. Thus, $\theta < (N - 4 + 2\sqrt{N - 1})/2$ always holds. If $N \leq 10$, then $(N - 4 - 2\sqrt{N - 1})/2 \leq 0 (< \theta)$, hence (2.7) holds. When $N \geq 11$, (2.7) holds if

$$p < 1 + \frac{4}{N - 4 - 2\sqrt{N - 1}} (= p_{JL}).$$

If $p \geq p_{JL}$, then two eigenvalues are negative. Thus, we have the following:

$$\begin{cases} \text{If } N \leq 10, \text{ then } (1, 0) \text{ is a spiral point for } p > p_S. \\ \text{If } N \geq 11, \text{ then } (1, 0) \text{ is a } \begin{cases} \text{spiral point for } p_S < p < p_{JL}. \\ \text{(resp. degenerate) node for} \\ \text{(resp. } p = p_{JL}) p > p_{JL}. \end{cases} \end{cases}$$

Let

$$(2.8) \quad E(y, z) := \frac{1}{2}z^2 - \frac{1}{2}y^2 + \frac{1}{p+1}y^{p+1}.$$

By direct calculation we have $\frac{d}{dt}E(y(t), z(t)) = -\alpha z^2(t)$. If $p > p_S$, then $\alpha > 0$ and E is a Lyapunov function. A phase plane analysis with this Lyapunov function reveals that there is a heteroclinic orbit from $(0, 0)$ to $(1, 0)$ which is in the right half-plane. Note that the eigenvalue θm is compatible with the limit condition in (2.3), because $y(t) \sim C_0 e^{\theta m t}$ ($t \rightarrow -\infty$). This orbit corresponds to a regular solution $\bar{u}(\rho)$ of (1.16) for some γ , and \bar{u} is positive. Since $\bar{u}(\rho)$ is a solution, $\gamma \bar{u}(\gamma^{\frac{p-1}{2}} \rho)$ is also a solution. Without loss of generality, we assume that $\bar{u}(0) = 1$. Every solution of (1.16) can be written as $\gamma \bar{u}(\gamma^{\frac{p-1}{2}} \rho)$. Since the equilibrium point $(1, 0)$ corresponds to the singular solution $\bar{u}^*(\rho)$ of (1.16) and $(1, 0)$ is a spiral point for $p_S < p < p_{JL}$, the intersection number of $\bar{u}(\rho, \gamma)$ and $\bar{u}^*(\rho)$ is infinite for each γ , i.e., $\mathcal{Z}_{(0, \infty)}[\bar{u}^*(\cdot) - \bar{u}(\cdot, \gamma)] = \infty$. We have

Proposition 2.1. *Suppose that $p > p_S$. Then the problem (1.16) has a unique (positive) solution $\bar{u}(\rho, \gamma)$ and $\bar{u}(\rho, \gamma) = \gamma \bar{u}(\gamma^{\frac{p-1}{2}} \rho, 1)$.*

Moreover, if $p_S < p < p_{JL}$, then, for each $\gamma > 0$, $\mathcal{Z}_{(0,\infty)}[\bar{u}^*(\cdot) - \bar{u}(\cdot, \gamma)] = \infty$.

In Section 4 we use Proposition 2.1 in order to prove (1.17).

2.2. Nonexistence of the entire solution.

Proposition 2.2. *If $u(s) \in C^2(\varepsilon, R) \cap C^0[\varepsilon, R]$ satisfies*

$$(2.9) \quad u_{ss} + \frac{N-1}{s}u_s + f(u) = 0 \quad (\varepsilon < s < R)$$

and $u(s) > 0$ ($\varepsilon < s < R$), then

$$(2.10) \quad \begin{aligned} 0 &= \left(\frac{N}{2} - 1 - \frac{N}{p+1} \right) \int_{\varepsilon}^R s^{N-1} u_s^2 ds + \left(\frac{N}{2} - \frac{N}{p+1} \right) \int_{\varepsilon}^R s^{N-1} u^2 ds \\ &+ \frac{N}{p+1} (R^{N-1} u(R) u_s(R) - \varepsilon^{N-1} u(\varepsilon) u_s(\varepsilon)) + \frac{1}{2} (R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2) \\ &- \frac{1}{2} (R^N u(R)^2 - \varepsilon^N u(\varepsilon)^2) + \frac{1}{p+1} (R^N u(R)^{p+1} - \varepsilon^N u(\varepsilon)^{p+1}). \end{aligned}$$

Proof of Proposition 2.2. Multiplying (2.9) by $s^N u_s$ and integrating it over $[\varepsilon, R]$, we have

$$\begin{aligned} 0 &= \int_{\varepsilon}^R s^N u_s \left\{ \frac{1}{s^{N-1}} (s^{N-1} u_s)_s - u + u^p \right\} ds \\ &= \int_{\varepsilon}^R (s^{N-1} u_s)_s s u_s ds - \int_{\varepsilon}^R s^N u u_s ds + \int_{\varepsilon}^R s^N u^p u_s ds \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

Then,

$$\begin{aligned} I_1 &= [s^{N-1} u_s s u_s]_{\varepsilon}^R - \int_{\varepsilon}^R s^{N-1} u_s (s u_{ss} + u_s) ds \\ &= R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2 - \int_{\varepsilon}^R s^N u_s u_{ss} ds - \int_{\varepsilon}^R s^{N-1} u_s^2 ds. \end{aligned}$$

Since

$$\begin{aligned} \int_{\varepsilon}^R s^N u_s u_{ss} ds &= \left[s^N \frac{u_s^2}{2} \right]_{\varepsilon}^R - \int_{\varepsilon}^R N s^{N-1} \frac{u_s^2}{2} ds \\ &= \frac{1}{2} (R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2) - \frac{N}{2} \int_{\varepsilon}^R s^{N-1} u_s^2 ds, \end{aligned}$$

we have

$$\begin{aligned}
I_1 &= R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2 - \int_{\varepsilon}^R s^{N-1} u_s^2 ds \\
&\quad - \frac{1}{2} (R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2) + \frac{N}{2} \int_{\varepsilon}^R s^{N-1} u_s^2 ds \\
&= \left(\frac{N}{2} - 1 \right) \int_{\varepsilon}^R s^{N-1} u_s^2 ds + \frac{1}{2} (R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2).
\end{aligned}$$

Next, we have

$$\begin{aligned}
I_2 &= - \left[s^N \frac{u^2}{2} \right]_{\varepsilon}^R + \int_{\varepsilon}^R N s^{N-1} \frac{u^2}{2} ds \\
&= \frac{N}{2} \int_{\varepsilon}^R s^{N-1} u^2 ds - \frac{1}{2} (R^N u(R)^2 - \varepsilon^N u(\varepsilon)^2), \\
I_3 &= \left[s^N \frac{u^{p+1}}{p+1} \right]_{\varepsilon}^R - \int_{\varepsilon}^R N s^{N-1} \frac{u^{p+1}}{p+1} ds \\
&= - \frac{N}{p+1} \int_{\varepsilon}^R s^{N-1} u^{p+1} ds + \frac{1}{p+1} (R^N u(R)^{p+1} - \varepsilon^N u(\varepsilon)^{p+1}).
\end{aligned}$$

Adding I_1 , I_2 , and I_3 , we have

$$\begin{aligned}
(2.11) \quad 0 &= I_1 + I_2 + I_3 \\
&= \left(\frac{N}{2} - 1 \right) \int_{\varepsilon}^R s^{N-1} u_s^2 ds + \frac{N}{2} \int_{\varepsilon}^R s^{N-1} u^2 ds \\
&\quad - \frac{N}{p+1} \int_{\varepsilon}^R s^{N-1} u^{p+1} ds \\
&\quad + \frac{1}{2} (R^N u_s(R)^2 - \varepsilon^N u_s(\varepsilon)^2) - \frac{1}{2} (R^N u(R)^2 - \varepsilon^N u(\varepsilon)^2) \\
&\quad + \frac{1}{p+1} (R^N u(R)^{p+1} - \varepsilon^N u(\varepsilon)^{p+1}).
\end{aligned}$$

Multiplying (2.9) by $s^{N-1}u$ and integrating it over $[\varepsilon, R]$, we have

$$\begin{aligned}
0 &= \int_{\varepsilon}^R s^{N-1} u \left\{ \frac{1}{s^{N-1}} (s^{N-1} u_s)_s - u + u^p \right\} ds \\
&= \int_{\varepsilon}^R (s^{N-1} u_s)_s u ds - \int_{\varepsilon}^R s^{N-1} u^2 ds + \int_{\varepsilon}^R s^{N-1} u^{p+1} ds.
\end{aligned}$$

Since

$$\begin{aligned}
\int_{\varepsilon}^R (s^{N-1} u_s)_s u ds &= [s^{N-1} u_s u]_{\varepsilon}^R - \int_{\varepsilon}^R s^{N-1} u_s^2 ds \\
&= - \int_{\varepsilon}^R s^{N-1} u_s^2 ds + R^{N-1} u(R) u_s(R) - \varepsilon^{N-1} u(\varepsilon) u_s(\varepsilon),
\end{aligned}$$

we have

$$(2.12) \quad 0 = - \int_{\varepsilon}^R s^{N-1} u_s^2 ds - \int_{\varepsilon}^R s^{N-1} u^2 ds + \int_{\varepsilon}^R s^{N-1} u^{p+1} ds \\ + R^{N-1} u(R) u_s(R) - \varepsilon^{N-1} u(\varepsilon) u_s(\varepsilon).$$

Calculating (2.11) + $\frac{N}{p+1}$ (2.12), we have (2.10). \square

Proposition 2.3. *Suppose that $p \geq p_S$. The problem*

$$(2.13) \quad \begin{cases} u_{ss} + \frac{N-1}{s} u_s + f(u) = 0, & 0 < s < \infty, \\ u_s(0) = 0, \quad u(0) < \infty, \quad u(\infty) = 0, \\ u > 0, & 0 \leq s < \infty \end{cases}$$

does not admit a solution.

Proof of Proposition 2.3. Since $f'(0) = -1 < 0$, it is well known that $u(s)$ and $u_s(s)$ decay exponentially as $s \rightarrow \infty$. Thus, the four terms $R^{N-1} u(R) u_s(R)$, $R^N u_s(R)^2$, $R^N u(R)^2$, and $R^N u(R)^{p+1}$ converge to zero as $R \rightarrow \infty$. It is clear that the four terms $\varepsilon^{N-1} u(\varepsilon) u_s(\varepsilon)$, $\varepsilon^N u_s(\varepsilon)^2$, $\varepsilon^N u(\varepsilon)$, and $\varepsilon^N u(\varepsilon)^{p+1}$ converge to zero as $\varepsilon \downarrow 0$. Taking the limit of (2.10) as $\varepsilon \downarrow 0$ and $R \rightarrow \infty$, we have

$$\left(\frac{N}{2} - 1 - \frac{N}{p+1} \right) \int_0^\infty s^{N-1} u_s^2 ds + \left(\frac{N}{2} - \frac{N}{p+1} \right) \int_0^\infty s^{N-1} u^2 ds = 0.$$

Since $p \geq p_S$, $\frac{N}{2} - 1 - \frac{N}{p+1} \geq 0$. The above equality implies that (2.13) does not admit a solution. \square

3. LOCAL AND GLOBAL PARAMETRIZATION RESULTS

In this section we study the parametrization of each branch \mathcal{C}_n of the solution of (1.3). Let $U(r)$ be a regular solution of (1.3). We sometimes use the stretched variable $s := \sqrt{\lambda} r (= r/\varepsilon)$. We define $u(s) := U(r)$. Then $u(s)$ satisfies (1.14). We denote the solution of the initial value problem (1.7) by $u(s, \gamma)$.

3.1. Local parametrization result. To begin with, we study the local parametrization of the branch.

Proposition 3.1 ([24, Proposition 3.1]). *Let (λ_0, U_0) be a nonconstant regular solution of (1.3), and let $\gamma_0 := U_0(0)$. Then, all solutions near (λ_0, U_0) can be parametrized as $\{(\lambda(\gamma), U(r, \gamma))\}_{|\gamma - \gamma_0| < \varepsilon}$ ($U(0, \gamma) = \gamma$, $\lambda(\gamma_0) = \lambda_0$, $U(r, \gamma_0) = U_0(r)$).*

This proposition was obtained in [24]. We prove this proposition for readers' convenience.

Proof of Proposition 3.1. Let $s := \sqrt{\lambda}r$ and $u(s) := U(r)$. Then, u is not constant and u satisfies (1.14). We consider the initial value problem (1.7) and denote the solution by $u(s, \gamma)$. Then, u is a C^2 -function of (s, γ) and $U(r, \lambda, \gamma) (= u(\sqrt{\lambda}r, \gamma))$ satisfies (1.3). Since $U_r(1, \lambda, \gamma) = 0$, $\sqrt{\lambda}u_s(\sqrt{\lambda}, \gamma) = 0$. We will show that $\partial_\lambda u_s(\sqrt{\lambda}, \gamma) \neq 0$. Since $\partial_\lambda u_s(\sqrt{\lambda}, \gamma) = u_{ss}(\sqrt{\lambda}, \gamma)/(2\sqrt{\lambda})$, it is enough to show that $u_{ss}(\sqrt{\lambda}, \gamma) \neq 0$. Suppose the contrary, i.e., $u_{ss}(\sqrt{\lambda}, \gamma) = 0$. Differentiating (1.7) with respect to s , we have

$$u_{sss} + \frac{N-1}{s}u_{ss} + \left(f'(u) - \frac{N-1}{s^2} \right) u_s = 0.$$

Since $u_s(\sqrt{\lambda}, \gamma) = u_{ss}(\sqrt{\lambda}, \gamma) = 0$, we see by the uniqueness of the solution of the ODE that $u_s(s, \gamma) \equiv 0$. Thus, u is a constant solution of (1.7) which contradicts that U is not constant. Hence, $u_{ss}(\sqrt{\lambda}, \gamma) \neq 0$. We can apply the implicit function theorem to $u_s(\sqrt{\lambda}, \gamma) = 0$. We see that there is a C^1 -function $\lambda = \lambda(\gamma)$, which is defined in a neighborhood of γ_0 , such that $u_s(\sqrt{\lambda(\gamma)}, \gamma) = 0$ and that all the solutions near (λ_0, U_0) can be written as $(\lambda(\gamma), u(\sqrt{\lambda(\gamma)}r, \gamma))$ ($|\gamma - \gamma_0| < \varepsilon$). The positivity $u(\sqrt{\lambda(\gamma)}r, \gamma) > 0$ ($0 \leq r \leq 1$) follows from the maximum principle. \square

Remark 3.2. *It follows from Proposition 3.1 that another radial branch does not emanate from \mathcal{C}_n . However, the branch may have a turning point.*

3.2. Local bifurcation from the trivial branch. We work on the space of radial functions. Let ϕ_n be the eigenfunction of the linearization of (1.3) at $(\bar{\lambda}_n, 1)$. Since

$$\int_0^1 \phi_n \partial_{U\lambda}(\lambda f(U))|_{(\lambda, U)=(\bar{\lambda}_n, 1)} \phi_n r^{N-1} dr = (p-1) \int_0^1 \phi_n^2 r^{N-1} dr \neq 0,$$

the transversality condition of Crandall-Rabinowitz type holds, hence $(\bar{\lambda}_n, 1)$ is a bifurcation point.

3.3. Extension to $1 < \gamma < \infty$. For each $n \geq 1$, the branch \mathcal{C}_n emanates from $(\bar{\lambda}_n, 1)$ and it is locally parametrized by $\gamma = U_n(0)$, hence

$\mathcal{C}_n = \{(\lambda_n(\gamma), U_n(r, \gamma))\}$ ($\gamma = U_n(0, \gamma)$). In this subsection we extend the branch in the direction $\gamma \rightarrow \infty$. In particular, we show that, for each $\gamma^* > 1$, $\lambda_n(\gamma)$ does not diverge as $\gamma \uparrow \gamma^*$.

Lemma 3.3. *For each $\gamma^* > 1$, there are $C > 0$ and $\varepsilon > 0$ such that $|\lambda_n(\gamma)| < C$ for $\gamma \in (\gamma^* - \varepsilon, \gamma^*)$.*

Proof. First, we prove the case $n = 1$. We use a contradiction argument. Suppose the contrary, i.e.,

$$(3.1) \quad \begin{aligned} \text{there is a sequence } \{\gamma_j\}_{j=1}^{\infty} \text{ (} \gamma_j < \gamma^*, \gamma_j \uparrow \gamma^* \text{)} \\ \text{such that } \lambda_1(\gamma_j) \rightarrow \infty \text{ (} j \rightarrow \infty \text{)}. \end{aligned}$$

Let $u(s, \gamma_j)$ denote the solution of (1.7), and let $u_j(s) := u(s, \gamma_j)$. Then, $u_j(0) = \gamma_j$. We show that there exists a solution $u_*(s) \in C^2(0, \infty) \cap C^0[0, \infty)$ of the problem

$$(3.2) \quad \begin{cases} u_{ss} + \frac{N-1}{s}u_s + f(u) = 0, & 0 < s < \infty, \\ u_s(0) = 0, \quad u(0) = \gamma^*, \quad u(\infty) = 0, \\ u > 0, \quad u_s < 0, & 0 < s < \infty. \end{cases}$$

Since the problem (1.7) is well-posed, there exists $\tilde{u}_*(s) \in C^2(0, \infty) \cap C^0[0, \infty)$ such that \tilde{u}_* satisfies

$$\begin{cases} (\tilde{u}_*)_{ss} + \frac{N-1}{s}(\tilde{u}_*)_s + f(\tilde{u}_*) = 0, & 0 < s < \infty, \\ (\tilde{u}_*)_s(0) = 0, \quad \tilde{u}_*(0) = \gamma^*, \end{cases}$$

and $u_j(s) \rightarrow \tilde{u}_*(s)$ in $C_{loc}^2(0, \infty) \cap C_{loc}^0[0, \infty)$ as $j \rightarrow \infty$. Therefore, in order to show that $\tilde{u}_*(s) \equiv u_*(s)$ we show that \tilde{u}_* satisfies

$$(3.3) \quad \begin{cases} \tilde{u}_*(\infty) = 0, \\ \tilde{u}_* > 0, \quad (\tilde{u}_*)_s < 0, \quad 0 < s < \infty. \end{cases}$$

If there is $\tilde{s}_0 > 0$ such that $\tilde{u}_*(\tilde{s}_0) < 0$, then, for large $j \geq 1$, $u_j(\tilde{s}_0) < 0$, since $u_j \rightarrow \tilde{u}_*$ in $C_{loc}^0[0, \infty)$. We obtain a contradiction, because $\tilde{s}_0 < \sqrt{\lambda_1(\gamma_j)}$ for large $j \geq 1$ and $u_j(s) > 0$ ($0 < s < \sqrt{\lambda_1(\gamma_j)}$) for every $j \geq 1$. Thus $\tilde{u}_*(s) \geq 0$ ($0 \leq s < \infty$). By the maximum principle we see that $\tilde{u}_*(s) > 0$ ($0 \leq s < \infty$). If there is $\tilde{s}_1 > 0$ such that $\partial_s \tilde{u}_*(\tilde{s}_1) > 0$, then, for large $j \geq 1$, $\partial_s u_j(\tilde{s}_1) > 0$. We obtain a contradiction, because $\tilde{s}_1 < \sqrt{\lambda_1(\gamma_j)}$ for large $j \geq 1$ and $\partial_s u_j(s) < 0$ ($0 < s < \sqrt{\lambda_1(\gamma_j)}$). Thus, $\partial_s \tilde{u}_*(s) \leq 0$ ($0 < s < \infty$). Let $v := \partial_s \tilde{u}_*$. If there is $\tilde{s}_2 > 0$ such that $v(\tilde{s}_2) = 0$, then $v_s(\tilde{s}_2) = 0$, since $v(s) \leq 0$ ($0 < s < \infty$). We

obtain a contradiction, because v satisfies

$$v_{ss} + \frac{N-1}{s}v_s + \left(f'(\tilde{u}_*) - \frac{N-1}{s^2} \right) v = 0 \quad (0 < s < \infty)$$

and Hopf's lemma says that $v_s(\tilde{s}_2) \neq 0$. Therefore, $v(s) < 0$ ($0 < s < \infty$) and $\partial_s \tilde{u}_*(s) < 0$ ($0 < s < \infty$). Since $\tilde{u}_* > 0$ and $\partial_s \tilde{u}_* < 0$, $\tilde{u}_*(\infty)$ exists and $\partial_s \tilde{u}_*(\infty) = 0$. Since \tilde{u}_* satisfies (1.7), $\partial_{ss} \tilde{u}_*(\infty)$ exists and it follows from the boundedness of \tilde{u}^* that $\partial_{ss} \tilde{u}_*(\infty) = 0$. Because of (1.7), $f(\tilde{u}_*(\infty)) = 0$, hence $\tilde{u}_*(\infty) = 0$ or 1. Suppose that $\tilde{u}_*(\infty) = 1$. Let $w := u - 1$. Then w satisfies

$$(3.4) \quad w_{ss} + \frac{N-1}{s}w_s + \frac{(w+1)^p - (w+1)}{w}w = 0.$$

If w is close to 0, then $\{(w+1)^p - (w+1)\}/w$ is close to $p-1$. Since $p-1 > 0$, by Sturm's oscillation theorem we see that w oscillates around 0 which contradicts that $w_s < 0$. Thus, $\tilde{u}_*(\infty) = 0$. We have shown that (3.3) holds and $\tilde{u}_* \equiv u_*$.

By Proposition 2.3 we see that if $p \geq p_S$, then (3.2) has no solution. We obtain a contradiction. Then, (3.1) does not hold, and $|\lambda_1(\gamma)|$ is bounded for $\gamma \in (\gamma^* - \varepsilon, \gamma^*)$. Therefore, there exists a subsequence, which is still denoted by $\{\gamma_j\}$, such that $\lambda_1(\gamma_j)$ converges as $j \rightarrow \infty$. We define $\lambda_1(\gamma^*) = \lim_{j \rightarrow \infty} \lambda_1(\gamma_j)$. Because of Proposition 3.1, $\lambda_1(\gamma^*)$ does not depend on the subsequence and it is uniquely determined. Moreover, $u_s(\sqrt{\lambda_1(\gamma^*)}, \gamma^*) = 0$.

Second, we prove the case $n \geq 2$. Let E be as given by (2.8). Then

$$(3.5) \quad \frac{d}{ds} E(u(s), v(s)) = -\frac{N-1}{s}v^2 \leq 0.$$

Therefore, $E(u(s), v(s)) \leq E(u(\sqrt{\lambda_1(\gamma^*)}), 0) < 0$ ($s > \sqrt{\lambda_1(\gamma^*)}$), which implies that

$$(3.6) \quad u(s) \geq u(\sqrt{\lambda_1(\gamma^*)}) (> 0).$$

Let $w(s) := u(s) - 1$. Then w satisfies (3.4). Because of (3.6), there is $\delta > 0$ such that

$$\frac{(w+1)^p - (w+1)}{w} > \frac{u(\sqrt{\lambda_1(\gamma^*)})^p - u(\sqrt{\lambda_1(\gamma^*)})}{u(\sqrt{\lambda_1(\gamma^*)}) - 1} > \delta > 0 \quad (\gamma > 0)$$

provided that $s > \sqrt{\lambda_1(\gamma^*)}$. Sturm's oscillation theorem says that w oscillates around 0 infinitely many times. Therefore, $u(s)$ has the n -th positive critical point s_n . Since $\lambda_n(\gamma^*) = s_n^2$ and $\lambda_n(\gamma)$ is continuous at γ^* (Proposition 3.1), we see that $|\lambda_n(\gamma)| < C$ for $\gamma \in (\gamma^* - \varepsilon, \gamma^*)$. \square

Using Lemma 3.3, we prove the global parametrization result.

Lemma 3.4. *For each $n \geq 1$, $\{(\lambda_n(\gamma), U_n(\gamma))\}$ can be defined in $1 < \gamma < \infty$.*

Proof. Let $\mathcal{C}_n := \{(\lambda_n(\gamma), U_n(\gamma))\}$ be the branch which emanates from $(\bar{\lambda}_n, 1)$. We extend \mathcal{C}_n . We define

$$\gamma^* := \sup\{\bar{\gamma} > 1; \lambda_n(\gamma) < \infty \text{ for every } \gamma \in (1, \bar{\gamma})\}.$$

Suppose the contrary, i.e., $\gamma^* < \infty$. Let $\{\gamma_j\}_{j=1}^\infty$ ($\gamma_j < \gamma^*$, $\gamma_j \uparrow \gamma^*$) be a sequence. Let $U_{n,j}(r)$ be a solution of (1.3) such that $U_{n,j}(0) = \gamma_j$ and $\partial_r U_{n,j}(0) = 0$. Because of Lemma 3.3, there are $\lambda_* \geq 0$ and a subsequence of $\{\gamma_j\}$, which is still denoted by $\{\gamma_j\}$, such that $\lambda_n(\gamma_j) \rightarrow \lambda_*$ ($j \rightarrow \infty$). Note that it is clear that $\lambda_* > 0$. Because of the continuous dependence of the solution $U(r, \gamma, \lambda)$ of

$$(3.7) \quad \begin{cases} U_{rr} + \frac{N-1}{r}U_r + \lambda f(U) = 0, & 0 < r < 1, \\ U_r(0) = 0, \quad U(0) = \gamma \end{cases}$$

on (γ, λ) in $C_{loc}^1[0, \infty)$, there exists a solution U_* of (3.7) with $(\gamma, \lambda) = (\gamma^*, \lambda_*)$ such that $\partial_r U_*(1) = 0$. Since each zero of $U_n(\cdot, \gamma) - 1$ is simple, $\mathcal{Z}_{(0,1)}[U_*(\cdot) - 1] = \mathcal{Z}_{(0,1)}[U_n(\cdot, \gamma_j) - 1] = n$ for large $j \geq 0$. Therefore, (λ_*, U_*) is a solution of (1.3) and $(\lambda_*, U_*) \in \mathcal{C}_n$. By Proposition 3.1 we can extend the branch \mathcal{C}_n to $1 < \gamma < \gamma^* + \varepsilon$ for small $\varepsilon > 0$. This contradicts to the definition of γ^* . Thus, $\gamma^* = \infty$. \square

The branch \mathcal{C}_n^+ is unbounded in γ .

3.4. Extension to $0 < \gamma < 1$. We extend \mathcal{C}_n^- .

Lemma 3.5. *For each $n \geq 1$, $\{(\lambda_n(\gamma), U_n(\gamma))\}$ can be defined in $0 < \gamma < 1$.*

Proof. Let

$$\gamma_* := \inf\{\underline{\gamma} > 0; \lambda_n(\gamma) < \infty \text{ for every } \underline{\gamma} < \gamma < 1\}.$$

Suppose the contrary, i.e., $\gamma_* > 0$. Let $u(s)$ be the solution of the (1.7) with $\gamma = \underline{\gamma}$, and let $v(s) := u_s(s)$. Let E be as given by (2.8). Then (3.5) holds. Therefore, $E(u(s), v(s)) \leq E(\underline{\gamma}, 0) < 0$, which implies that

$$(3.8) \quad u(s) \geq u(0) (= \underline{\gamma} > 0).$$

Let $w(s) := u(s) - 1$. Then w satisfies (3.4). Because of (3.8), there is $\delta > 0$ such that $\{(w+1)^p - (w+1)\}/w > (u(0)^p - u(0))/(u(0) - 1) > \delta > 0$ ($\gamma > 0$). Sturm's oscillation theorem says that w oscillates around 0 infinitely many times. Therefore, $u(s)$ has the n -th positive critical point s_n . Let $\lambda_n := s_n^2$. Then $(\lambda(\underline{\gamma}), u)$ is a solution of (1.7) with $\gamma = \underline{\gamma}$. By Proposition 3.1 we can extend $\lambda_n(\underline{\gamma})$, which contradicts to the definition of $\underline{\gamma}$. Thus, $\underline{\gamma} = 0$, and $\lambda_n(\gamma)$ can be defined in $0 < \gamma < 1$. \square

The branch \mathcal{C}_n^- is defined in $0 < \gamma < 1$.

4. ENTIRE SINGULAR SOLUTION

In this section we prove Theorem A. Let $(\lambda, U(r))$ be a solution of (1.3). Then $u(s) := U(r)$ ($s := \sqrt{\lambda}r$) satisfies (1.14). We use the same change of variables $y(t) := A(p, N)^{-1}s^\theta u(s)$ and $t := m^{-1} \log s$ as in Section 2. Here

$$m := \{\theta(N - 2 - \theta)\}^{-\frac{1}{2}},$$

and θ is defined by (1.6). The function $y(t)$ satisfies

$$(4.1) \quad y'' + \alpha y' - y + y^p - m^2 e^{2mt} y = 0,$$

where $\alpha := m(N - 2 - 2\theta)$. We mainly consider (4.1) in this section.

4.1. Existence of the singular solution near $r = 0$. We construct the singular solution of (1.14) near $s = 0$.

Lemma 4.1. *The problem*

$$(4.2) \quad \begin{cases} y'' + \alpha y' - y + y^p - m^2 e^{2mt} y = 0, \\ y(t) \rightarrow 1 \quad (t \rightarrow -\infty) \end{cases}$$

has a unique solution.

The proof is essentially the same as one of [22, Theorem A]. We use the following lemma to prove Lemma 4.1.

Lemma 4.2. *Assume that (4.2) has a solution $y^*(t)$. Then, $y^*(t)$ satisfies*

$$(4.3) \quad y^*(t) = 1 + O(e^{2mt}) \quad (t \rightarrow -\infty).$$

Proof. Let $\zeta := -t$ and $\eta(\zeta) := y(t) - 1$. The function $\eta(\zeta)$ satisfies

$$(4.4) \quad \begin{cases} \eta'' - \alpha\eta' + (p-1)\eta = g(\eta), & \zeta_0 < \zeta < \infty, \\ \eta(\zeta) \rightarrow 0 & (\zeta \rightarrow \infty), \end{cases}$$

where ζ_0 is a large number,

$$(4.5) \quad g(\zeta) := m^2 e^{-2m\zeta} (1 + \eta(\zeta)) - \varphi(\eta(\zeta)),$$

$$(4.6) \quad \varphi(\eta) := (1 + \eta)^p - 1 - p\eta.$$

There are three possibilities:

$$(4.7) \quad (1) p - 1 > \left(\frac{\alpha}{2}\right)^2, \quad (2) p - 1 < \left(\frac{\alpha}{2}\right)^2, \quad (3) p - 1 = \left(\frac{\alpha}{2}\right)^2.$$

Case (1): Let $\beta := \sqrt{(p-1) - \left(\frac{\alpha}{2}\right)^2}$. Because the linearly independent solutions of the homogeneous equations associated with the equation in (4.4) becomes unbounded as $\zeta \rightarrow \infty$, we have

$$(4.8) \quad \eta(\zeta) = \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \sin(\beta(\sigma - \zeta)) g(\sigma) d\sigma.$$

If $|\eta|$ is small, then there are $\varepsilon > 0$ and $\zeta_\varepsilon > 0$ such that

$$(4.9) \quad |\varphi(\eta)| \leq |(1 + \eta)^p - 1 - p\eta| \leq \varepsilon|\eta| \quad (\zeta > \zeta_\varepsilon).$$

By (4.5) and (4.9) we have

$$(4.10) \quad |g(\zeta)| \leq C_0 e^{-2m\zeta} + \varepsilon|\eta(\zeta)| \quad (\zeta > \zeta_\varepsilon).$$

Using (4.8), we have

$$(4.11) \quad \begin{aligned} \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma &\leq \int_{\zeta}^{\infty} \frac{e^{\frac{\alpha}{2}\sigma}}{\beta} \int_{\sigma}^{\infty} e^{-\frac{\alpha}{2}\tau} |g(\tau)| d\tau d\sigma \\ &= \left[\frac{2e^{\frac{\alpha}{2}\sigma}}{\alpha\beta} \int_{\sigma}^{\infty} e^{-\frac{\alpha}{2}\tau} |g(\tau)| d\tau \right]_{\zeta}^{\infty} + \int_{\zeta}^{\infty} \frac{2e^{\frac{\alpha}{2}\sigma}}{\alpha\beta} e^{-\frac{\alpha}{2}\sigma} |g(\sigma)| d\sigma \\ &\leq -\frac{2e^{\frac{\alpha}{2}\zeta}}{\alpha\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} |g(\sigma)| d\sigma + \frac{2}{\alpha\beta} \int_{\zeta}^{\infty} |g(\sigma)| d\sigma \\ &\leq \frac{2}{\alpha\beta} \int_{\zeta}^{\infty} |g(\sigma)| d\sigma, \end{aligned}$$

where we use

$$\lim_{\sigma \rightarrow \infty} \frac{2 \int_{\sigma}^{\infty} e^{-\frac{\alpha}{2}\tau} |g(\tau)| d\tau}{\alpha\beta e^{-\frac{\alpha}{2}\sigma}} = \lim_{\sigma \rightarrow \infty} \frac{-2e^{-\frac{\alpha}{2}\sigma} |g(\sigma)|}{\alpha\beta \left(-\frac{\alpha}{2}\right) e^{-\frac{\alpha}{2}\sigma}} = 0.$$

By (4.10) and (4.11) we have

$$\int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma \leq \frac{2}{\alpha\beta} \int_{\zeta}^{\infty} (C_0 e^{-2m\sigma} + \varepsilon|\eta(\sigma)|) d\sigma \quad (\zeta > \zeta_\varepsilon).$$

Hence,

$$(4.12) \quad \begin{aligned} \left(1 - \frac{2\varepsilon}{\alpha\beta}\right) \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma &\leq \frac{2C_0}{\alpha\beta} \int_{\zeta}^{\infty} e^{-2m\sigma} d\sigma \\ &= \frac{C_0 e^{-2m\zeta}}{\alpha\beta m} \quad (\zeta > \zeta_\varepsilon). \end{aligned}$$

On the other hand, by (4.8) and (4.10) we have

$$(4.13) \quad \begin{aligned} |\eta(\zeta)| &\leq \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} (C_0 e^{-2m\sigma} + \varepsilon |\eta(\sigma)|) d\sigma \\ &= \frac{C_0}{\beta} e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2}+2m)\sigma} d\sigma + \frac{\varepsilon}{\beta} \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma \\ &= \frac{C_0 e^{-2m\zeta}}{\beta(\frac{\alpha}{2}+2m)} + \frac{\varepsilon}{\beta} \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma. \end{aligned}$$

By (4.12) and (4.13) we have

$$(4.14) \quad \begin{aligned} |\eta(\zeta)| &\leq \frac{C_0 e^{-2m\zeta}}{\beta(\frac{\alpha}{2}+2m)} + \frac{C_0 \varepsilon}{\left(1 - \frac{2\varepsilon}{\alpha\beta}\right) \alpha\beta^2 m} e^{-2m\zeta} \\ &=: C_1 e^{-2m\zeta}. \end{aligned}$$

Case (2): Let $\beta := \sqrt{\left(\frac{\alpha}{2}\right)^2 - (p-1)}$. Then, $\frac{\alpha}{2} - \beta > 0$. We have

$$(4.15) \quad \eta(\zeta) = \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \sinh(\beta(\sigma - \zeta)) g(\sigma) d\sigma.$$

Using (4.15) and $|\sinh(\beta(\tau - \sigma))| \leq e^{\beta(\tau - \sigma)}/2$ ($\tau \geq \sigma$), we have

$$\begin{aligned}
 \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma &\leq \int_{\zeta}^{\infty} \frac{e^{\frac{\alpha}{2}\sigma}}{\beta} \int_{\sigma}^{\infty} e^{-\frac{\alpha}{2}\tau} |\sinh(\beta(\tau - \sigma))| |g(\tau)| d\tau d\sigma \\
 &\leq \int_{\zeta}^{\infty} \frac{e^{\frac{\alpha}{2}\sigma}}{\beta} \int_{\sigma}^{\infty} e^{-\frac{\alpha}{2}\tau} \frac{e^{\beta(\tau - \sigma)}}{2} |g(\tau)| d\tau d\sigma \\
 &= \int_{\zeta}^{\infty} \frac{e^{(\frac{\alpha}{2} - \beta)\sigma}}{2\beta} \int_{\sigma}^{\infty} e^{-(\frac{\alpha}{2} - \beta)\tau} |g(\tau)| d\tau d\sigma \\
 (4.16) \quad &= \left[\frac{e^{(\frac{\alpha}{2} - \beta)\sigma}}{\beta(\alpha - 2\beta)} \int_{\sigma}^{\infty} e^{-(\frac{\alpha}{2} - \beta)\tau} |g(\tau)| d\tau \right]_{\zeta}^{\infty}
 \end{aligned}$$

$$\begin{aligned}
 &+ \frac{1}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} |g(\sigma)| d\sigma \\
 (4.17) \quad &= -\frac{e^{(\frac{\alpha}{2} - \beta)\zeta}}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2} - \beta)\tau} |g(\tau)| d\tau \\
 &+ \frac{1}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} |g(\sigma)| d\sigma
 \end{aligned}$$

$$\begin{aligned}
 (4.18) \quad &\leq \frac{1}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} |g(\sigma)| d\sigma,
 \end{aligned}$$

where we use $\frac{\alpha}{2} - \beta > 0$ and

$$\lim_{\sigma \rightarrow \infty} \frac{\int_{\sigma}^{\infty} e^{-(\frac{\alpha}{2} - \beta)\tau} |g(\tau)| d\tau}{\beta(\alpha - 2\beta) e^{-(\frac{\alpha}{2} - \beta)\sigma}} = \lim_{\sigma \rightarrow \infty} \frac{-e^{-(\frac{\alpha}{2} - \beta)\sigma} |g(\sigma)|}{\beta(\alpha - 2\beta) (-\frac{\alpha}{2} + \beta) e^{-(\frac{\alpha}{2} - \beta)\sigma}} = 0.$$

In the case (2) (4.10) also holds. By (4.10) and (4.18) we have

$$\int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma \leq \frac{1}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} (C_0 e^{-2m\sigma} + \varepsilon |\eta(\sigma)|) d\sigma \quad (\zeta > \zeta_{\varepsilon}).$$

Hence,

$$\begin{aligned}
 \left(1 - \frac{\varepsilon}{\beta(\alpha - 2\beta)}\right) \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma &\leq \frac{C_0}{\beta(\alpha - 2\beta)} \int_{\zeta}^{\infty} e^{-2m\sigma} d\sigma \\
 (4.19) \quad &= \frac{C_0}{2m\beta(\alpha - 2\beta)} e^{-2m\zeta} \quad (\zeta > \zeta_{\varepsilon}).
 \end{aligned}$$

On the other hand, by (4.15) and (4.10) we have

$$\begin{aligned}
|\eta(\zeta)| &\leq \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \frac{e^{\beta(\sigma-\zeta)}}{2} (C_0 e^{-2m\sigma} + \varepsilon |\eta(\sigma)|) d\sigma \\
&= \frac{C_0}{2\beta} e^{(\frac{\alpha}{2}-\beta)\zeta} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2}-\beta+2m)\sigma} d\sigma \\
&\quad + \frac{\varepsilon}{2\beta} e^{(\frac{\alpha}{2}-\beta)\zeta} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2}-\beta)\sigma} |\eta(\sigma)| d\sigma \\
(4.20) \quad &\leq \frac{C_0 e^{-2m\zeta}}{2\beta (\frac{\alpha}{2} - \beta + 2m)} + \frac{\varepsilon}{2\beta} \int_{\zeta}^{\infty} |\eta(\sigma)| d\sigma.
\end{aligned}$$

By (4.19) and (4.20) we have

$$\begin{aligned}
|\eta(\zeta)| &\leq \frac{C_0 e^{-2m\zeta}}{\beta(\alpha - 2\beta + 4m)} + \frac{C_0 \varepsilon}{4m\beta(\beta(\alpha - 2\beta) - \varepsilon)} e^{-2m\zeta} \\
(4.21) \quad &=: C_2 e^{-2m\zeta} \quad (\zeta > \zeta_\varepsilon).
\end{aligned}$$

Case (3): In this case we have

$$(4.22) \quad \eta(\zeta) = e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} (\sigma - \zeta) g(\sigma) d\sigma.$$

For each $\beta > 0$, we have

$$|\sigma - \zeta| \leq \frac{1}{\beta} |\sinh(\beta(\sigma - \zeta))|.$$

Hence,

$$|\eta(\zeta)| \leq \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} |\sinh(\beta(\sigma - \zeta))| |g(\sigma)| d\sigma.$$

Therefore, by the same argument as in Case (2) we have (4.21). However, β is not the same value as in Case (2).

We have verified all the cases. \square

Proof of Lemma 4.1. There are three cases (4.7) as in the proof of Lemma 4.2. We consider only the case (1).

We transform (4.2) to the integral equation

$$(4.23) \quad \eta(\zeta) = \mathcal{F}(\eta)(\zeta).$$

The integral operator \mathcal{F} is different in each case. In this proof we write $g(\eta, \sigma)$ in order to stress the dependence of g on η . In the case (1) \mathcal{F} becomes

$$\mathcal{F}(\eta)(\zeta) = \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \sin(\beta(\sigma - \zeta)) g(\eta, \sigma) d\sigma.$$

We show by contraction mapping theorem that (4.23) has a unique solution. Let $\zeta_0 > 0$ be large. Hereafter, by $\|\cdot\|$ we denote $\|\cdot\|_{C^0[\zeta_0, \infty)}$. We set $X := \{\eta(\zeta) \in C^0[\zeta_0, \infty); \|\eta(\zeta)\| < \infty\}$ and $\mathcal{B} := \{\eta(\zeta) \in X; \|\eta\| < \bar{\eta}\}$. Let $\eta_1, \eta_2 \in \mathcal{B}$. If $\bar{\eta} > 0$ is small, then there is a small $\varepsilon > 0$ such that

$$(4.24) \quad |\varphi(\eta_1) - \varphi(\eta_2)| \leq \varepsilon|\eta_1 - \eta_2|.$$

Using (4.24), we have

$$(4.25) \quad \begin{aligned} |g(\eta_1, \sigma) - g(\eta_2, \sigma)| &\leq m^2 e^{-2m\zeta} |\eta_1 - \eta_2| + \varepsilon |\eta_1 - \eta_2| \\ &\leq 2\varepsilon |\eta_1 - \eta_2|, \end{aligned}$$

provided that $\zeta > 0$ is large. By (4.25) we have

$$(4.26) \quad \begin{aligned} \|\mathcal{F}(\eta_1) - \mathcal{F}(\eta_2)\| &\leq \left\| \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} |g(\eta_1, \sigma) - g(\eta_2, \sigma)| d\sigma \right\| \\ &\leq 2\varepsilon \left\| \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} d\sigma \right\| \|\eta_1 - \eta_2\| \\ &= \frac{4\varepsilon}{\alpha\beta} \|\eta_1 - \eta_2\|. \end{aligned}$$

On the other hand,

$$(4.27) \quad \begin{aligned} \|\mathcal{F}(0)\| &\leq \left\| \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} |g(0, \sigma)| d\sigma \right\| \\ &\leq \left\| \frac{e^{\frac{\alpha}{2}\zeta}}{\beta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} m^2 e^{-2m\sigma} d\sigma \right\| \\ &= \frac{m^2}{\beta \left(\frac{\alpha}{2} + 2m\right)} e^{-2m\zeta} \\ &< \varepsilon, \end{aligned}$$

provided that $\zeta > 0$ is large. By (4.26) and (4.27) we see that if $\varepsilon > 0$ is small, then

$$(4.28) \quad \begin{aligned} \|\mathcal{F}(\eta_1)\| &= \|\mathcal{F}(\eta_1) - \mathcal{F}(0) + \mathcal{F}(0)\| \\ &\leq \|\mathcal{F}(\eta_1) - \mathcal{F}(0)\| + \|\mathcal{F}(0)\| \\ &\leq \frac{4\varepsilon}{\alpha\beta} \|\eta_1\| + \varepsilon \\ &< \bar{\eta}. \end{aligned}$$

Because of (4.26) and (4.28), if $\varepsilon > 0$ is small, then \mathcal{F} is a contraction mapping from \mathcal{B} into itself. Therefore, (4.23) has a unique solution. \square

Hereafter, let $y^*(t)$ be the solution given by Lemma 4.1, and

$$(4.29) \quad u^*(s) := As^{-\theta} y^*(m^{-1} \log s).$$

Then the following corollary is an immediate consequence of Lemmas 4.1 and 4.2.

Corollary 4.3.

$$(4.30) \quad u^*(s) = As^{-\theta} (1 + o(1)) \quad \text{as } s \downarrow 0.$$

Here, $A = A(p, N)$.

4.2. Positivity of the entire singular solution.

Lemma 4.4. *The domain of the singular solution $y^*(t)$ obtained in Lemma 4.1 can be extended to $t \in (-\infty, \infty)$, and $y^*(t) > 0$ ($-\infty < t < \infty$). Therefore, the domain of $u^*(s)$ can be extended to $s \in (0, \infty)$, and $u^*(s) > 0$ ($0 < s < \infty$).*

Proof. In the proof we denote y^* by y for simplicity. We extend the domain of $y(t)$. The equation (4.1) is equivalent to

$$(4.31) \quad \begin{cases} y' = z, \\ z' = -\alpha z + y - y^p + m^2 e^{2mt} y. \end{cases}$$

Then, the solution of (4.1) can be considered as the orbit $(y(t), z(t))$ of (4.31). Let

$$(4.32) \quad \tilde{E}(y, z) := \frac{z^2}{2} - \frac{\beta(t)}{2} y^2 + \frac{y^{p+1}}{p+1},$$

where $\beta(t) := 1 + m^2 e^{2mt}$. Then

$$(4.33) \quad \frac{d}{dt} \tilde{E}(y(t), z(t)) = -\alpha z^2 - m^3 e^{2mt} y^2 \leq 0.$$

Since $(y(-\infty), z(-\infty)) = (1, 0)$, $\tilde{E}(y(-\infty), z(-\infty)) = -1/2 + 1/(p+1) < 0$. Because of (4.33), $(y(t), z(t)) \in \{(y, z); \tilde{E}(y, z) < 0\}$. Since $-\beta(t)y^2/2 + y^{p+1}/(p+1) < 0$,

$$(4.34) \quad 0 < y(t) < \left(\frac{p+1}{2} \beta(t) \right)^{\frac{1}{p-1}} \quad (-\infty < t < \infty).$$

Therefore, $y(t) > 0$ ($-\infty < t < \infty$). Since $\tilde{E}(y, z) < 0$, $z^2 < \beta(t)y^2 - 2y^{p+1}/(p+1)$, hence

$$(4.35) \quad 0 < z(t) < \left(\frac{p-1}{p+1} \beta(t)^{1+\theta} \right)^{\frac{1}{2}} \quad (-\infty < t < \infty).$$

Because of (4.34) and (4.35), neither $y(t)$ nor $z(t)$ blows up. Hence, the solution $(y(t), z(t))$ of (4.31) can be extended to $t \in (-\infty, \infty)$. Because of (4.29), the conclusion about $u^*(s)$ also holds. \square

4.3. Nonexistence of the entire singular solution.

Lemma 4.5. *Let $y^*(t)$ be as in Lemma 4.1. Then,*

$$(4.36) \quad (y^*)'(t) = \frac{m}{2(N-1) - 3\theta} e^{2mt} + O(e^{2m(1+\nu)t}) \quad (t \rightarrow -\infty),$$

where $\nu = \min\{p-1, 1\}$.

Proof. Let $\zeta := -t$ and $\eta(\zeta) = y(t) - 1$. By (4.6) and (4.14) we have

$$(4.37) \quad |\varphi(\eta)| \leq C_1 e^{-2m(1+\nu)\zeta}.$$

Because of (4.6) and (4.37), we have

$$(4.38) \quad g(\zeta) = m^2 e^{-2m\zeta} + O(e^{-2m(1+\nu)\zeta}).$$

There are three cases (4.7) as in the proof of Lemma 4.2. We consider only the case (1).

Differentiating (4.8) with respect to ζ , we have

$$(4.39) \quad \eta'(\zeta) = \frac{\alpha}{2\beta} e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \sin(\beta(\sigma - \zeta)) g(\sigma) d\sigma \\ - e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-\frac{\alpha}{2}\sigma} \cos(\beta(\sigma - \zeta)) g(\sigma) d\sigma.$$

By (4.38) and (4.39) we have

$$\begin{aligned} \eta'(\zeta) &= \frac{\alpha m^2}{2\beta} e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2}+2m)\sigma} \sin(\beta(\sigma - \zeta)) d\sigma \\ &\quad - m^2 e^{\frac{\alpha}{2}\zeta} \int_{\zeta}^{\infty} e^{-(\frac{\alpha}{2}+2m)\sigma} \cos(\beta(\sigma - \zeta)) d\sigma + O(e^{-2m(1+\nu)\zeta}) \\ &= \frac{-2m^3}{(\frac{\alpha}{2} + 2m)^2 + \beta^2} e^{-2m\zeta} + O(e^{-2m(1+\nu)\zeta}) \\ &= -\frac{m}{2(N-1) - 3\theta} e^{-2m\zeta} + O(e^{-2m(1+\nu)\zeta}). \end{aligned}$$

Since $(y^*)'(t) = -\eta'(\zeta)$, we obtain (4.36). In the other cases (2) and (3) we also obtain (4.36), using (4.15) and (4.22). \square

Corollary 4.6. *Suppose that $p > p_s$. Let $\delta > 0$. Then, $u^* \in H^1(B_\delta)$, and*

$$(4.40) \quad u_s^*(s) = -\theta A s^{-\theta-1} (1 + o(1)) \quad (s \downarrow 0).$$

Proof. Differentiating (4.29) with respect to s , we have

$$\begin{aligned} u_s^*(s) &= -\theta A s^{-\theta-1} y^* \left(\frac{1}{m} \log s \right) + A s^{-\theta} (y^*)' \left(\frac{1}{m} \log s \right) \frac{1}{s} \\ &= -\theta A s^{-\theta-1} (1 + o(1)) + A s^{-\theta-1} o(s) \\ &= -\theta A s^{-\theta-1} (1 + o(1)), \end{aligned}$$

where we use Lemma 4.5. Then, we obtain (4.40). We show that

$$\int_0^\delta \{ (u^*)^2 + (u_s^*)^2 \} s^{N-1} ds < \infty.$$

We have

$$\begin{aligned} \{ u^*(s)^2 + u_s^*(s)^2 \} s^{N-1} &\leq (C_0 s^{-2\theta} + C_1 s^{-2\theta-2}) s^{N-1} \\ &= C_0 s^{-2\theta+N-1} + C_1 s^{-2\theta-2+N-1}. \end{aligned}$$

Since $p > p_S$, $-2\theta - 2 + N - 1 > -1$ and $-2\theta + N - 1 > -1$. Hence, $\int_0^\delta (C_0 s^{-2\theta+N-1} + C_1 s^{-2\theta-2+N-1}) ds < \infty$, which indicates that $u^* \in H^1(B_\delta)$. \square

Lemma 4.7. *Suppose that $p > p_S$. The problem*

$$(4.41) \quad \begin{cases} u_{ss} + \frac{N-1}{s} u_s + f(u) = 0, & 0 < s < \infty, \\ u(s) = A s^{-\theta} (1 + o(1)) \quad (s \downarrow 0), \\ u(\infty) = 0, \\ u > 0, & 0 < s < \infty \end{cases}$$

does not admit a solution.

This lemma was proved by [25]. See [5] for details of nonexistence results. However, we briefly prove the lemma.

Proof. Suppose that (4.41) has a solution. Because $u(s) = A s^{-\theta} (1 + o(1))$ ($s \downarrow 0$) and the uniqueness of the solution $u^*(s)$ obtained by Lemma 4.1, the solution of (4.41) is $u^*(s)$. We denote u^* by u for simplicity. Because $f'(0) = -1 < 0$, $u(s)$ and $u_s(s)$ decay exponentially as $s \rightarrow \infty$. Thus, the four terms $R^{N-1} u(R) u_s(R)$, $R^N u_s(R)^2$, $R^N u(R)^2$, and $R^N u(R)^{p+1}$ converge to zero as $R \rightarrow \infty$. Because of (4.30) and (4.40), we see that the four terms $\varepsilon^{N-1} u(\varepsilon) u_s(\varepsilon)$, $\varepsilon^N u_s(\varepsilon)^2$, $\varepsilon^N u(\varepsilon)^2$, and $\varepsilon^N u(\varepsilon)^{p+1}$ converge to zero as $\varepsilon \downarrow 0$. Taking the limit of (2.10) as $\varepsilon \downarrow 0$ and $R \rightarrow \infty$, we have

$$\left(\frac{N}{2} - 1 - \frac{N}{p+1} \right) \int_0^\infty s^{N-1} u_s^2 ds + \left(\frac{N}{2} - \frac{N}{p+1} \right) \int_0^\infty s^{N-1} u^2 ds = 0.$$

Since $p > p_S$, $\frac{N}{2} - 1 - \frac{N}{p+1} > 0$. The above equality implies that (4.41) does not admit a solution. \square

4.4. Existence of critical points of $u^*(s)$. The main technical lemma in this section is the following:

Lemma 4.8. *Suppose that $p_S < p < p_{JL}$. There is a sequence $\{s_n^*\}_{n=0}^\infty$ ($0 = s_0^* < s_1^* < s_2^* < \dots \rightarrow \infty$) such that $\partial_s u^*(s_n^*) = 0$ ($n \geq 1$),*

$$u^*(s) \begin{cases} < 1 & (s \in \{s_1^*, s_3^*, \dots\}), \\ > 1 & (s \in \{s_2^*, s_4^*, \dots\}) \end{cases}$$

and

$$\partial_s u^*(s) \begin{cases} < 0 & (s \in (s_0^*, s_1^*) \cup (s_2^*, s_3^*) \cup \dots), \\ > 0 & (s \in (s_1^*, s_2^*) \cup (s_3^*, s_4^*) \cup \dots). \end{cases}$$

Proof. First, we show that u^* has the first positive critical point. We use a contradiction argument. Suppose the contrary, i.e.,

$$(4.42) \quad \partial_s u^*(s) < 0 \quad (0 < s < \infty).$$

Since $u^*(s) > 0$ ($0 < s < \infty$), $u^*(s)$ converges as $s \rightarrow \infty$. Since $\partial_s u^*$ does not change sign, $\partial_s u^*(s) \rightarrow 0$ ($s \rightarrow \infty$). We see by (1.14) that $\partial_{ss} u^*(s)$ converges as $s \rightarrow \infty$. Since $u^*(s)$ converges, $\partial_{ss} u^*(s) \rightarrow 0$ ($s \rightarrow \infty$). Therefore, $f(u^*(\infty)) = 0$, and $u^*(\infty) = 0$ or 1 . If $u^*(\infty) = 1$, then by the same argument as in the proof of Lemma 3.3 we see that u^* should oscillate around 1 . This oscillation contradicts that $\partial_s u^*(s) < 0$ ($0 < s < \infty$). Thus, $u^*(\infty) = 0$. We see that u^* is a solution of (4.41), which contradicts to Lemma 4.7. Therefore, there is $s_1^* > 0$ such that $\partial_s u^*(s) < 0$ ($0 < s < s_1^*$) and $\partial_s u^*(s_1^*) = 0$.

Second, we show that there exists s_n^* ($n \geq 2$). Let $v^* := u_s^*$, and let E be as given in (2.8). Then

$$\frac{d}{ds} E(u^*(s), v^*(s)) = -\frac{N-1}{s} (v^*(s))^2 < 0.$$

Therefore, $E(u^*(s), v^*(s)) < E(u^*(s_1^*), 0) < 0$ ($s > s_1^*$). This implies that

$$(4.43) \quad u^*(s) > u^*(s_1^*) \quad (s > s_1^*).$$

Let $w(s) := u^*(s) - 1$. Then w satisfies (3.4). Because of (4.43), $\{(w+1)^p - (w+1)\}/w > \{(u^*(s_1^*))^p - u^*(s_1^*)\}/(u^*(s_1^*) - 1) > 0$. Sturm's oscillation theorem says that w oscillates around 0 infinitely many times

as $s \rightarrow \infty$. Consequently, u^* has infinitely many critical points. Because u^* satisfies the ODE, we easily see that u^* does not have a critical point on $\{u^* = 1\}$ and that if u^* has a critical point on $\{u^* > 1\}$ (resp. $\{u^* < 1\}$), then it is a local maximum point (resp. local minimum point). Thus, the conclusion of the lemma holds. \square

4.5. Proof of Theorem A. Since the convergence property in Theorem 1.1 is proved in Section 5, the proof of Theorem 1.1 is postponed until Section 5. Here we prove only Theorem A.

Proof of Theorem A. Let $\lambda_n^* := (s_n^*)^2$. Then, $(\lambda_n^*, u^*(s))$ is a singular solution of (1.14) which satisfies (4.30). We define $U_n^*(r) := u^*(\sqrt{\lambda_n^*}r)$. Then by Corollary 4.3 and Lemmas 4.4 and 4.8 we see that U_n^* satisfies (i), (ii), and (iii). The uniqueness of (λ_n^*, U_n^*) follows from the uniqueness of the solution of (4.2) which was shown in Lemma 4.1. \square

5. CONVERGENCE TO THE SINGULAR SOLUTION AS $\gamma \rightarrow \infty$

Let $u(s, \gamma)$ denote the solution of (1.7). Note that $u(0, \gamma) = \gamma$. Let $y(t) = A(p, N)^{-1} s^\theta u(s, \gamma)$ and $t = m^{-1} \log s$. Then $y(t)$ satisfies (4.1). Let $n \in \mathbb{N}$ be fixed. Let λ_n^* be the number given in Theorem A, and let $y^*(t)$ be the singular solution given in Lemma 4.1. Then by Lemma 4.4 we see that $y(t)$ can be defined in \mathbb{R} . We set $t_n^* := m^{-1} \log \sqrt{\lambda_{n+1}^*}$.

5.1. Convergence to u^* . Our goal in this section is to prove the following:

Theorem 5.1. *Let $n \geq 1$ be fixed. Let $r_n^* := e^{mt_n^*}$. Suppose that $\{\gamma_j\}_{j=1}^\infty$ is an arbitrary sequence diverging to ∞ . Let $u^*(s)$ be the singular solution defined by (4.29), and let $u_j(s) := u(s, \gamma_j)$ be the solution of (1.7). Then, as $j \rightarrow \infty$,*

(5.1)

$$u_j \rightarrow u^*, \quad \partial_s u_j \rightarrow \partial_s u^*, \quad \text{and} \quad \partial_{ss} u_j \rightarrow \partial_{ss} u^* \quad \text{in} \quad C_{loc}^0(0, r_n^*].$$

We define

$$\tau(\gamma) := \frac{1}{m\theta}(\log \gamma - \log A).$$

Lemma 5.2. *Suppose that $\gamma > 0$ is large. Then, $y(t)$ satisfies*

$$y(t, \gamma) \leq e^{m\theta(t+\tau(\gamma))} \quad (-\infty < t \leq t_n^*).$$

Proof. Let $u(s) := u(s, \gamma)$ and $v(s) := u_s(s)$. Since $(u(s), v(s))$ satisfies (3.5),

$$(5.2) \quad -\frac{u(s)^2}{2} + \frac{u(s)^{p+1}}{p+1} \leq E(u(s), v(s)) \leq E(\gamma, 0) = -\frac{\gamma^2}{2} + \frac{\gamma^{p+1}}{p+1} \quad (s \geq 0).$$

Since $\gamma > 0$ is large, (5.2) tells us that $u(s) \leq \gamma$ ($s \geq 0$). By the definition of $y(t)$ we have

$$y(t) = A^{-1} s^\theta u(s) \leq A^{-1} e^{m\theta t} \gamma = e^{m\theta(t+\tau(\gamma))} \quad (-\infty < t \leq t_n^*).$$

□

We define

$$\xi := t + \tau(\gamma) \quad \text{and} \quad w(\xi, \gamma) := y(t, \gamma).$$

Then $w(\xi, \gamma)$ is a solution of the problem

$$(5.3) \quad \begin{cases} w'' + \alpha w' - w + w^p - m^2 e^{2m(\xi-\tau(\gamma))} w = 0, & -\infty < \xi < t_n^* + \tau(\gamma), \\ w > 0, & -\infty < \xi < t_n^* + \tau(\gamma), \\ e^{-m\theta\xi} w(\xi) \rightarrow 1 \quad (\xi \rightarrow -\infty). \end{cases}$$

Because of Lemma 5.2,

$$(5.4) \quad w(\xi, \gamma) \leq e^{m\theta\xi} \quad (-\infty < \xi \leq t_n^* + \tau(\gamma)).$$

Using the apriori estimate (5.4), we prove the following uniform convergence:

Lemma 5.3. *Let $\bar{w}(\xi)$ be a solution of*

$$(5.5) \quad \begin{cases} w'' + \alpha w' - w + w^p = 0, & -\infty < \xi < t_n^* + \tau(\gamma), \\ w > 0, & -\infty < \xi < t_n^* + \tau(\gamma), \\ e^{-m\theta\xi} w(\xi) \rightarrow 1 \quad (\xi \rightarrow -\infty). \end{cases}$$

Then, for each fixed $\bar{\xi} \in \mathbb{R}$,

$$w(\xi, \gamma) \rightarrow \bar{w}(\xi) \quad \text{and} \quad w'(\xi, \gamma) \rightarrow \bar{w}'(\xi) \quad \text{in} \quad C^0(-\infty, \bar{\xi})$$

as $\gamma \rightarrow \infty$.

Proof. Let $\varepsilon > 0$ be small. Since $\bar{w}(\xi)$ is a solution of (5.5),

$$(5.6) \quad \bar{w}(\xi) \rightarrow 0 \quad (\xi \rightarrow -\infty).$$

Because of (5.4) and (5.6), there is $\underline{\xi} (< \bar{\xi})$ such that

$$|w(\xi, \gamma)| \leq \frac{\varepsilon}{2} \quad \text{and} \quad |\bar{w}(\xi)| \leq \frac{\varepsilon}{2} \quad (-\infty < \xi \leq \underline{\xi}).$$

Then,

$$\begin{aligned}
 |w(\xi, \gamma) - \bar{w}(\xi)| &\leq |w(\xi, \gamma)| + |\bar{w}(\xi)| \\
 &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\
 (5.7) \qquad &= \varepsilon \quad (-\infty < \xi < \underline{\xi}).
 \end{aligned}$$

We show that

$$(5.8) \qquad |w(\xi, \gamma) - \bar{w}(\xi)| < \varepsilon \quad \text{in} \quad [\underline{\xi}, \bar{\xi}].$$

Since $y'(t) = A^{-1}ms^\theta(\theta u(s) + su'(s))$, $y'(t, \gamma) \rightarrow 0$ ($t \rightarrow -\infty$). Hence, $w'(\xi, \gamma) \rightarrow 0$ ($\xi \rightarrow -\infty$). Integrating the equation in (5.3) over $(-\infty, \xi]$, we have

$$(5.9) \qquad w'(\xi) + \alpha w(\xi) = \int_{-\infty}^{\xi} (w - w^p + m^2 e^{2m(\eta - \tau(\gamma))} w) d\eta,$$

where we use $w'(-\infty) = 0$ and $w(-\infty) = 0$. There is $C_0 > 0$ independent of large $\gamma > 0$ such that

$$\int_{-\infty}^{\xi} |w - w^p + m^2 e^{2m(\eta - \tau(\gamma))} w| d\eta < C_0 \quad (\xi \leq \bar{\xi}),$$

because of (5.4). Then $|w'(\xi)| \leq |\alpha w(\xi)| + C_0$, which indicates that w is equicontinuous in $[\underline{\xi}, \bar{\xi}]$. It follows from (5.4) that $w(\xi, \gamma)$ is uniformly bounded in $[\underline{\xi}, \bar{\xi}]$. By Arzelà-Ascoli theorem we see that $w(\xi, \gamma)$ converges to a certain function in $C^0[\underline{\xi}, \bar{\xi}]$ which satisfies the equation in (5.5). This function is a solution of (5.5). Using the same argument as in the proof of Lemma 4.1, we see that the solution of (5.5) is unique, hence the limit function is $\bar{w}(\xi)$. Therefore, (5.8) holds for large $\gamma > 0$. The estimates (5.7) and (5.8) indicate that $w(\xi, \gamma) \rightarrow \bar{w}(\xi)$ in $C^0(-\infty, \bar{\xi}]$ as $\gamma \rightarrow \infty$. In particular, this convergence is uniform.

Next, we show that

$$(5.10) \qquad w'(\xi, \gamma) \rightarrow \bar{w}'(\xi) \quad \text{in} \quad C^0(-\infty, \bar{\xi}] \quad \text{as} \quad \gamma \rightarrow \infty.$$

Because of (5.4), we see by (5.9) that $|w'(\xi, \gamma)|$ is uniformly bounded in each bounded interval. Since $w'' = -\alpha w' + w - w^p + m^2 e^{2m(\xi - \tau(\gamma))} w$, $|w''(\xi, \gamma)|$ is also uniformly bounded in each bounded interval. By Arzelà-Ascoli theorem we see that $w'(\xi, \gamma)$ uniformly converges in a bounded interval as $\gamma \rightarrow \infty$. Since the solution of (5.5) is unique, the limit function is $\bar{w}'(\xi)$. Hence

$$(5.11) \qquad w'(\xi, \gamma) \rightarrow \bar{w}'(\xi) \quad \text{in} \quad C_{loc}^0(-\infty, \bar{\xi}].$$

By the same argument as in the case $w'(-\infty)$ we can show that $\bar{w}'(-\infty) = 0$. Therefore, there is ξ_0 such that

$$(5.12) \quad |w'(\xi, \gamma) - \bar{w}'(\xi)| < \varepsilon \quad (-\infty < \xi < \xi_0).$$

Because of (5.11) and (5.12), (5.10) holds. The proof is complete. \square

Since $(\bar{w}, \bar{w}') \rightarrow (1, 0)$ ($\xi \rightarrow \infty$), it follows from Lemma 5.3 that (y, z) approaches $(1, 0)$ as $\gamma \rightarrow \infty$ along $t = \bar{\xi} - \tau(\gamma)$ provided that $\bar{\xi}$ is large enough. In the next lemma we show that once near $(1, 0)$, the orbit (y, z) remains near $(1, 0)$ until t reaches some T , which is large and negative, but independent of γ . We set

$$\Gamma_\rho := \{(y, z); E(y, z) < E(1, 0) + \rho\}, \quad \rho > 0,$$

where E is defined by (2.8).

Lemma 5.4. *For every $\varepsilon > 0$, there exists $T_\varepsilon (< t^*)$ such that the following holds: If there is $t_0 (< T_\varepsilon)$ such that $(y(t_0), z(t_0)) \in \Gamma_\varepsilon$, then*

$$(y(t), z(t)) \in \Gamma_{2\varepsilon} \quad (t_0 \leq t \leq T_\varepsilon).$$

Proof. Let $E(t) := E(y(t), z(t))$. We have

$$(5.13) \quad E'(t) = -\alpha z^2 + m^2 e^{2mt} y z \leq \frac{m^4}{4\alpha} e^{4mt} y^2.$$

Hence, for $t > t_0$,

$$(5.14) \quad E(t) \leq E(t_0) + \frac{m^4}{4\alpha} \int_{t_0}^t e^{4m\zeta} y(\zeta)^2 d\zeta.$$

However, there is $K > 0$ such that

$$(5.15) \quad E(y, z) \geq -K + y^2.$$

Therefore,

$$(5.16) \quad E(t) \leq E(t_0) + \frac{m^4}{4\alpha} \left(\frac{K}{4m} e^{4mt} + \int_{t_0}^t e^{4m\zeta} E(\zeta) d\zeta \right).$$

We choose T_ε so that

$$\frac{m^4}{4\alpha} \left\{ \frac{K}{4m} + \frac{1}{4m} (E(1, 0) + 2\varepsilon) \right\} e^{4mT_\varepsilon} < \varepsilon.$$

Let $t_0 (< T_\varepsilon)$ and define

$$t_1 := \sup\{t > t_0; E(t) < E(1, 0) + 2\varepsilon\}.$$

We show by contradiction that $t_1 > T_\varepsilon$. Suppose the contrary, i.e., $t_1 \leq T_\varepsilon$. Then by (5.16) and the definition of t_1 we have

$$\begin{aligned} E(t_1) &\leq E(t_0) + \frac{m^4}{4\alpha} \left(\frac{K}{4m} e^{4mt} + \int_{t_0}^t e^{4m\zeta} E(\zeta) d\zeta \right) \\ &< E(t_0) + \varepsilon \\ &\leq E(1, 0) + \varepsilon + \varepsilon \\ &= E(1, 0) + 2\varepsilon, \end{aligned}$$

which contradicts to the definition of t_1 . Hence, $t_1 > T_\varepsilon$, which indicates that $(y, z) \in \Gamma_{2\varepsilon} (t_0 \leq t \leq T_\varepsilon)$. \square

Lemma 5.5. *Suppose that $\gamma > 0$ is large. There exists a constant $M > 0$ independent of $\gamma > 0$ such that*

$$|y(t)| + |z(t)| \leq M \quad (t \leq t_n^*).$$

Proof. By the same argument as in the proof of Lemma 5.3 we show that $y'(-\infty) = 0$. Let $\tilde{E} := \tilde{E}(y(t), z(t))$. Since $\tilde{E}(t) \leq \tilde{E}(-\infty) = 0$, hence

$$(5.17) \quad \frac{z(t)^2}{2} - \frac{\beta(t)}{2} y(t)^2 + \frac{y(t)^{p+1}}{p+1} \leq 0 \quad \text{for } t \in \mathbb{R}.$$

When $t \in (-\infty, t_n^*]$, $\beta(t)$ is bounded, hence there is $K > 0$ such that

$$(5.18) \quad -\frac{\beta(t)}{2} y^2 + \frac{y^{p+1}}{p+1} \geq -\frac{K}{2} + \frac{y^2}{2} \quad (-\infty < t \leq t_n^*).$$

We see by (5.17) and (5.18) that for $t \in (-\infty, t_n^*]$,

$$\frac{z(t)^2}{2} + \frac{y(t)^2}{2} - \frac{K}{2} \leq \frac{z(t)^2}{2} - \frac{\beta(t)}{2} y(t)^2 + \frac{y(t)^{p+1}}{p+1} \leq 0,$$

hence $y(t)^2 + z(t)^2 \leq K$. The conclusion holds. \square

Proof of Theorem 5.1. Let (y_j, z_j) be a solution of (4.31) such that $e^{-m\theta t} y(t) \rightarrow \frac{\gamma_j}{A}$ ($t \rightarrow -\infty$). By Lemma 5.5, the sequence $\{(y_j, z_j)\}$ is uniformly bounded in $(-\infty, t_n^*]$. Since (y_j, z_j) is a solution of (4.31), the sequence $\{(y_j, z_j)\}$ is uniformly bounded in $(C^2(-\infty, t_n^*))^2$. By Arzelà-Ascoli theorem we can extract a subsequence, which is still denoted by $\{(y_j, z_j)\}$, converges in $(C^1(I))^2$ to some function (\bar{y}, \bar{z}) , where I is an arbitrary compact subset of $(-\infty, t_n^*]$. By taking the limit, we find that (\bar{y}, \bar{z}) satisfies (4.31).

It remains to show that

$$(5.19) \quad \bar{y}(t) \rightarrow 1 \quad \text{as } t \rightarrow -\infty,$$

for then \bar{y} is the solution of the problem

$$\begin{cases} y'' + \alpha y' - y + y^p - m^2 e^{2mt} y = 0, & -\infty < t < t_n^*, \\ y(-\infty) = 1. \end{cases}$$

We have shown in Lemma 4.1 that this problem has a unique solution $y^*(t)$. Thus, $\bar{y}(t) \equiv y^*(t)$ ($-\infty < t \leq t_n^*$). Because of this uniqueness, the entire sequence $\{y_j\}$ converges to y^* . Thus, as $j \rightarrow \infty$,

$$y_j \rightarrow y^* \quad \text{and} \quad y_j' \rightarrow (y^*)' \quad \text{in} \quad C_{loc}^0(-\infty, t_n^*].$$

Hence, the first two convergences of the statement of the theorem are obtained. The third convergence holds, since u_j satisfies (1.7).

We show (5.19) by contradiction argument. Suppose that (5.19) does not hold. Then there exists a sequence $\{t_k\}$ such that $t_k \rightarrow -\infty$ as $k \rightarrow \infty$, and a constant $\delta > 0$ so that

$$(5.20) \quad (\bar{y}(t_k), \bar{z}(t_k)) \notin \Gamma_\delta \quad \text{for all} \quad k \geq 1.$$

Let $\varepsilon := \frac{\delta}{4}$. Then by Lemma 5.3, there exist $\bar{\xi} \in \mathbb{R}$ and $j_0 \in \mathbb{N}$ such that

$$(y_j(\bar{\xi} - \tau(\gamma_j)), z_j(\bar{\xi} - \tau(\gamma_j))) \in \Gamma_\varepsilon \quad \text{if} \quad j > j_0.$$

By Lemma 5.4, this implies that if $j > j_0$, then

$$(y_j(t), z_j(t)) \in \Gamma_{2\varepsilon} \subset \Gamma_\delta \quad \text{in} \quad (\bar{\xi} - \tau(\gamma_j), T_\varepsilon),$$

where T_ε depends only on ε . By choosing j large enough, the interval $(\bar{\xi} - \tau(\gamma_j), T_\varepsilon)$ can be made to include an element of the sequence $\{t_k\}$. We obtain a contradiction, because of (5.20). \square

We are in a position to prove Theorem 1.1.

Proof of Theorem 1.1. The existence of $u^*(s)$ near $s = 0$ follows from Lemma 4.1. Lemma 4.4 says that u^* is positive and it is defined in $s \in (0, \infty)$. It follows from Lemma 4.8 that u^* oscillates around 1. The convergence property follows from Theorem 5.1. \square

5.2. Convergence to λ_n^* . Using Theorem 5.1, we prove the following:

Theorem 5.6. *Let $\{\lambda_n(\gamma)\}_{\gamma>1}$ be as in Subsection 3.3, and let λ_n^* be as in Theorem A. Then*

$$\lambda_n(\gamma) \rightarrow \lambda_n^* \quad \text{as} \quad \gamma \rightarrow \infty.$$

Proof. Let \tilde{s}_1^* be the first positive zero of $u^*(s) - \left(\frac{p+1}{2}\right)^{\frac{1}{p-1}}$, and let $I_1 := (0, \frac{\tilde{s}_1^*}{2})$. Let s_1 be the first positive critical point of $u(s, \gamma)$. We show by contradiction that if $\gamma > 0$ is large, then $u(s, \gamma)$ does not have a critical point in I_1 . Suppose that $s_1 \in I_1$. Then, s_1 should be a local minimum point of $u(s, \gamma)$, hence by (1.7), $u(s_1, \gamma) < 1$. Then the corresponding orbit in the (u, u_s) -plane is in $\{E \leq 0\}$ at $s = s_1$, and it cannot go out when $s > s_1$. This contradicts to Theorem 5.1, because $u(s, \gamma) \rightarrow u^*(s)$ in $C^0[\frac{\tilde{s}_1^*}{2}, \tilde{s}_1^*]$. We have shown that u does not have a critical point in I_1 .

Let s_n^* be the n -th positive critical point of $u^*(s)$. Let $\delta > 0$ be small. Because of Theorem 5.1, $u(s, \gamma) \rightarrow u^*(s)$ in $C^1[\frac{\tilde{s}_1^*}{2}, s_n^* + \delta]$ as $\gamma \rightarrow \infty$. Since each zero of u_s^* is simple, $s_n \rightarrow s_n^*$ ($\gamma \rightarrow \infty$), where s_n is the n -th positive critical point of $u(s, \gamma)$. Since $\sqrt{\lambda_n(\gamma)} = s_n$ and $\sqrt{\lambda_n^*} = s_n^*$, we obtain the conclusion. \square

6. OSCILLATION OF $\lambda_n(\gamma)$

Let $u(s, \gamma)$ be the solution of (1.7), and let (λ_n^*, u^*) be a singular solution of (1.14). Let $\{\lambda_n(\gamma)\}_{\gamma > 1}$ be as given in Subsection 3.3. The goal of this section is to prove the following:

Theorem 6.1. *Suppose that $p_S < p < p_{JL}$. $\lambda_n(\gamma)$ oscillates around λ_n^* infinitely many times as $\gamma \rightarrow \infty$.*

Proof. Let $I := (0, \sqrt{\lambda_n^*}]$. We define $\tilde{u}(\rho, \gamma) := u(s, \gamma)/\gamma$, $\tilde{u}^*(\rho, \gamma) := u^*(s)/\gamma$, and $\rho := \gamma^{\frac{p-1}{2}} s$. We show that

(6.1) for each $M \geq 1$, there are $\rho_0 \in [0, 1]$ and $\gamma_0 > 0$ such that

$$\mathcal{Z}_{(0, \rho_0)}[\tilde{u}^*(\cdot, \gamma) - \tilde{u}(\cdot, \gamma)] \geq M \quad \text{for } \gamma \geq \gamma_0.$$

The function $\tilde{u}(\rho)$ satisfies (1.15). Since the energy

$$\tilde{G}(\tilde{u}, \tilde{u}_\rho) = \frac{\tilde{u}_\rho^2}{2} - \frac{\tilde{u}^2}{2\gamma^{p-1}} + \frac{\tilde{u}^{p+1}}{p+1}$$

decreases, $\tilde{G}(\tilde{u}, \tilde{u}_\rho) \leq \tilde{G}(1, 0) < 0$, hence $\tilde{u} \leq \left(\frac{p+1}{2\gamma^{p-1}}\right)^{\frac{1}{p-1}}$. Since $\tilde{u}/\gamma^{p-1} \rightarrow 0$ as $\gamma \rightarrow \infty$,

(6.2) $\tilde{u}(\rho, \gamma) \rightarrow \bar{u}(\rho, 1)$ in $C_{loc}^2(0, \infty) \cap C_{loc}^0[0, \infty)$ ($\gamma \rightarrow \infty$),

where $\bar{u}(\cdot, 1)$ is the solution of (1.16) with $\gamma = 1$. We apply the same change of variables to the singular solution $u^*(s)$. We have

$$\begin{aligned} \tilde{u}^*(\rho, \gamma) &:= \frac{1}{\gamma} A \left(\frac{\rho}{\gamma^{\frac{p-1}{2}}} \right)^{-\theta} (1 + o(1)) \quad (s \downarrow 0) \\ (6.3) \quad &= A\rho^{-\theta}(1 + o(1)) \quad (\rho\gamma^{-\frac{p-1}{2}} \downarrow 0). \end{aligned}$$

If ρ is in a compact interval in \mathbb{R}_+ , then $\rho\gamma^{-\frac{p-1}{2}}$ uniformly converges to 0 on the interval as $\gamma \rightarrow \infty$. Therefore, (6.3) indicates that

$$(6.4) \quad \tilde{u}^*(\rho, \gamma) \rightarrow \bar{u}^*(\rho) \quad \text{in } C_{loc}^2(0, \infty) \quad (\gamma \rightarrow \infty),$$

where \bar{u}^* is given by (2.2) which is a singular solution of the equation in (1.16). Because of Proposition 2.1, $\mathcal{Z}_{[0, \infty)}[\bar{u}^*(\cdot) - \bar{u}(\cdot, 1)] = \infty$. Therefore, (6.2) and (6.4) indicate that (6.1) holds.

Since $0 < \rho < \rho_0$ is equivalent to $0 < s < \rho_0\gamma^{-\frac{p-1}{2}}$ and $\rho_0\gamma^{-\frac{p-1}{2}} \downarrow 0$ ($\gamma \rightarrow \infty$), (6.1) implies that

$$(6.5) \quad \mathcal{Z}_I[u^*(\cdot) - u(\cdot, \gamma)] \rightarrow \infty \quad (\gamma \rightarrow \infty).$$

Since each zero of $u^*(\cdot) - u(\cdot, \gamma)$ is simple, the number of the zeros in I is preserved provided that another zero does not enter I from ∂I . Since $u^*(0) - u(0, \gamma) = \infty$, a zero cannot enter I from $s = 0$. Since (6.5) says that the number of the zeros increases, a simple zero enter I from $s = \sqrt{\lambda_n^*}$ infinitely many times as $\gamma \rightarrow \infty$. Therefore, there exists a sequence of large numbers $\{\gamma_j\}_{j=1}^\infty$ ($\gamma_1 < \gamma_2 < \dots \rightarrow \infty$) such that $u(\sqrt{\lambda_n^*}, \gamma_j) = u^*(\sqrt{\lambda_n^*})$ and

$$\partial_s u(\sqrt{\lambda_n^*}, \gamma_j) \begin{cases} < 0, & (j \in \{1, 3, 5, \dots\}), \\ > 0, & (j \in \{2, 4, 6, \dots\}). \end{cases}$$

Note that $\partial_s u(\sqrt{\lambda_n^*}, \gamma_j) \neq 0$, because of the simplicity of the zero. We consider only the case where n is odd, since the proof of the case where n is even is similar to this case. In the proof of Theorem 5.6 we already saw that $u(s, \gamma_j)$ does not have a critical point in $(0, \frac{\bar{s}_1}{2})$. $\sqrt{\lambda_n^*}$ is a local minimum point of u^* . Because of Theorem 5.1, $u(s, \gamma_j)$ is close to $u^*(s)$ in $C^2[\frac{\bar{s}_1}{2}, \sqrt{\lambda_n^*}]$. We easily see that the n -th positive critical point of $u(s, \gamma)$, which is $\sqrt{\lambda_n(\gamma_j)}$, is close to $\sqrt{\lambda_n^*}$. We consider the case where $\partial_s u(\sqrt{\lambda_n^*}, \gamma_j) < 0$. We show that $\lambda_n(\gamma_j) > \lambda_n^*$. Suppose the contrary, i.e., $\lambda_n(\gamma_j) < \lambda_n^*$. Since $\partial_s u(\sqrt{\lambda_n^*}, \gamma_j) < 0$ and $s = \sqrt{\lambda_n^*}$ is a local minimum point of $u(\sqrt{\lambda_n(\gamma_j)}, \gamma_j)$, there is a local maximum point

$s^* \in (\sqrt{\lambda_n(\gamma_j)}, \sqrt{\lambda_n^*})$. Since s^* is a local maximum point, $u(s^*, \gamma_j) > 1$ which contradicts to Theorem 5.1. Thus, $\lambda_n(\gamma_j) > \lambda_n^*$.

When $\partial_s u(\sqrt{\lambda_n^*}, \gamma_j) > 0$, we suppose the contrary, i.e., $\lambda_n(\gamma_j) > \lambda_n^*$. Since $\sqrt{\lambda_n(\gamma_j)}$ is a local minimum point, there is a local maximum point $s^* \in (\sqrt{\lambda_n(\gamma_j)}, \sqrt{\lambda_n^*})$ and $u(s^*, \gamma_j) > 1$. We obtain a contradiction, because of Theorem 5.1. Thus, $\lambda_n(\gamma_j) < \lambda_n^*$.

We have proved (1.18), which implies that $\lambda_n(\gamma)$ oscillates around λ_n^* infinitely many times as $\gamma \rightarrow \infty$. \square

7. BOUNDARY CONCENTRATING SOLUTION

We construct a boundary concentrating solution which is radially increasing when λ is large. We show that this solution belongs to \mathcal{C}_1 . We use different notation only in this section.

7.1. One-dimensional problem. We consider the one-dimensional problem

$$(7.1) \quad \begin{cases} \varepsilon^2 w_{xx} + f(w) = 0, & 0 < x < 1, \\ w_x(0) = w_x(1) = 0. \end{cases}$$

Let $\bar{w}(y) := w(x)$ and $y := x/\varepsilon$. The function $\bar{w}(y)$ satisfies

$$(7.2) \quad \begin{cases} \bar{w}_{yy} + f(\bar{w}) = 0, & 0 < y < d_\varepsilon, \\ \bar{w}_y(0) = \bar{w}_y(d_\varepsilon) = 0, \end{cases}$$

where $d_\varepsilon := 1/\varepsilon$.

For $p > 1$ the system of equations for (\bar{w}, \bar{z}) ($\bar{z} := \bar{w}_y$) in the phase plane

$$\begin{cases} \bar{w}_y = \bar{z} \\ \bar{z}_y = -f(\bar{w}) \end{cases}$$

has a saddle point at $(0, 0)$ and a center $(1, 0)$. There is a unique homoclinic solution around the center connecting the saddle to itself. This homoclinic solution can be written explicitly as

$$w^*(y) := \left(\frac{p+1}{2} \right)^{\frac{1}{p-1}} \left(\cosh \left(\frac{p-1}{2} y \right) \right)^{-\frac{2}{p-1}}.$$

From the phase portrait for (\bar{w}, \bar{z}) it is clear that all the orbits on $\{\bar{w} > 0\}$ satisfying the Neumann boundary conditions are inside the region enclosed by the homoclinic orbit and that every orbit in this region is periodic one. Hence, \bar{w} is a solution of (7.2) if and only if an integral multiple of its half period is equal to the interval length d_ε .

Now, we find an increasing solution. Let $\bar{w}(y)$ be an increasing solution that has maximum α and minimum β . Then,

$$0 < \beta < 1 < \alpha < \bar{\alpha} := \left(\frac{p+1}{2}\right)^{\frac{1}{p-1}}.$$

Multiplying (7.2) by \bar{w}_y and integrating it, we have

$$\bar{w}_y^2 = F(\bar{w}) - F(\beta), \quad F(\alpha) = F(\beta), \quad \text{and} \quad F(\bar{w}) = \bar{w}^2 - \frac{2}{p+1}\bar{w}^{p+1}.$$

The half period is given by the integral

$$(7.3) \quad T(\alpha) := \int_{\beta}^{\alpha} \frac{d\bar{w}}{\sqrt{F(\bar{w}) - F(\beta)}}.$$

Thus, \bar{w} is an increasing solution of (7.2) if and only if $T(\alpha) = d_\varepsilon$. The integral (7.3) was studied by De Groen and Karadzhov [8]. Among other things, they obtained

Proposition 7.1. *There is a small $\varepsilon_0 > 0$ such that the problem (7.1) has a smooth curve of increasing solutions $\{w(x; \varepsilon)\}_{0 < \varepsilon < \varepsilon_0}$, which can be described as a graph of ε , satisfying the following: For $\delta > 0$, there exists $\varepsilon_1 > 0$ such that, if $0 < \varepsilon < \varepsilon_1$, then*

$$(7.4) \quad |\tilde{w}(s; \varepsilon) - w^*(s)| < \delta \quad \text{for } s \in [0, d_\varepsilon],$$

where $\tilde{w}(s; \varepsilon) := w(x; \varepsilon)$ ($x = 1 - \varepsilon s$). Moreover, the first two eigenvalues of the eigenvalue problem

$$\varepsilon^2 \phi_{xx} + f'(w)\phi = \kappa\phi \quad \text{in } (0, 1), \quad \phi_x = 0 \quad \text{at } x = 0, 1$$

are

$$\begin{aligned} \kappa_0(\varepsilon) &= \frac{(p-1)(p+3)}{4} + O\left(e^{-\frac{2}{\varepsilon}}\right), \\ \kappa_1(\varepsilon) &\begin{cases} = -\frac{(p-1)(5-p)}{4} + O\left(e^{-\frac{3-p}{\varepsilon}}\right) & (1 < p < 3), \\ \leq -1 + O\left(e^{-\frac{2}{\varepsilon}}\right) & (3 \leq p). \end{cases} \end{aligned}$$

In particular, if $\varepsilon > 0$ is small, then $\kappa_1(\varepsilon) < 0 < \kappa_0(\varepsilon)$ and $w(x; \varepsilon)$ is nondegenerate.

Using (7.4), we immediately have

Corollary 7.2. *Let $w(x; \varepsilon)$ be the increasing solution. Then, w satisfies $(\varepsilon w_x)^2 - F(w) = -F(w(0; \varepsilon))$ for $x \in (0, 1)$, and $-F(w(0; \varepsilon)) \rightarrow 0$ ($\varepsilon \downarrow 0$). Moreover, $F(w(\cdot; \varepsilon)) \xrightarrow{\varepsilon \downarrow 0} 0$ pointwisely in $[0, 1]$.*

We recall some known results of the “limiting” operator $L^* := \partial_{ss} + f'(w^*(s))$ on \mathbb{R}_+ with the Neumann boundary condition. The operator L^* has a continuous spectrum $(-\infty, -1]$ and may have discrete eigenvalues outside $(-\infty, -1]$ ([16, p. 140]). We study the nonnegative eigenvalues of L^* .

Proposition 7.3. *The eigenvalue problem*

$$(7.5) \quad L^* \phi^* = \kappa^* \phi^* \text{ in } \mathbb{R}_+, \quad \phi_s^*(0) = 0, \quad \phi^* \in H^1(\mathbb{R}_+)$$

has the unique nonnegative eigenvalue. Moreover, this eigenvalue is the first one which is simple, $\kappa_0^ := (p-1)(p+3)/4$, and the associated eigenfunction is $\phi_0^*(s) := (w^*)^{(p+1)/2}$.*

By direct calculation we see that

$$\phi_1^* := (w^*)^{\frac{3-p}{2}} - \frac{p+3}{2(p+1)} (w^*)^{\frac{p+1}{2}} \quad \text{and} \quad \kappa_1^* := -\frac{(p-1)(5-p)}{4}$$

satisfy

$$L^* \phi_1^* = \kappa_1^* \phi_1^* \text{ in } \mathbb{R}_+, \quad \partial_s \phi_1^*(0) = 0, \quad \phi_1^* \in H^1(\mathbb{R}_+) \quad (1 < p < 3).$$

It is known that if $1 < p < 3$, then κ_1^* is the second eigenvalue and that if $p \geq 3$, then L^* has only one eigenvalue above -1 . In particular, 0 is not an eigenvalue and L^* is invertible.

Corollary 7.4. *For each $p > 1$, there is $\delta > 0$ such that L^* has no eigenvalue in $[-\delta, \infty)$ except κ_0^* .*

7.2. Notation. From now on we assume $q \in (\max\{N/2, 2\}, N)$. Under this assumption the inclusion $W^{2,q}(B) \hookrightarrow C^0(B)$ is continuous ($q > N/2$) and $1/r \in L^q(B)$ ($q < N$). Let \mathcal{R} denote the set of the radial functions on B . Let $\mathbb{X} := \{u \in W^{2,q}(B) \cap \mathcal{R}; \partial_\nu u = 0 \text{ on } \partial B\}$ and $\mathbb{Y} := L^q(B) \cap \mathcal{R}$. By $\mathcal{B}(\mathbb{X}, \mathbb{Y})$ we denote the space of the bounded operators from \mathbb{X} to \mathbb{Y} equipped with the operator norm $\|\cdot\|_{\mathcal{B}(\mathbb{X}, \mathbb{Y})}$. In this section we work on \mathbb{X} and \mathbb{Y} and define $f(U) := -U + U|U|^{p-1}$ in order that f is defined in \mathbb{R} . The regular radial solutions of (1.3) satisfy

$$(7.6) \quad \begin{cases} \varepsilon^2 (\partial_{rr} + \frac{N-1}{r} \partial_r) U + f(U) = 0, & 0 < r < 1, \\ U_r(0) = U_r(1) = 0. \end{cases}$$

Let $w(r)$ be the increasing solution given in Proposition 7.1. We find a solution of (7.6) near $w(r)$. Substituting $U(r) = v(r) + w(r)$ into (7.6), we obtain the following equation of the error term v :

$$(7.7) \quad \mathcal{L}v + \mathcal{M}w + \mathcal{N}(v, w) = 0,$$

where

$$\begin{aligned} \mathcal{L}v &:= \varepsilon^2 \left(\partial_{rr} + \frac{N-1}{r} \partial_r \right) v + f'(w)v, \\ \mathcal{M}w &:= \varepsilon^2 \frac{N-1}{r} w_r, \\ \mathcal{N}(v, w) &:= f(v+w) - f(w) - f'(w)v. \end{aligned}$$

\mathcal{L} is an approximation of the linearized operator $\varepsilon^2 \left(\partial_{rr} + \frac{N-1}{r} \partial_r \right) + f'(U)$. We easily see that $\mathcal{L} \in \mathcal{B}(\mathbb{X}, \mathbb{Y})$ and that $\mathcal{N}(\cdot, w)$ is a nonlinear mapping from \mathbb{X} to \mathbb{Y} , since the inclusion $\mathbb{X} \hookrightarrow C^0(B)$ is continuous.

7.3. Apriori estimates. We use the following apriori estimate in order to use the dominated convergence theorem.

Proposition 7.5. *Let Ω be a bounded domain. Let $\phi \in C^0(\Omega) \cap H^1(\Omega)$. If*

$$\varepsilon^2 \Delta \phi - V(x)\phi = 0 \quad \text{in } \Omega,$$

and if $V(x) > C_0 > 0$ in Ω , then there is $C_1 > 0$ independent of ε such that

$$|\phi(x)| \leq 2 \|\phi\|_\infty \exp\left(-\frac{C_1 \text{dist}(x, \partial\Omega)}{\varepsilon}\right),$$

where $\text{dist}(x, \partial\Omega)$ is the distance from x to $\partial\Omega$.

When $\phi \in C^2(\Omega)$, Proposition 7.5 was proved in [10, Lemma 4.2] for a general operator. The proof of [10, Lemma 4.2] works in the generalized sense, hence Proposition 7.5 holds. This proposition was also used in [27].

Lemma 7.6. *Let $(\kappa_\varepsilon, \phi) \in \mathbb{R} \times \mathbb{X}$ ($\|\phi\|_\infty = 1$) be the eigenpair of*

$$\begin{cases} \mathcal{L}\phi = \kappa_\varepsilon \phi & \text{in } B, \\ \partial_\nu \phi = 0 & \text{on } \partial B. \end{cases}$$

If there is a small $\delta_0 > 0$ such that $|\kappa_\varepsilon| < \delta_0$ for small $\varepsilon > 0$, then there are $C_0 > 0$ and $C_1 > 0$ independent of $\varepsilon > 0$ such that

$$|\phi(r)| \leq C_0 e^{-C_1(1-r)/\varepsilon} \quad (0 \leq r \leq 1).$$

Proof. We see by Proposition 7.1 that there are $C_2 > 0$ and $\delta_1 > 0$ such that $f'(w) < -C_2 < 0$ ($0 \leq r \leq 1 - \delta_1\varepsilon$). The function ϕ satisfies

$$\varepsilon^2 \left(\partial_{rr} + \frac{N-1}{r} \partial_r \right) \phi + f'(w)\phi - \kappa_\varepsilon \phi = 0.$$

Since $\delta_0 > 0$ is small, there is $C_3 > 0$ such that $f'(w) - \kappa_\varepsilon < f'(w) + \delta_0 < -C_3 < 0$ ($0 \leq r \leq 1 - \delta_1\varepsilon$). It follows from Proposition 7.5 that there are $C_4 > 0$ and $C_5 > 0$ such that $\text{dist}(r, \partial B) = 1 - r - \delta_1\varepsilon$ and

$$|\phi(r)| \leq C_4 e^{-C_5(1-r-\varepsilon\delta_1)/\varepsilon} = C_4 e^{\delta_1 C_5} e^{-C_5(1-r)/\varepsilon} \quad (0 \leq r \leq 1 - \delta_1\varepsilon).$$

If we take C_4 large enough, then this inequality holds for $r \in [0, 1]$, because $\|\phi\|_\infty = 1 < C_4 \leq C_4 e^{-C_5(1-r-\varepsilon\delta_1)/\varepsilon}$ ($1 - \delta_1\varepsilon < r < 1$). The proof is complete. \square

7.4. Invertibility of \mathcal{L} . The main technical lemma in Section 7 is the invertibility of \mathcal{L} .

Lemma 7.7. *The operator $\mathcal{L} \in \mathcal{B}(\mathbb{X}, \mathbb{Y})$ is invertible, and there is $C > 0$ independent of small $\varepsilon > 0$ such that $\|\mathcal{L}^{-1}\|_{\mathcal{B}(\mathbb{Y}, \mathbb{X})} \leq C$.*

Proof. It is enough to show that there is a small $\delta > 0$ such that \mathcal{L} has no eigenvalue in $[-\delta, \delta]$. We show this by contradiction.

Suppose the contrary, i.e., there is a small $\delta > 0$ such that \mathcal{L} has an eigenvalue κ_ε in $[-\delta, \delta]$ for small $\varepsilon > 0$. Then there is an eigenfunction ϕ such that

$$\begin{cases} \varepsilon^2 \Delta \phi + f'(w)\phi = \kappa_\varepsilon \phi & \text{in } B, \\ \partial_\nu \phi = 0 & \text{on } \partial B. \end{cases}$$

Without loss of generality we can assume $\|\phi\|_\infty = 1$. Because of Proposition 7.1, $f'(w) - \kappa_\varepsilon < -\delta_0 < 0$ ($0 \leq r \leq 1 - \delta_1\varepsilon$). By Lemma 7.6 we have

$$|\phi(r)| \leq C_0 e^{-C_1(1-r)/\varepsilon} \quad (0 \leq r \leq 1).$$

We let $\tilde{\phi}(s) := \phi(r)$, $\tilde{w}(s) := w(r)$, and $r = 1 - \varepsilon s$. Moreover, we set $\bar{\phi}(s) := \tilde{\phi}(|s|)$ and $\bar{w}(s) := \tilde{w}(|s|)$. Then $\bar{\phi}$ satisfies

$$(7.8) \quad \begin{cases} \bar{\phi}_{ss} + a(s)\bar{\phi}_s + f'(\bar{w}(s))\bar{\phi} = 0, & -\frac{1}{\varepsilon} < s < \frac{1}{\varepsilon}, \\ \bar{\phi}_s(0) = \bar{\phi}_s(\pm \frac{1}{\varepsilon}) = 0, \\ \|\bar{\phi}\|_\infty = 1, \end{cases}$$

where

$$a(s) := -\text{sign}(s) \frac{(N-1)\varepsilon^2}{1 - \varepsilon|s|}.$$

Let $R > 0$. If $\varepsilon > 0$ is small, then $2R < 1/\varepsilon$, $a(s) \in L^\infty(-2R, 2R)$. By [13, Theorem 9.11] we see that $\|\bar{\phi}\|_{H^2(-R, R)} \leq C_2$. Since the inclusion $H^2(-R, R) \hookrightarrow C^{1, \gamma}(-R, R)$ ($0 < \gamma < 1/2$) is continuous, $\|\bar{\phi}\|_{C^{1, \gamma}(-R, R)} \leq C_3$. By the Arzelá-Ascoli theorem we can choose a subsequence $\{\bar{\phi}\}_{\varepsilon > 0}$ such that the following holds: There is $\bar{\phi}_R \in C^1(-R, R)$ such that $\bar{\phi} \xrightarrow{\varepsilon \downarrow 0} \bar{\phi}_R$ in $C^1(-R, R)$. It follows from (7.8) that $\bar{\phi}_{ss}$ converges in $C^0(-R, R)$ as $\varepsilon \downarrow 0$. Since the operator ∂_{ss} that is defined on $\{u \in C^2(-R, R); u_s(0) = 0\}$ is a closed operator in $C^1(-R, R)$, $\bar{\phi}_R \in C^2$ and $\bar{\phi}_{ss} \xrightarrow{\varepsilon \downarrow 0} \partial_{ss}\bar{\phi}_R$ in $C^0(-R, R)$. Let $\{R_j\}_{j \geq 1}$ ($0 < R_1 < R_2 < \dots \rightarrow \infty$) be a sequence diverging to ∞ . For each fixed R_j , there is $\varepsilon > 0$ such that $[0, R_j) \subset (-1/\varepsilon, 1/\varepsilon)$. Using the expanding domains $\{[0, R_j)\}_{j \geq 1}$ and a diagonal argument, we see that there is $\bar{\phi}^* \in C^2(\mathbb{R})$ and $\kappa^* \in [-\delta, \delta]$ such that $\bar{\phi} \xrightarrow{\varepsilon \downarrow 0} \bar{\phi}^*$ in $C_{loc}^2(\mathbb{R})$ and $\kappa_\varepsilon \rightarrow \kappa^*$. We set $\phi^*(s) := \bar{\phi}^*(s)$ ($s \in \overline{\mathbb{R}_+}$). Since, for each $\varphi \in C_0^1(\overline{\mathbb{R}_+})$,

$$\int_0^{1/\varepsilon} \left(\tilde{\phi}_{ss} - \frac{(N-1)\varepsilon^2}{1-\varepsilon s} \tilde{\phi}_s + f'(\tilde{w})\tilde{\phi} - \kappa_\varepsilon \tilde{\phi} \right) \varphi ds = 0,$$

we have

$$\int_{\overline{\mathbb{R}_+}} (\phi_{ss}^* + f'(w^*)\phi^* - \kappa^*\phi^*) \varphi dx = 0, \quad \|\phi^*\|_\infty = 1, \quad \phi_s^*(0) = 0.$$

Since $\|\phi^*\|_\infty = 1$, $\phi^* \not\equiv 0$.

We will show that $\phi^* \in H^1(\mathbb{R}_+)$. Multiplying $\mathcal{L}\phi = \kappa_\varepsilon\phi$ by $r^{N-1}\phi$ and integrating it, we have

$$\int_0^1 (\varepsilon^2 \phi_r^2 - f'(w)\phi^2) r^{N-1} dr = - \int_0^1 \kappa_\varepsilon \phi^2 r^{N-1} dr.$$

Making the change of variables $r = 1 - \varepsilon s$, we have

$$\int_0^{1/\varepsilon} (\tilde{\phi}_s^2 + \tilde{\phi}^2) (1 - \varepsilon s)^{N-1} ds = \int_0^{1/\varepsilon} (p\tilde{w}^{p-1} - \kappa_\varepsilon) \tilde{\phi}^2 (1 - \varepsilon s)^{N-1} ds.$$

By Lemma 7.6 we have $|\tilde{\phi}(s)| \leq C_0 e^{-C_1 s}$. The right-hand side is bounded uniformly in ε , hence

$$\left| \int_0^{1/\varepsilon} (p\tilde{w}^{p-1} - \kappa_\varepsilon) \tilde{\phi}^2 (1 - \varepsilon s)^{N-1} ds \right| \leq C_2.$$

Using Fatou's lemma, we have

$$\begin{aligned} \int_{\mathbb{R}_+} \{(\phi_s^*)^2 + (\phi^*)^2\} ds &= \int_{\mathbb{R}_+} \liminf_{\varepsilon \downarrow 0} \chi_{[0,1/\varepsilon]} \left(\tilde{\phi}_s^2 + \tilde{\phi}^2 \right) (1 - \varepsilon s)^{N-1} ds \\ &\leq \liminf_{\varepsilon \downarrow 0} \int_0^{1/\varepsilon} \left(\tilde{\phi}_s^2 + \tilde{\phi}^2 \right) (1 - \varepsilon s)^{N-1} ds \leq C_2, \end{aligned}$$

where $\chi_{[0,1/\varepsilon]}$ is the characteristic function. Thus $\phi^* \in H^1(\mathbb{R}_+)$. Therefore (κ^*, ϕ^*) is an eigenpair of (7.5). Since (7.5) has no eigenvalue in $[-\delta, \delta]$ (Corollary 7.4), we obtain a contradiction. \square

7.5. Contraction mapping. Because of Lemma 7.7, \mathcal{L} is invertible provided that $\varepsilon > 0$ is small. The equation (7.7) can be transformed to

$$v = \mathcal{T}(v),$$

where

$$\mathcal{T}(v) := -\mathcal{L}^{-1}[\mathcal{M}w] - \mathcal{L}^{-1}[\mathcal{N}(v, w)].$$

In order to show that \mathcal{T} is a contraction mapping on a small ball in \mathbb{X} we show that $\|\mathcal{L}^{-1}[\mathcal{M}w]\|_{\mathbb{X}}$ is small.

Lemma 7.8. $\|\mathcal{L}^{-1}[\mathcal{M}w]\|_{\mathbb{X}} = o(\varepsilon)$.

Proof. By Lemma 7.7 we have

$$\|\mathcal{L}^{-1}[\mathcal{M}w]\|_{\mathbb{X}} \leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(\mathbb{X}, \mathbb{Y})} \|\mathcal{M}w\|_{\mathbb{Y}} \leq C \|\mathcal{M}w\|_{\mathbb{Y}}.$$

We show that $\|\mathcal{M}w\|_{\mathbb{Y}} = o(\varepsilon)$. Using Corollary 7.2, we have

$$\begin{aligned} \|\mathcal{M}w\|_{\mathbb{Y}}^q &= \left\| \varepsilon^2 \frac{N-1}{r} w_r \right\|_q^q \\ &= (N-1)^q \varepsilon^q \int_0^1 (\varepsilon w_r)^q r^{N-q-1} dr \\ &= (N-1)^q \varepsilon^q \int_0^1 (F(w(r)) - F(w(0)))^{\frac{q}{2}} r^{N-q-1} dr. \end{aligned}$$

Since $|F(w(r)) - F(w(0))| \leq C$ ($0 \leq r \leq 1$) and $|F(w(r)) - F(w(0))| \xrightarrow{\varepsilon \downarrow 0} 0$ pointwisely in $[0, 1]$, we see that $(F(w(r)) - F(w(0)))^{q/2} r^{N-q-1} \xrightarrow{\varepsilon \downarrow 0} 0$ pointwisely in $(0, 1]$ and that

$$|(F(w(r)) - F(w(0)))^{q/2} r^{N-q-1}| \leq C^{q/2} r^{N-q-1} \in L^1(0, 1).$$

Here, we use the condition $q < N$. The dominated convergence theorem says that

$$\int_0^1 (F(w(r)) - F(w(0)))^{\frac{q}{2}} r^{N-q-1} dr \rightarrow 0 \quad (\varepsilon \downarrow 0).$$

Since $\|\mathcal{M}w\|_{\mathbb{Y}}^q = \varepsilon^q \cdot o(1) = o(\varepsilon^q)$, the proof is complete. \square

Let $\mathbb{B}_\varepsilon := \{u \in \mathbb{X}; \|u\|_{\mathbb{X}} < \varepsilon\}$. We show that the Lipschitz constant of $\mathcal{N}(\cdot, w)$ in \mathbb{B}_ε is small.

Lemma 7.9. *Let $\varepsilon > 0$ be small. If $v_1, v_2 \in \mathbb{B}_\varepsilon$, then*

$$\|\mathcal{L}^{-1}[\mathcal{N}(v_1, w)] - \mathcal{L}^{-1}[\mathcal{N}(v_2, w)]\|_{\mathbb{X}} \leq o(1) \|v_1 - v_2\|_{\mathbb{X}} \quad (\varepsilon \downarrow 0).$$

In particular, the Lipschitz constant of $\mathcal{L}^{-1}[\mathcal{N}(\cdot, w)] : \mathbb{B}_\varepsilon \rightarrow \mathbb{X}$ is less than one.

Proof. By Lemma 7.7 we have

$$\begin{aligned} \|\mathcal{L}^{-1}[\mathcal{N}(v_1, w)] - \mathcal{L}^{-1}[\mathcal{N}(v_2, w)]\|_{\mathbb{X}} &\leq \|\mathcal{L}^{-1}\|_{\mathcal{B}(\mathbb{X}, \mathbb{Y})} \|\mathcal{N}(v_1, w) - \mathcal{N}(v_2, w)\|_{\mathbb{Y}} \\ &\leq C \|\mathcal{N}(v_1, w) - \mathcal{N}(v_2, w)\|_{\mathbb{Y}}. \end{aligned}$$

We show that $\|\mathcal{N}(v_1, w) - \mathcal{N}(v_2, w)\|_{\mathbb{Y}} \leq o(\varepsilon) \|v_1 - v_2\|_{\mathbb{X}}$ for $v_1, v_2 \in \mathbb{B}_\varepsilon$. We have

$$\begin{aligned} \|\mathcal{N}(v_1, w) - \mathcal{N}(v_2, w)\|_q &= \|f(v_1 + w) - f(v_2 + w) - f'(w)(v_1 - v_2)\|_q \\ &= \|f'(\tilde{w})(v_1 - v_2) - f'(w)(v_1 - v_2)\|_q \\ &\leq \|f'(\tilde{w}) - f'(w)\|_\infty \|v_1 - v_2\|_q. \end{aligned}$$

Here, $\tilde{w}(x)$ is a function such that

$$\min\{v_1(x) + w(x), v_2(x) + w(x)\} \leq \tilde{w}(x) \leq \max\{v_1(x) + w(x), v_2(x) + w(x)\},$$

hence $\|w(x) - \tilde{w}(x)\|_\infty \leq C\varepsilon$. We have $\|f'(\tilde{w}) - f'(w)\|_\infty = o(1)$ ($\varepsilon \downarrow 0$). Thus,

$$\begin{aligned} \|\mathcal{N}(v_1, w) - \mathcal{N}(v_2, w)\|_q &\leq o(1) \|v_1 - v_2\|_q \\ &\leq o(1)C \|v_1 - v_2\|_{\mathbb{X}}. \end{aligned}$$

The proof is complete. \square

Using Lemma 7.9, we will show that \mathcal{T} is a contraction mapping in \mathbb{B}_ε .

Lemma 7.10. *Let $\varepsilon > 0$ be small. Then $T(\mathbb{B}_\varepsilon) \subset \mathbb{B}_\varepsilon$. Moreover, the Lipschitz constant of $\mathcal{T} : \mathbb{B}_\varepsilon \rightarrow \mathbb{B}_\varepsilon$ is less than one, i.e., there is $\delta \in (0, 1)$ such that if $v_1, v_2 \in \mathbb{B}_\varepsilon$, then $\|\mathcal{T}(v_1) - \mathcal{T}(v_2)\|_{\mathbb{X}} \leq \delta \|v_1 - v_2\|_{\mathbb{X}}$.*

Proof. First, we will show that $T(\mathbb{B}_\varepsilon) \subset \mathbb{B}_\varepsilon$. By Lemma 7.9 we have that, for $v \in \mathbb{B}_\varepsilon$,

$$(7.9) \quad \begin{aligned} \|\mathcal{L}^{-1}[\mathcal{N}(v, w)]\|_{\mathbb{X}} &= \|\mathcal{L}^{-1}[\mathcal{N}(v, w)] - \mathcal{L}^{-1}[\mathcal{N}(0, w)]\|_{\mathbb{X}} \\ &\leq o(1) \|v - 0\|_{\mathbb{X}}, \end{aligned}$$

where we use Lemma 7.9. We have that, for $v \in \mathbb{B}_\varepsilon$,

$$\begin{aligned} \|\mathcal{T}(v)\|_{\mathbb{X}} &= \|\mathcal{L}^{-1}[\mathcal{M}w] + \mathcal{L}^{-1}[\mathcal{N}(v, w)]\|_{\mathbb{X}} \\ &\leq \|\mathcal{L}^{-1}[\mathcal{M}w]\|_{\mathbb{X}} + \|\mathcal{L}^{-1}[\mathcal{N}(v, w)]\|_{\mathbb{X}} \\ &\leq o(\varepsilon) + o(1) \|v\|_{\mathbb{X}} = o(\varepsilon), \end{aligned}$$

where we use Lemma 7.8 and (7.9). Thus, if $\varepsilon > 0$ is small, then, for all $v \in \mathbb{B}_\varepsilon$, $\mathcal{T}(v) \in \mathbb{B}_\varepsilon$.

Second, we show that the Lipschitz constant of $\mathcal{T}(\cdot)$ in \mathbb{B}_ε is $o(1)$. We have that, for $v_1, v_2 \in \mathbb{B}_\varepsilon$,

$$\begin{aligned} \|\mathcal{T}(v_1) - \mathcal{T}(v_2)\|_{\mathbb{X}} &= \|\mathcal{L}^{-1}[\mathcal{N}(v_1, w)] - \mathcal{L}^{-1}[\mathcal{N}(v_2, w)]\|_{\mathbb{X}} \\ &\leq o(1) \|v_1 - v_2\|_{\mathbb{X}}, \end{aligned}$$

where we use Lemma 7.9. Then the later part of the lemma holds. \square

Applying the contraction mapping theorem to \mathcal{T} which is defined in \mathbb{B}_ε , we obtain the following:

Corollary 7.11. *There is a large $\lambda_0 > 0$ such that (7.6) has a one-parameter family of positive solutions $\{(\lambda, U(r, \lambda))\}_{\lambda > \lambda_0}$ and $\|U - w\|_\infty < C\varepsilon (= C/\sqrt{\lambda})$.*

Proof. Since the condition $\|U - w\|_\infty < C\varepsilon$ does not guarantee the positivity of U , we have to check the positivity of U . We see by a priori estimate that U is a classical solution. If there is $r_0 \in [0, 1]$ such that $U(r_0) < 0$, then there is $r_1 \in [0, 1]$ such that $\min_{0 \leq r < 1} U(r) = U(r_1) < 0$. Then, $0 \leq \varepsilon^2 U_{rr}(r_1) = -f(U(r_1))$, hence $U(r_1) \leq -1$. We obtain a contradiction, because $\|U - w\|_\infty < C\varepsilon$. Therefore $U \geq 0$. The equality does not hold, because of the strong maximum principle. Thus, $U > 0$. \square

7.6. Nondegeneracy.

Lemma 7.12. *Let $\{(\lambda, U)\}$ be a family of solutions obtained in Corollary 7.11, and let $L := \varepsilon^2 (\partial_{rr} + \frac{N-1}{r} \partial_r) + f'(U)$. Then there is small*

$\delta > 0$ such that L has no eigenvalue in $[-\delta, \delta]$ if $\varepsilon > 0$ is small. In particular, U is nondegenerate in \mathbb{X} .

Proof. Let $\delta > 0$ be a constant used in the proof of Lemma 7.7. Then, \mathcal{L} has no eigenvalue in $[-\delta, \delta]$. We show that L has no eigenvalue in $[-\delta/2, \delta/2]$. Suppose the contrary, i.e., there is $\kappa_\lambda \in [-\delta/2, \delta/2]$ such that $L\phi = \kappa_\lambda\phi$ ($\|\phi\|_{\mathbb{X}} = 1$). Then,

$$(7.10) \quad (\mathcal{L} - \kappa_\lambda)\phi + (f'(U) - f'(w))\phi = 0.$$

Because of Lemma 7.7, $(\mathcal{L} - \kappa_\lambda)$ is invertible and there is $C > 0$ independent of small $\varepsilon > 0$ such that $\|(\mathcal{L} - \kappa_\lambda)^{-1}\|_{\mathcal{B}(\mathbb{X}, \mathbb{Y})} \leq C$. It follows from Corollary 7.11 that $\|f'(U) - f'(w)\|_\infty \rightarrow 0$ as $\varepsilon \downarrow 0$. By (7.10) we have

$$\begin{aligned} \|\phi\|_{\mathbb{X}} &= \|(\mathcal{L} - \kappa_\lambda)^{-1} [(f'(U) - f'(w))\phi]\|_{\mathbb{X}} \\ &\leq \|(\mathcal{L} - \kappa_\lambda)^{-1}\|_{\mathcal{B}(\mathbb{X}, \mathbb{Y})} \| (f'(U) - f'(w))\phi \|_{\mathbb{Y}} \\ &\leq C \|f'(U) - f'(w)\|_\infty \|\phi\|_{\mathbb{Y}} \rightarrow 0 \quad (\varepsilon \downarrow 0), \end{aligned}$$

which contradicts that $\|\phi\|_{\mathbb{X}} = 1$. \square

7.7. Asymptotic behavior of $\lambda_1(\gamma)$ ($\gamma \downarrow 0$).

Theorem 7.13. *Let $(\lambda_1(\gamma), U_1(r, \gamma)) \in \mathcal{C}_1$. Then, $U_1(r, \gamma)$ is a boundary concentrating solution obtained in Corollary 7.11 if $\gamma > 0$ is small.*

Proof. Since $|w(r) - w(r)^p| \leq C$ ($0 \leq r \leq 1$) and $w(r) \xrightarrow{\varepsilon \downarrow 0} 0$ pointwisely in $[0, 1)$, by the dominated convergence theorem we see that if $r > 1/2$, then

$$\begin{aligned} \varepsilon^2 \frac{|w_r(r)|}{r} &\leq \frac{1}{r} \int_0^r (|w(s)| + |w(s)|^p) ds \\ &\leq 2 \int_0^1 (|w(s)| + |w(s)|^p) ds \rightarrow 0 \quad (\varepsilon \downarrow 0). \end{aligned}$$

Hence, $\|\mathcal{M}w\|_\infty \rightarrow 0$ ($\varepsilon \downarrow 0$). Since v satisfies (7.7), by the elliptic regularity we see that $\|v\|_{C^1} \rightarrow 0$ ($\varepsilon \downarrow 0$). If $v(x_0) + w(x_0) = 1$, then $x_0 > 1/2$ and by phase plane argument we see that $w'(x_0) (> 0)$ is large. Since $v'(x_0)$ is small, $v'(x_0) + w'(x_0) > 0$. Therefore, $\mathcal{Z}_{[0,1]}[v(\cdot) + w(\cdot) - 1] = 1$ provided that $\varepsilon > 0$ is small. The boundary concentrating solution obtained by Corollary 7.11 belongs to \mathcal{C}_1 , because of (1.10). The proof of (1.10) is postponed until the proof of Theorem B in Section 8. \square

Corollary 7.14. *Let $(\lambda_1(\gamma), U_1(r, \gamma)) \in \mathcal{C}_1$. Then, $\lambda_1(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$). Moreover, $\partial_\gamma \lambda_1(\gamma) < 0$ for small $\gamma > 0$.*

Proof. Because of Lemma 7.12, if $\varepsilon > 0$ is small, then U is nondegenerate, which indicates the following: If $\lambda_1(\gamma) > 0$ is large, then

$$(7.11) \quad \partial_\gamma \lambda_1(\gamma) \neq 0.$$

Because of Corollary 7.11, $\|U - w\|_\infty < C\varepsilon$. Since $\varepsilon = (\lambda_1(\gamma))^{-\frac{1}{2}}$, we have

$$\begin{aligned} (\gamma(\lambda_1) =) U(0) &\leq |U(0) - w(0)| + |w(0)| \\ &< O\left(\frac{1}{\sqrt{\lambda_1}}\right) + o(1) \quad (\lambda_1 \rightarrow \infty). \end{aligned}$$

Therefore,

$$(7.12) \quad \gamma(\lambda_1) \rightarrow 0 \quad (\lambda_1 \rightarrow \infty).$$

By (7.11) and (7.12) we see that

$$(7.13) \quad \partial_\gamma \lambda_1(\gamma) < 0 \quad \text{if } \lambda_1(\gamma) > 0 \text{ is large.}$$

By (7.12) and (7.13) we see that $\lambda_1(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$). \square

8. PROOF OF THEOREM B

Proof of Theorem B. (i) is mentioned in Section 1. (ii) follows from Theorem 5.6. (iii) is proved in Theorem 6.1. (v) follows from Theorem 7.13 and Corollary 7.14. Let $u(s, \gamma)$ be the solution of (1.7). Then, we already saw in Lemmas 3.3 and 3.5 that if $\gamma \neq 1$, then $u(s, \gamma)$ has infinitely many critical points. Let $\{s_n\}_{n=1}^\infty$ ($0 < s_1 < s_2 < \dots$) denote the set of the critical points of $u(s, \gamma)$. Since $\lambda_n(\gamma) = s_n^2$, (vi) holds. Corollary 7.14 says that $\lambda_1(\gamma) \rightarrow \infty$ ($\gamma \downarrow 0$). Since $\lambda_1(\gamma) < \lambda_2(\gamma) < \dots$, (iv) holds.

We prove (1.10). Let $(\lambda_0, U_0) \in \mathcal{S}$, and let $m := \mathcal{Z}_{[0,1]}[U(\cdot) - 1]$. Proposition 3.1 says that there is a one-parameter family of solutions $\mathcal{C} := (\lambda(\gamma), U(\gamma))$ such that $(\lambda(0), U(0)) = (\lambda_0, U_0)$. This branch \mathcal{C} can be extended to $\gamma = 1$, because of Lemmas 3.3 and 3.5. Note that $\mathcal{Z}_{[0,1]}[U(\cdot, \gamma) - 1] = m$ if $(\lambda, U) \in \mathcal{C}$ and $U \neq 1$. It follows from the uniqueness of the branch near $(\bar{\lambda}_m, 1)$ that $\mathcal{C} \subset \mathcal{C}_m$. Thus, $(\lambda_0, U_0) \in \mathcal{C} \subset \mathcal{C}_m$, and the (1.10) holds. The proof is complete. \square

Proof of Corollary 1.3. Let $\underline{\lambda} := \inf_{\gamma \in \mathbb{R}_+} \lambda_1(\gamma) > 0$. Because of (ii) and (iv), $\underline{\lambda} > 0$. For each $n \geq 2$, $\lambda_n(\gamma) > \lambda_1(\gamma) (\geq \underline{\lambda})$. This implies that the first assertion holds. The other assertion follows from the boundedness of $\{\lambda_1(\gamma)\}_{\gamma > 1}$. \square

Proof of Corollary 1.4. The statement clearly follows from (ii), (iii), and (iv) of Theorem B. \square

REFERENCES

- [1] Adimurthi and S. Yadava, *Existence and nonexistence of positive radial solutions of Neumann problems with critical Sobolev exponents*, Arch. Rational Mech. Anal. **115** (1991), 275–296.
- [2] Adimurthi and S. Yadava, *Nonexistence of positive radial solutions of a quasilinear Neumann problem with a critical Sobolev exponent*, Arch. Rational Mech. Anal. **139** (1997), 239–253.
- [3] C. Budd, M. Knaap, and A. Peletier, *Asymptotic behavior of solutions of elliptic equations with critical exponents and Neumann boundary conditions*, Proc. Roy. Soc. Edinburgh Sect. A **117** (1991), 225–250.
- [4] C. Budd and J. Norbury, *Semilinear elliptic equations and supercritical growth*, J. Differential Equations **68** (1987), 169–197.
- [5] J. Chern, Z. Chen, J. Chen, and Y. Tang, *On the classification of standing wave solutions for the Schrödinger equation*. Comm. Partial Differential Equations **35** (2010), 275–301.
- [6] M. Crandall and P. Rabinowitz, *Bifurcation from simple eigenvalues*, J. Functional Analysis **8** (1971), 321–340.
- [7] J. Dolbeault and I. Flores, *Geometry of phase space and solutions of semilinear elliptic equations in a ball*, Trans. Amer. Math. Soc. **359** (2007), 4073–4087.
- [8] P. de Groen and G. Karadzhov, *Metastability in the shadow system for Gierer-Meinhardt’s equations*, Electron. J. Differential Equations (2002), No. 50, 22 pp. (electronic).
- [9] M. del Pino, M. Musso, and A. Pistoia, *Super-critical boundary bubbling in a semilinear Neumann problem*, Ann. Inst. H. Poincaré Anal. Non Linéaire **22** (2005), 45–82.
- [10] P. Fife, *Semilinear elliptic boundary value problems with small parameters*, Arch. Rational Mech. Anal. **52** (1973), 205–232.
- [11] M. Grossi and B. Noris, *Positive constrained minimizers for supercritical problems in the ball*, Proc. Amer. Math. Soc. **140** (2012), 2141–2154.
- [12] A. Gierer and H. Meinhardt, *A theory of biological pattern formation*, Kybernetik (Berlin) **12** (1972), 30–39.
- [13] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*. Second edition, *Grundlehren der Mathematischen Wissenschaften, 224*. Springer-Verlag, Berlin, 1983.
- [14] C. Gui and J. Wei, *On multiple mixed interior and boundary peak solutions for some singularly perturbed Neumann problems*, Canad. J. Math. **52** (2000), 522–538.
- [15] Z. Guo and J. Wei, *Global solution branch and Morse index estimates of a semilinear elliptic equation with super-critical exponent*, Trans. Amer. Math. Soc. **363** (2011), 4777–4799.
- [16] D. Henry, *Geometric theory of semilinear parabolic equations*, *Lecture Notes in Mathematics, 840*. Springer-Verlag, Berlin-New York, 1981.

- [17] D. Joseph and T. Lundgren, *Quasilinear Dirichlet problems driven by positive sources*, Arch. Rational Mech. Anal. **49** (1973), 241–269.
- [18] M. Kwong, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbf{R}^n* , Arch. Rational Mech. Anal. **105** (1989), 243–266.
- [19] E. Keller and L. Segal, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol. **26** (1970), 399–415.
- [20] C.-S. Lin and W.-M. Ni, *On the diffusion coefficient of a semilinear Neumann problem*, Calculus of variations and partial differential equations (Trento, 1986), 160–174, Lecture Notes in Math., 1340, Springer, Berlin, 1988.
- [21] A. Malchiodi and M. Montenegro, *Boundary concentration phenomena for a singularly perturbed elliptic problem*, Comm. Pure Appl. Math. **55** (2002), 1507–1568.
- [22] F. Merle and L. Peletier, *Positive solutions of elliptic equations involving supercritical growth*, Proc. Roy. Soc. Edinburgh Sect. A **118** (1991), 49–62.
- [23] Y. Miyamoto, *Structure of the positive solutions for supercritical elliptic equations in a ball*, preprint.
- [24] Y. Miyamoto and K. Yagasaki, *Monotonicity of the first eigenvalue and the global bifurcation diagram for the branch of interior peak solutions*, J. Differential Equations **254** (2013), 342–367.
- [25] W.-M. Ni and J. Serrin, *Nonexistence theorems for singular solutions of quasilinear partial differential equations*, Comm. Pure Appl. Math. **39** (1986), 379–399.
- [26] W.-M. Ni, *On the positive radial solutions of some semilinear elliptic equations on R^n* , Appl. Math. Optim. **9** (1983), 373–380.
- [27] W.-M. Ni and I. Takagi, *On the shape of least-energy solutions to a semilinear Neumann problem*, Comm. Pure Appl. Math. **44** (1991), 819–851.
- [28] S. Pohožaev, *On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$* , Dokl. Akad. Nauk SSSR **165** (1965) 36–39.
- [29] J. Wei, *On the boundary spike layer solutions to a singularly perturbed Neumann problem*, J. Differential Equations **134** (1997), 104–133.
- [30] J. Wei, *On the interior spike layer solutions to a singularly perturbed Neumann problem*, Tohoku Math. J. **50** (1998), 159–178.
- [31] J. Wei, *On single interior spike solutions of the Gierer-Meinhardt system: uniqueness and spectrum estimates*, European J. Appl. Math. **10** (1999), 353–378.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,
3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN
E-mail address: miyamoto@ms.u-tokyo.ac.jp

UTMS

- 2012–13 H. Kawakami, A. Nakamura, and H. Sakai: *Degeneration scheme of 4-dimensional Painlevé-type equations.*
- 2012–14 Yusaku Tiba: *The second main theorem for entire curves into Hilbert modular surfaces.*
- 2012–15 O. Yu. Imanuvilov and M. Yamamoto: *Inverse boundary value problem for Schrödinger equation in cylindrical domain by partial boundary data.*
- 2012–16 Nao Hamamuki: *A discrete isoperimetric inequality on lattices.*
- 2012–17 Nao Hamamuki: *Asymptotically self-similar solutions to curvature flow equations with prescribed contact angle and their applications to groove profiles due to evaporation-condensation.*
- 2012–18 Guanyu Zhou: *Analysis of the fictitious domain method with L^2 -penalty for elliptic and parabolic problems.*
- 2013–1 Yasuko Hasegawa: *The critical values of exterior square L -functions on $GL(2)$.*
- 2013–2 Toshio Oshima: *An elementary approach to the Gauss hypergeometric function.*
- 2013–3 Kazuki Okamura: *Large deviations for simple random walk on percolations with long-range correlations .*
- 2013–4 Kiyoomi Kataoka and Nobuko Takeuchi: *A system of fifth-order PDE's describing surfaces containing 2 families of circular arcs and the reduction to a system of fifth-order ODE's .*
- 2013–5 Takushi Amemiya: *Orders of meromorphic mappings into Hopf and Inoue surfaces.*
- 2013–6 Yasuhito Miyamoto: *Structure of the positive radial solutions for the supercritical Neumann problem $\varepsilon^2 \Delta u - u + u^p = 0$ in a ball.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012