


UTMS 2017–3

October 03, 2017

**An application of the value distribution
theory for semi-abelian varieties to
problems of Ax-Lindemann and
Manin-Mumford types**

by

Junjiro NOGUCHI



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

An application of the value distribution theory for semi-abelian varieties to problems of Ax-Lindemann and Manin-Mumford types

Junjiro Noguchi*

The University of Tokyo

Abstract

The aim of this paper is to prove a theorem of Ax-Lindemann type for complex semi-abelian varieties as an application of a big Picard theorem proved by the author in 1981, and then apply it to prove a theorem of classical Manin-Mumford Conjecture for semi-abelian varieties, which was proved by M. Raynaud 1983, M. Hindry 1988, . . . , and Pila-Zannier 2008 by a different method from others, which is most relevant to ours. The present result might be a first instance of a *direct connection* between the value distribution theory of holomorphic maps and the arithmetic (Diophantine) theory over algebraic number fields, while there have been many *analogies* between them.

1 Introduction and main results

The purpose of this paper is to prove a theorem of Ax-Lindemann type for complex semi-abelian varieties as an application of a big Picard theorem obtained in [6], 1981. We then apply it to prove a theorem of Manin-Mumford type¹⁾ for the distribution of torsion points on a subvariety of a semi-abelian variety defined over a number field, combined with extending a part of the arguments in Pila-Zannier [11] for abelian varieties (cf. §3). In the course of the proof the Kawamata structure theorem for semi-abelian varieties by [6], Lemma (4.1) works quite effectively (see §3 (b)).

Theorem 1.1 (Ax-Lindemann type). *Let $\exp : \mathbf{C}^n \rightarrow A$ be an exponential map of a complex semi-abelian variety A . Let $V \subset \mathbf{C}^n$ be a complex irreducible affine algebraic subvariety with the restricted map $\exp|_V : V \rightarrow A$. Then the Zariski closure $X(\exp|_V)$ of the image of $\exp|_V$ in A is a translate of a complex semi-abelian subvariety of A .*

Theorem 1.2 (Manin-Mumford type). *Let $X \subset A$ be a proper algebraic reduced subvariety of a semi-abelian variety A defined over an algebraic number field. Then, the Zariski closure $\overline{X} \cap A_{\text{tor}}^{\text{Zar}}$ of torsion points on X is a finite union of translates of semi-abelian subvarieties by torsion points on X .*

In view of the above two theorems, it is naturally led to study a denseness property of the value-distribution of $\exp|_V$ in Theorem 1.1. In fact, for an ample divisor D on a projective compactification of A , we will prove that $D \cap \exp(V)$ is Zariski dense in $D \cap X(\exp|_V)$ (cf. Theorem 4.2 in §4).

*Research supported in part by Grant-in-Aid for Scientific Research (C) 15K04917.

MSC2010: 11J95, 32H30, 03C64

Key words: Ax-Lindemann, Manin-Mumford, big Picard, Nevanlinna theory, semi-abelian variety

¹⁾ It was called the Manin-Mumford Conjecture and proved by M. Raynaud [12], M. Hindry [3], . . . , Pila-Zannier [11]; cf. [11], Introduction, S. Lang [4], Chap. I §6, and e.g., P. Tzermias [13] for surveys of the Manin-Mumford Conjecture.

In §2 we will introduce a new notion of “*strictly transcendental*” for holomorphic maps into semi-abelian varieties (see Definition 2.3). By making use of a Big Picard Theorem due to [6] we prove the image structure stated in Theorem 1.1 for strictly transcendental holomorphic maps into semi-abelian varieties (see Theorem 2.4). We then prove that the map $\exp|_V$ in Theorem 1.1 is strictly transcendental (Proposition 2.5).

In §3 we will prove Theorem 1.2 by induction in $\dim X$, in which we will use [6], Lemma (4.1). The proof of Theorem 1.2 roughly consists of two parts: The first is a decomposition of the torsion points on X to an algebraic part and its compliment, done by the arguments due to Pila-Zannier [11], Theorem 2.1, Step 1, being extended to the semi-abelian case. The second part is the application of Theorem 1.1 to the algebraic part of torsion points, and for its compliment we use the lower and upper estimates due to Masser and Pila-Wilkie (cf. [11], §3) to deduce the statement of Theorem 1.2.

In the last §5 we give some example of a transcendental but not strictly transcendental map into an abelian variety, and study its image structure.

The present result might be a first instance of a *direct connection* between the value distribution theory of holomorphic maps and the arithmetic (Diophantine) theory over algebraic number fields, although there have been many *analogies* between them. It is the point of interest of this paper to observe that the direct connection is provided by the theory of o-minimal structures in model theory (Pila-Wilkie [9]; this approach evolved from Bombieri-Pila [1] (cf. Zannier [14])). This aspect should be of some interest for both sides of the theories.

Acknowledgment. The present paper has grown up through the interesting and helpful discussions with Professors Pietro Corvaja and Umberto Zannier since Conference “Specialization Problems in Diophantine Geometry”, Cetraro (Italy), 9–14 July 2017. In particular, the proof of Theorem 1.2 in §3 is based on their arguments suggested through the discussions. It is a great pleasure of the author to express his deep gratitude to both of them.

2 Big Picard and Ax-Lindemann

(i) Big Picard. Let Δ^m be the unit polydisk of \mathbf{C}^m with center at 0 and let $E \subset \Delta^m$ be a thin complex analytic subset. Let Y be a reduced complex space with a compactification \bar{Y} , a compact complex space with reduced structure. Let $f : \Delta^m \setminus E \rightarrow Y$ be a holomorphic map. If there is a meromorphic map $\bar{f} : \Delta^m \rightarrow \bar{Y}$ with the restriction to $\Delta^m \setminus E$, $\bar{f}|_{(\Delta^m \setminus E)} = f$, we say that E is *removable* for f or f is (meromorphically) extendable over E : Otherwise, f is said to be *transcendental*.

We denote by $X(f)$ the Zariski closure of the image of f in Y with respect to the Zariski topology on \bar{Y} . Note that $X(f)$ is irreducible.

Remark 2.1 (Hartogs extension theorem). Let Y be quasi-projective algebraic. If $\text{codim } E \geq 2$, E is always removable for any holomorphic map $f : \Delta^m \setminus E \rightarrow Y$. Therefore, as the removability is concerned with quasi-projective Y , it suffices to deal with smooth E of codimension 1.

Let A be a complex semi-abelian variety with a smooth projective compactification \bar{A} , and let $X \subset A$ be a non-empty complex algebraic subset. We denote by $\text{St}^0(X)$ the connected component of 0 of the group $\{a \in A : a + X = X\}$, and simply call it the *stabilizer* of X . Then, $\text{St}^0(X)$ is an algebraic subgroup of A and a semi-abelian subvariety by itself. Note that X is of general type if and only if $\text{St}^0(X) = \{0\}$.

We recall:

Theorem 2.2 ([6], Corollary (4.7)). *Assume that X is an irreducible algebraic subvariety of A . Let $f : \Delta^m \setminus E \rightarrow X \hookrightarrow A$ be a holomorphic map. If $\text{St}^0(X(f)) = \{0\}$, then f is extendable over E ; in the other words, if f is transcendental, then $\dim \text{St}^0(X(f)) > 0$.*

Definition 2.3 (strictly transcendental). A transcendental holomorphic map $f : \Delta^m \setminus E \rightarrow A$ is said to be *strictly transcendental* if for every semi-abelian subvariety $B \subset A$ the composite $q_B \circ f : \Delta^m \setminus E \rightarrow A/B$ with the quotient map $q_B : A \rightarrow A/B$ is either constant or transcendental.

Theorem 2.4. *Let $f : \Delta^m \setminus E \rightarrow A$ be a strictly transcendental holomorphic map. Then, $X(f) = f(c) + \text{St}^0(X(f))$ with $c \in \Delta^m \setminus E$; i.e., $X(f)$ is a translate of a semi-abelian subvariety of A .*

Proof. Set $B = \text{St}^0(X(f))$ and $g = q_B \circ f : \Delta^m \setminus E \rightarrow A/B$. It suffices to show that g is constant. Otherwise, g would be transcendental by the assumption for f . Since $\text{St}^0(X(g)) = \{0\}$, it follows from Theorem 2.2 that g is extendable over E ; this is a contradiction. \square

(ii) Ax-Lindemann. We keep the notation used above. Here we always assumes that “varieties” are irreducible with reduced structure. Let

$$\exp : \mathbf{C}^n \rightarrow A$$

be an exponential map. A map from a complex algebraic variety into another complex algebraic variety is called a *transcendental* map if it is *not* a rational map. We then define a *strictly transcendental map* from a complex algebraic variety into A as in Definition 2.3.

Proposition 2.5. *Let $V \subset \mathbf{C}^n$ be a positive dimensional irreducible complex algebraic subvariety of \mathbf{C}^n . Then, the restricted map $\exp|_V : V \rightarrow A$ of \exp to V is strictly transcendental.*

Proof. Let $B \subsetneq A$ be a complex semi-abelian subvariety. Suppose that

$$q_B \circ \exp|_V : V \rightarrow A_1 = A/B$$

is a non-constant rational map. We shall deduce a contradiction.

There is an algebraic curve $C \subset V$ (irreducible and $\dim C = 1$) such that the restriction

$$f := q_B \circ \exp|_C : C \rightarrow A_1$$

is non-constant rational.

(a) Case of $B = \{0\}$: There is an exact sequence

$$(2.6) \quad 0 \rightarrow (\mathbf{C}^*)^t \rightarrow A \rightarrow A_0 \rightarrow 0,$$

where A_0 is an abelian variety. Set $L = \exp^{-1}(\mathbf{C}^*)^t$. Then L is a t -dimensional vector subspace of \mathbf{C}^n . Let $p_0 : \mathbf{C}^n \rightarrow \mathbf{C}^n/L \cong \mathbf{C}^m$ ($m = n - t$) and $q_0 : A \rightarrow A_0$ be the quotient maps. Then \exp naturally induces an exponential map

$$\exp_0 : \mathbf{C}^m \rightarrow A_0.$$

We are going to infer

Claim 2.7. $q_0 \circ \exp(C)$ is a point.

Suppose that it is not the case. Then, the Zariski closure C_0 of the image $p_0(C)$ in \mathbf{C}^m is an algebraic curve in \mathbf{C}^m . Let ω_0 be a flat Kähler metric on A_0 such that $\exp_0^* \omega_0 = \alpha$, where $\alpha = \sum_{j=1}^m \frac{i}{2\pi} dz_j \wedge d\bar{z}_j$ with the natural coordinate system (z_1, \dots, z_m) of \mathbf{C}^m . Since $\exp|_C$ is rational, $\exp_0|_{C_0}$ is non-constant rational. Then, the Zariski closure $W := \overline{\exp_0(C_0)}^{\text{Zar}}$ in A_0 is an algebraic curve in A_0 with

$$(2.8) \quad \int_{\exp_0(C_0)} \omega_0 = \int_W \omega_0 = M < \infty.$$

If k denotes the degree of the rational map $\exp_0|_{C_0} : C_0 \rightarrow W$, we have

$$(2.9) \quad \int_{C_0} \alpha = kM.$$

Let $B(r) \subset \mathbf{C}^m$ be an open ball of radius $r(> 0)$ with center at a point $a \in C_0$ and set $C_0(r) = C_0 \cap B(r)$. Wirtinger's inequality implies

$$\int_{C_0(r)} \alpha \geq \nu(a; C_0) r^2,$$

where $\nu(a; C_0) (\geq 1)$ denotes the order of C_0 at a . Letting $r \rightarrow \infty$, we have a contradiction to (2.9). Therefore, Claim 2.7 follows.

Thus, we have a non-constant rational map after a translation:

$$\exp|_C : C(\subset L \cong \mathbf{C}^t) \rightarrow (\mathbf{C}^*)^t,$$

which is a restriction of exponential map

$$(\zeta_1, \dots, \zeta_t) \in \mathbf{C}^t \rightarrow (e^{\zeta_1}, \dots, e^{\zeta_t}) \in (\mathbf{C}^*)^t.$$

Let \bar{C} be the closure of C in $(\mathbf{P}^1(\mathbf{C}))^t \supset (\mathbf{C}^*)^t$. Here we write $\mathbf{P}^1(\mathbf{C}) = \mathbf{C} \cup \{\infty\}$. Then there is a point $b = (b_1, \dots, b_t) \in \bar{C} \setminus C$ with some $b_j = \infty$. Since e^{ζ_j} has an isolated *essential* singularity at $\zeta_j = \infty$, $\exp|_C$ cannot be rational.

(b) Case of general B : Set

$$\begin{aligned} F &= \exp^{-1} B, \\ p' : \mathbf{C}^n &\rightarrow \mathbf{C}^n / F \cong \mathbf{C}^{n'}, \\ q' : A &\rightarrow A/B = A'. \end{aligned}$$

Let $\exp' : \mathbf{C}^{n'} \rightarrow A'$ be the naturally induced exponential from $\exp : \mathbf{C}^n \rightarrow A$. Let C' be the Zariski closure of $p'(C)$ in $\mathbf{C}^{n'}$. Then, it follows from the assumption that $\pi'|_{C'} : C' \rightarrow A'$ would be a non-constant rational map: This contradicts to what was proved in (a) above. \square

Proof of Theorem 1.1: This is now immediate by Proposition 2.5 and Theorem 2.4. \square

Example 2.10. Here we give a simple example of a strictly transcendental map. Set

$$\begin{aligned} \exp : (z, w) \in \mathbf{C}^2 &\rightarrow (e^z, e^w) \in (\mathbf{C}^*)^2 = A, \\ V &= \{(z, w) \in \mathbf{C}^2 : zw = 1\} = \{(z, 1/z) : z \in \mathbf{C}^*\} \cong \mathbf{C}^*. \end{aligned}$$

Then V is affine algebraic,

$$\exp_V : (z, 1/z) \in V \rightarrow (e^z, e^{1/z}) \in A$$

is strictly transcendental, and $X(\exp|_V) = A$.

3 Proof of Theorem 1.2

Let K be a number field over which A is defined. Let

$$\exp : \mathbf{C}^n \rightarrow A$$

be an exponential map. The reduction of $\exp^{-1} X \cap A_{\text{tor}}$ to the algebraic part is done in parallel to the proof of Pila-Zannier [11], §3, and then we apply Theorem 1.1 to conclude the theorem.

The proof is done by induction on $\nu = \dim X \geq 0$. If $\nu = 0$, it is trivial. Suppose that the case of $\dim X \leq \nu - 1 (\nu \geq 1)$ holds. Let $\dim X = \nu$.

We consider the stabilizer $\text{St}^0(X)$.

(a) The case of $\dim \text{St}^0(X) > 0$. We set the quotient map $q : A \rightarrow A_1 = A/\text{St}^0(X)$ and $X_1 = q(X) = X/\text{St}^0(X)$. Since $\dim X_1 < \nu$, the induction hypothesis implies that there are at most finitely many semi-abelian subvarieties $B_j \subset A_1, 1 \leq j \leq l$, and torsion points $b_j \in X_1 \cap A_{1\text{tor}}$ such that

$$X_1 \cap A_{1\text{tor}} = \bigcup_{j=1}^l (b_j + B_{j\text{tor}}).$$

Taking any elements $a_j \in q^{-1}b_j \cap X \cap A_{\text{tor}}, 1 \leq j \leq l$, we have

$$X \cap A_{\text{tor}} \subset \bigcup_{j=1}^l (a_j + (q^{-1}B_j)_{\text{tor}}).$$

Since $X \supset a_j + q^{-1}B_j$, the opposite inclusion above also holds; thus,

$$\overline{X \cap A_{\text{tor}}}^{\text{Zar}} = \bigcup_{j=1}^l (a_j + q^{-1}B_j).$$

(b) The case of $\text{St}^0(X) = \{0\}$. Note that X is of (log) general type. We put $\Lambda = \exp^{-1}\{0\} (\subset \mathbf{C}^n)$ and $Z = \exp^{-1}X$ which is a Λ -periodic analytic subset of \mathbf{C}^n . We consider a basis of Λ as \mathbf{Z} -module, which is also a basis of the real vector space $\mathbf{R}\Lambda$ of dimension $d = 2n - t$. Thus we have identifications

$$\mathbf{Z}^d \cong \mathbf{Z}\Lambda \subset \mathbf{R}\Lambda \cong \mathbf{R}^d.$$

The torsion points on X lift to points on $\mathbf{Q}\Lambda \cong \mathbf{Q}^d \subset \mathbf{R}^d$. We consider the set $\check{Z} = Z \cap \mathbf{Q}^d$ of the preimage of all torsion points on X and its restriction $\check{Z}_1 := \check{Z} \cap [0, 1]^d$ of \check{Z} to the closed fundamental domain $[0, 1]^d$.

For a subset $W \subset \mathbf{Q}^d$ and a real number $T \geq 1$, we denote by $N_W(T)$ the number of rational points in W whose denominator is at most T .

By Pila-Wilkie [9] there is a so-called algebraic part \check{Z}_1^{alg} of \check{Z}_1 , satisfying the following properties:

(i) We have

$$(3.1) \quad \check{Z}_1^{\text{alg}} = \bigcup_{V \subset Z} (V \cap \{t = (t_j) \in [0, 1]^d : t_j \in \mathbf{Q}\}),$$

where V runs over all positive dimensional affine algebraic subsets of \mathbf{C}^n , contained in Z (cf. [11], Proposition 2.1²⁾).

(ii) For every $\varepsilon > 0$ there is a positive constant $c_1 (= c_1(\check{Z}_1, \varepsilon))$, depending on \check{Z}_1 and ε such that

$$(3.2) \quad N_{\check{Z}'_1}(T) \leq c_1 T^\varepsilon, \quad T \geq 1,$$

where $\check{Z}'_1 := \check{Z}_1 \setminus \check{Z}_1^{\text{alg}}$, called the transcendental part of \check{Z}_1 ([9]).

We first analyze the algebraic part \check{Z}_1^{alg} . If $\check{Z}_1^{\text{alg}} = \emptyset$, it is done; we assume $\check{Z}_1^{\text{alg}} \neq \emptyset$. Let V be as in (3.1). It follows from Theorem 1.1 that $X(\exp|_V) (\subset X)$ is a translate of a positive dimensional semi-abelian subvariety of A . Let Y denote the union of all translates $a + B'$ ($a \in X$) of positive dimensional semi-abelian subvarieties $B' \subset A$ such that $a + B' \subset X$. It follows from [6], Lemma (4.1) (cf. also [7], §5.6.4, and [11], p.160) that $Y (\subset X)$ is an algebraic subset of dimension $< \nu$ and for every irreducible component Y' of Y , $\dim \text{St}^0(Y') > 0$. We may assume that Y is defined over K . Applying the induction

²⁾ It is noted that the periodicity condition of Proposition 2.1 is not used in the proof, so that it can be applied to our Z .

hypothesis to Y , we have finitely many positive dimensional semi-abelian subvarieties $B'_j \subset A, 1 \leq j \leq l'$, and $P_j \in Y \cap A_{\text{tor}}$ such that

$$Y \cap A_{\text{tor}} = \bigcup_{j=1}^{l'} (P_j + B'_{j\text{tor}}).$$

Therefore, we have

$$\exp(Z_1^{\text{alg}}) = \bigcup_{j=1}^{l'} (P_j + B'_{j\text{tor}}), \quad \overline{\exp(\check{Z}'_1^{\text{alg}})}^{\text{Zar}} = \bigcup_{j=1}^{l'} (P_j + B'_j).$$

We may assume that all P_j and B'_j are defined over K after a finite extension of K . Then, we have:

3.3 (Invariance). For a torsion point $P \in \exp(\check{Z}'_1^{\text{alg}})$ (resp. $\exp(\check{Z}'_1)$), all of its conjugates over K lie in $\exp(\check{Z}'_1^{\text{alg}})$ (resp. $\exp(\check{Z}'_1)$).

To analyze the part \check{Z}'_1 we prepare:

Lemma 3.4. *Let $P \in A$ be a torsion point of exact order N . Then, there is a number $\rho > 0$ depending only on $\dim A$ such that*

$$(3.5) \quad [K(P) : K] \geq c_2 N^\rho, \quad T \geq 1,$$

where $c_2 = c_2(A, K)$ is a positive constant with dependence as signed.

Proof. For the semi-abelian variety A we have the following exact sequence

$$0 \rightarrow \mathbf{G}_m^t \rightarrow A \xrightarrow{\pi} A_0 \rightarrow 0,$$

where \mathbf{G}_m^t is an algebraic torus and A_0 is an abelian variety; we may assume that all of the above morphisms and algebraic groups are defined over K . Then, $P_0 := \pi(P)$ is a torsion point of A_0 , whose exact order is denoted by N_0 . Note that $N_0 P \in \mathbf{G}_m^t$ and $N = N_1 N_0$, where N_1 is the order of $N_0 P \in \mathbf{G}_m^t$. Then,

$$c_3 \varphi(N_1) \leq [K(N_0 P) : K] \leq [K(P) : K],$$

where $c_3 = c_3(t, K)$ is a positive constant and φ is the Euler function. It is known that

$$\varphi(N_1) \sim \frac{N_1}{\log \log N_1}.$$

If $N_1 \geq \sqrt{N}$, the proof is finished. Otherwise, we have $N_0 \geq \sqrt{N}$. We then apply Masser's estimate ([5]) for the following second inequality:

$$c_3 N^{\rho'/2} \leq c_3 N_0^{\rho'} \leq [K(P_0) : K] \leq [K(P) : K],$$

where $c_3 = c_3(A_0, K)$ is a positive constant. Thus, we have the lower estimate (3.5). \square

Finish of the proof of Theorem 1.2: Let now $P \in \exp(\check{Z}'_1)$ be a torsion point of exact order N . It follows from Invariance 3.3 that $N_{\check{Z}'_1}(N) \geq [K(P) : K]$. By Lemma 3.4 we see that

$$(3.6) \quad N_{\check{Z}'_1}(N) \geq c_2 N^\rho.$$

On the other hand, we have by (3.2)

$$(3.7) \quad N_{\check{Z}'_1}(N) \leq c_1 N^\varepsilon.$$

Taking $\varepsilon = \rho/2$ with ρ in (3.6), we conclude the boundedness of N by (3.6) and (3.7). Therefore, \check{Z}'_1 is a finite set. \square

4 Distribution of $\exp(V)$ on divisors

Let A be a semi-abelian variety with exponential map

$$\exp : \mathbf{C}^n \longrightarrow A,$$

and let $V \subset \mathbf{C}^g$ be an irreducible affine algebraic subvariety. Let \bar{A} be a projective compactification of A . Because of the results of the previous sections, it might be of some interest to look at the actual value-distribution of $\exp|_V(V)$ in its Zariski closure $X(\exp|_V)$.

We first deal with a transcendental holomorphic map $f : \Delta^* \rightarrow A$ from a punctured disk Δ^* into A . In [8] we dealt with entire holomorphic maps from the whole plane \mathbf{C} into A . Combining the method of [6] with the result and the arguments explored in [8] and [7] we see that the results of [8] (and [7], Chap. 6) hold for transcendental holomorphic maps from Δ^* into A . In particular, we have (cf., also [2], Theorem 5.2)

Theorem 4.1. *Let $f : \Delta^* \rightarrow A$ be a transcendental holomorphic map and let D be an effective ample divisor on \bar{A} such that $D \not\supset f(\Delta^*)$. Then $f(\Delta^*) \cap X(f) \cap D$ is Zariski dense in $X(f) \cap D$ (where D stands also for the support of D).*

By making use of this we prove:

Theorem 4.2. *Let $V \subset \mathbf{C}^n$ be an irreducible complex affine algebraic subvariety, and let D be an effective ample divisor on \bar{A} . Then, the intersection $D \cap \exp(V)$ is Zariski dense in $D \cap X(\exp|_V)$.*

Proof. Let $\zeta_0 \in V$ be fixed. We consider a pencil of affine algebraic curves $C_\gamma \subset V, \gamma \in \Gamma$, passing through ζ_0 , such that $\bigcup_\gamma C_\gamma$ contains a non-empty open subset of V in the sense of differential topology. By Theorem 1.1 $X(\exp|_V)$ and $X(\exp|_{C_\gamma})$ are all translates of semi-abelian subvarieties of A passing through $\exp(\zeta_0)$. Since there are at most countably many such semi-abelian subvarieties, one finds a curve $C_0 = C_\gamma$ such that $X(\exp|_{C_0}) = X(\exp|_V)$. Then it suffices to show the theorem for C_0 . Let $C_1 \rightarrow C_0$ be the normalization and let \bar{C}_1 be its smooth compactification. Then there is an analytic neighborhood $U (\subset \bar{C}_1)$ of a point Q of $\bar{C}_1 \setminus C_1$ such that $U \setminus \{Q\}$ is biholomorphic to a punctured disk Δ^* . Then our assertion is immediate by Theorem 4.1. \square

5 An example of a transcendental but not strictly transcendental map

Let \bar{C} be a smooth complex projective algebraic curve of genus $g \geq 1$ and let $q : \bar{C} \rightarrow J(\bar{C})$ be the Jacobian embedding; here, when $g = 1$, we simply take $q : \bar{C} \rightarrow \bar{C} (= J(\bar{C}))$ as the identity map. We set $A_1 = J(\bar{C})$, which is an abelian variety of dimension g . Let $Q \in \bar{C}$ be a point and set $C = \bar{C} \setminus \{Q\}$. Then C is affine algebraic and there is a finite map $p : C \rightarrow \mathbf{C}$.

Let $\exp : \mathbf{C}^g \rightarrow A_1$ be an exponential map. We take a linear embedding $\lambda : \mathbf{C} \rightarrow \mathbf{C}^g$ that is in sufficiently generic direction with respect to the period lattice of $\exp : \mathbf{C}^g \rightarrow A_1$. Then, $X(\exp \circ \lambda) = A_1$. We put

$$(5.1) \quad f : x \in C \longrightarrow (q(x), \exp \circ \lambda \circ p(x)) \in A_1 \times A_1 =: A.$$

Proposition 5.2. *Let $f : C \rightarrow A$ be as above.*

- (i) *The holomorphic map f is transcendental but not strictly transcendental.*
- (ii) *If $g \geq 2$, the Zariski closure $X(f)$ of the image $f(C)$ is not a translate of an abelian subvariety of A .*

(iii) If $g = 1$, $X(f) = A$.

Proof. (i) The first half is clear. For the latter, note that $X(f) \subset \bar{C} \times A_1$. With a subgroup $\{0\} \times A_1 \subset A_1 \times A_1 = A$ we consider the quotient map $\mu : A \rightarrow A/\{0\} \times A_1 \cong A_1$. Then, $\mu \circ f = q|_C : C \rightarrow A_1$ is rational. Therefore, f is not strictly transcendental.

(ii) Since $\mu(X(f)) = q(\bar{C})$, $X(f)$ is not a translate of an abelian subvariety.

(iii) Since $\dim X(f) = 1$, or 2, it suffices to deduce a contradiction with assuming $\dim X(f) = 1$. If so, $X(f)$ is a translate of an abelian subvariety of A . We consider an effective divisor $D = \bar{C} \times \{w\}$ with $w \in \bar{C}$. We infer from the definition of f that $X(f) \cap D$ is infinite. Therefore, $X(f) = D$; this is a contradiction. \square

Remark 5.3. ³⁾ Păun-Sibony [10] deals with a similar application of the Bloch-Ochiai Theorem to the abelian Ax-Lindemann statement ([10], Theorem 5.2). But with regard to Proposition 5.2 above, in Theorems 5.2 of [10] one might be able to have only the non-triviality of the stabilizer of the Zariski closure of the image as in Theorem 2.2, obtained in [6] (Corollary (4.7)).

References

- [1] E. Bombieri and J. Pila, The number of integral points on arcs and ovals, *Duke Math. J.* **59** (1989), 337–357.
- [2] P. Corvaja and J. Noguchi, A new unicity theorem and Erdős’ problem for polarized semi-abelian varieties, *Math. Ann.* **353** (2012), 439–464.
- [3] M. Hindry, Autour d’une conjecture de Serge Lang, *Invent. Math.* **94** (1988), 575–603.
- [4] S. Lang, Number Theory III, Diophantine Geometry, *Encycl. Math. Sci.* vol. **60**, Springer-Verlag, 1991.
- [5] D. Masser, Small values of the quadratic part of the Néron-Tate height on an abelian variety, *Compos. Math.* **53** (1984), 153–170.
- [6] J. Noguchi, Lemma on logarithmic derivatives and holomorphic curves in algebraic varieties, *Nagoya Math. J.* **83** (1981), 213–233.
- [7] J. Noguchi and J. Winkelmann, Nevanlinna Theory in Several Complex Variables and Diophantine Approximation, *Grundle. der Math. Wiss.* Vol. 350, Springer, Tokyo-Heidelberg-New York-Dordrecht-London, 2014.
- [8] J. Noguchi, J. Winkelmann and K. Yamanoi, The second main theorem for holomorphic curves into semi-abelian varieties II, *Forum Math.* **20** (2008), 469–503.
- [9] J. Pila and A.J. Wilkie, The rational points of a definable set, *Duke Math. J.* **133** (2006), 591–616.
- [10] M. Păun and N. Sibony, Value distribution theory for parabolic Riemann surfaces, preprint v4, arXiv, 8 April 2017.
- [11] J. Pila and U. Zannier, Rational points in periodic analytic sets and the Manin-Mumford conjecture, *Rend. Lincei Math. Appl.* **19** (2008), 149–162.
- [12] M. Raynaud, Sous-variété d’une variété abélienne et points de torsion, In *Arithmetic and Geometry*, Vol. I, M. Artin and J. Tate (Eds.), pp. 327–352, Birkhäuser, Boston, 1983.
- [13] P. Tzermias, The Manin-Mumford conjecture: a brief survey, *Bull. London Math. Soc.* **32** (2000), 641–652.
- [14] U. Zannier, Some problems of unlikely intersections in arithmetic and geometry, *Ann. Math. Studies* no. 181, Princeton Univ. Press, Princeton N.J., 2012.

Graduate School of Mathematical Sciences
University of Tokyo (Emeritus)
Komaba, Meguro-ku, Tokyo 153-8914
Japan
e-mail: noguchi@ms.u-tokyo.ac.jp

³⁾ The author is glad to acknowledge that the present work was inspired by the talks given by J. Pila and E. Ullmo at the Cetraro Conference 2017 referred at the end of §1, and the talk of Ullmo drew his attention to the preprint [10].

UTMS

- 2015–4 Koya SAKAKIBARA: *Analysis of the dipole simulation method for two-dimensional Dirichlet problems in Jordan regions with analytic boundaries* .
- 2015–5 Junjiro NOGUCHI: *A scalar associated with the inverse of some Abelian integrals on open Riemann surfaces and a ramified Riemann domain* .
- 2015–6 Tomoya KEMMOCHI: *Discrete maximal regularity for abstract Cauchy problems*.
- 2015–7 Tatsu-Hiko MIURA: *Zero width limit of the heat equation on moving thin domains*.
- 2016–1 Norikazu SAITO and Takiko SASAKI: *Finite difference approximation for nonlinear Schrödinger equations with application to blow-up computation*.
- 2016–2 Takiko SASAKI: *Regularity and singularity of blow-up curve for $\partial_t^2 u - \partial_x^2 u = |\partial_t u|^p$* .
- 2016–3 Atsushi KAWAMOTO: *Lipschitz stability estimates in inverse source problems for a fractional diffusion equation of half order in time by Carleman estimates*.
- 2016–4 Yoshikazu GIGA and Naoto KAJIWARA: *On a resolvent estimate for bidomain operators and its applications*.
- 2016–5 Yoshikazu GIGA and Tokinaga NAMBA : *Well-posedness of Hamilton-Jacobi equations with Caputo's time-fractional derivative*.
- 2017–1 Junjiro NOGUCHI: *A remark on Oka's coherence without Weierstrass' preparation theorem and the Oka theory* .
- 2017–2 Shigeo KUSUOKA: *Some regularity estimates for Diffusion semigroups with Dirichlet boundary conditions I*.
- 2017–3 Junjiro NOGUCHI: *An application of the value distribution theory for semi-abelian varieties to problems of Ax-Lindemann and Manin-Mumford types*.

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012