Japan. J. Math. 15, 121–167 (2020) DOI: 10.1007/s11537-019-1650-8



Hankel transform, Langlands functoriality and functional equation of automorphic *L*-functions*

Ngô Bảo Châu

Received: 2 February 2018 / Revised: 20 January 2019 / Accepted: 29 September 2019 Published online: 18 January 2020 © The Mathematical Society of Japan and Springer Japan KK, part of Springer Nature 2020

Communicated by: Takeshi Saito

Dedicated to the memory of Prof. Hiroshi Saito, with affection

Abstract This is a survey on recent works of Langlands's work on functoriality conjectures and related works including the works of Braverman and Kazhdan on the functional equation of automorphic L-functions. Efforts have been made to carry out in complete generality the construction of the L-monoid, and certain a kernel which is, we believe, related to the elusive Hankel kernel.

Keywords and phrases: Langlands functoriality, Hankel transform, functional equation

Mathematics Subject Classification (2010): 11M99, 22E35

B.C. NGÔ

Department of Mathematics, University of Chicago, 5734 University Avenue, Chicago, IL 60637-1514, USA (e-mail: ngo@uchicago.edu)

^{*} This article is based on the 18th Takagi Lectures that the author delivered at The University of Tokyo on November 5–6, 2016.