

# An operator algebra approach to the classification of certain fusion categories III

Masaki Izumi

Graduate School of Science, Kyoto University

August 3, 2016 at Sendai

# Classification of near-group categories

Let  $\mathcal{C} \subset \text{End}_0(M)$  be a near-group category with a finite group  $G$ .

$$[\rho^2] = \sum_{g \in G} [\alpha_g] + m[\rho].$$

Assume  $G$  is non-trivial and  $m \neq 0$ .

If  $d = d(\rho) = \frac{m + \sqrt{m^2 + 4|G|}}{2}$  is irrational, then  $G$  is abelian and  $m$  is a multiple of  $|G|$ .

Moreover, the categorifications of  $R(G, m)$  are completely classified by explicit polynomial equations.

A quadratic form on  $G$  appears in the polynomial equations:

$$\langle g, h \rangle = a(g)a(h)\overline{a(g+h)}, \quad a(-g) = a(g),$$

$$c^3 = \frac{1}{\sqrt{|G|}} \sum_{g \in G} \overline{a(g)}.$$

# Character formula

$$\alpha_g \circ \rho = \rho.$$

$$S_e \in (\text{id}, \rho^2), S_g = \alpha_g(S_e) \in (\alpha_g, \rho^2).$$

$$U(g) \in (\rho, \rho\alpha_g), U(g)S_e = S_e.$$

$$(\rho^2, \rho^2) = \bigoplus_{g \in G} \mathbb{C}S_gS_g^* \oplus B(\mathcal{K}),$$

$$U(g) = \sum_{h \in G} \chi_h(g)S_hS_h^* + U_{\mathcal{K}}(g),$$

where  $\mathcal{K} = (\rho, \rho^2)$ .

$$d = \frac{m + \sqrt{m^2 + 4|G|}}{2} \in \mathbb{R} \setminus \mathbb{Q} \Rightarrow$$

$$\bigoplus_{h \in G} \chi_h \cong \lambda, \quad U_{\mathcal{K}} \cong \frac{m}{|G|} \lambda.$$

# Irrational case

## Lemma

$\langle g, h \rangle := \chi_h(g)$  is a non-degenerate (in fact, symmetric) bicharacter.

## Proof.

$$\alpha_h(U(g)) \in (\rho, \rho\alpha_g) \Rightarrow \alpha_h(U(g)) = \exists c(g, h)U(g).$$

$$\begin{aligned} c(g, h)S_e &= c(g, h)U(g)S_e = \alpha_h(U(g))S_e \\ &= \alpha_h(U(g)S_{-h}) = \chi_{-h}(g)\alpha_h(S_{-h}) = \chi_{-h}(g)S_e, \end{aligned}$$

and  $\chi_{-h}(g) = c(g, h)$  is a non-degenerate character. □

$U_{\mathcal{K}}(g) \in B(\mathcal{K})$  is given by  $\mathcal{K} \ni T \mapsto U(g)T$ .

Definition of two other representations on board.

## Irrational case (continued)

### Definition

Let  $\mathcal{H}(G)$  be the universal  $C^*$ -algebra generated by three unitary representations  $v_0, v_1, v_2$  of  $G$ , and a unitary  $w$  of period 3 satisfying

$$v_{i+1}(g)v_i(h) = \langle h, g \rangle v_i(h)v_{i+1}(g),$$

$$w^*v_i(g)w = v_{i+1}(g),$$

where  $i \in \mathbb{Z}/3\mathbb{Z}$ .

We have a representation of  $\mathcal{H}(G)$  in  $\mathcal{K} = (\rho, \rho^2)$ .

# Irrational case (continued)

## Lemma

$\exists 3|G|$  irreducible representations of  $\mathcal{H}(G)$ , realized in  $\ell^2(G)$  as

$$\pi_{a,c}(v_0(g))f(h) = \langle g, h \rangle f(h),$$

$$\pi_{a,c}(v_1(g))f(h) = f(h + g),$$

$$\pi_{a,c}(v_2(g))f(h) = a(h)\overline{a(h-g)}f(h-g),$$

$$\pi_{a,c}(w)f(h) = \frac{c}{\sqrt{n}} \sum_k a(h)\overline{\langle h, k \rangle} f(k),$$

where  $a : G \rightarrow \mathbb{T}$  and  $c \in \mathbb{T}$  satisfy

$$a(g+h)\langle g, h \rangle = a(g)a(h),$$

$$c^3 \sum_{g \in G} a(g) = \sqrt{n}.$$

# Quadratic categories with $(G, \tau, m)$

## Definition

Let  $G$  be a finite group,  $\tau \in \text{Aut}(G)$  be an involution, and let  $m \in \mathbb{N}$ .

A **quadratic category of type  $(G, \tau, m)$**  is a fusion category  $\mathcal{C}$  with  $\text{Irr}(\mathcal{C}) = G \sqcup \{g \otimes \rho\}_{g \in G}$ , satisfying

$$[g][h] = [gh], \quad g, h \in G,$$

$$[g][\rho] = [\rho][g^\tau],$$

$$[\rho]^2 = [\text{id}] \oplus m \sum_{g \in G} [g][\rho].$$

The even part of the Haagerup subfactor is a quadratic category of type  $(\mathbb{Z}_3, -1, 1)$ .

Asaeda-Haagerup subfactor can be constructed from a quadratic category of type  $(\mathbb{Z}_4, -1, 2)$ .

# Obstructions

Let  $\mathcal{C} \subset \text{End}_0(M)$  be a quadratic category of type  $(G, \tau, m)$ .

$$\begin{aligned}\alpha_g \circ \alpha_h &= \text{Ad}^\exists U_{g,h} \circ \alpha_{gh}, \\ \rho \circ \alpha_{g\tau} &= \text{Ad}^\exists V_g \circ \alpha_g \circ \rho.\end{aligned}$$

We seek obstructions to making  $U_{g,h} = 1$  and  $V_g = 1$ .

A quadratic category of type “ $(G, \tau, 0)$ ” is  $\text{Vec}_{G \rtimes_\tau \mathbb{Z}_2}^\omega$ .

Recall that the  $E_2$ -term of the spectral sequence computing  $H^*(G \rtimes_\tau \mathbb{Z}_2, \mathbb{T})$  is  $E_2^{p,q} = H^p(\mathbb{Z}_2, H^q(G, \mathbb{T}))$ .

We use this analogy to define invariants of quadratic categories of type  $(G, \tau, m)$ .



$\exists \omega \in Z^3(G, \mathbb{C}^\times)$  satisfying  $\alpha_g(U_{h,k})U_{g,hk} = \omega(g, h, k)U_{g,h}U_{gh,k}$ .

### Lemma

$\exists \xi(g, h) \in \mathbb{T}$  satisfying

$$\omega(g, h, k) = \omega(g^\tau, h^\tau, k^\tau)\xi(h, k)\xi(gh, k)^{-1}\xi(g, hk)\xi(g, h)^{-1}.$$

In particular,  $[\omega] \in H^3(G, \mathbb{T})^{\mathbb{Z}_2} = H^0(\mathbb{Z}_2, H^3(G, \mathbb{T})) = E_2^{0,3}$ .

$$\rho \circ \alpha_{g^\tau} \circ \alpha_{h^\tau} = \rho \circ \text{Ad } U_{g^\tau, h^\tau} \circ \alpha_{(gh)^\tau} = \text{Ad}(\rho(U_{g,h})V_{gh}) \circ \alpha_{gh} \circ \rho,$$

$$\rho \circ \alpha_{g^\tau} \circ \alpha_{h^\tau} = \text{Ad}(V_g \alpha_g(V_h)) \circ \alpha_g \circ \alpha_h \circ \rho = \text{Ad}(V_g \alpha_g(V_h)U_{g,h}) \circ \alpha_{gh} \circ \rho,$$

$$\Rightarrow \exists \xi(g, h) \in \mathbb{T} \text{ satisfying } V_g \alpha_g(V_h)U_{g,h} = \xi(g, h)\rho(U_{g,h})V_{gh}.$$

### Definition

$$c^{0,3}(\mathcal{C}) := [\omega] \in H^3(G, \mathbb{T})^\tau.$$

Assume  $c^{0,3}(\mathcal{C}) = 0$ .

We may assume  $\alpha$  is an action,  $\xi \in Z^2(G, \mathbb{T})$  and

$$\rho \circ \alpha_{g^\tau} = \text{Ad } V_g \circ \alpha_g \circ \rho,$$

$$V_g \alpha_g(V_h) = \xi(g, h) V_{gh}.$$

### Lemma

$\exists \eta(g) \in \mathbb{T}$  satisfying

$$\xi(g^\tau, h^\tau) \xi(g, h) = \eta(gh) \eta(g)^{-1} \eta(h)^{-1}.$$

In particular, the 2-cocycle  $\xi \in Z^2(G, \mathbb{C}^\times)$  gives a class in  $H^1(\mathbb{Z}_2, H^2(G, \mathbb{C}^\times)) = E_2^{1,2}$ .

Using rigidity, we get  $\overline{V}_g \in (\alpha_{g^{\tau-1}} \circ \rho, \rho \circ \alpha_{g^{-1}})$ , and  $\eta(g) \in \mathbb{T}$  satisfying  $\overline{V_{(g^\tau)^{-1}}} = \eta(g) V_g$ .

### Definition

Define  $c^{1,2}(\mathcal{C}) \in H^1(\mathbb{Z}_2, H^2(G, \mathbb{T}))$  by the class given by  $\xi$ .

## Condition for Cuntz algebra models

Assume further that  $\mathfrak{c}^{1,2}(\mathcal{C}) = 0$ .

Then we can choose  $V_g$  satisfying

$$\rho \circ \alpha_{g^\tau} = \text{Ad } V_g \circ \alpha_g \circ \rho,$$

$$V_g \alpha_g(V_h) = V_{gh}.$$

Thus  $\exists W \in \mathcal{U}(M)$  satisfying  $V_g = W^{-1} \alpha_g(W)$ , and


$$\text{Ad } W \circ \rho \circ \alpha_{g^\tau} = \alpha_g \circ \text{Ad } W \circ \rho.$$

Replacing  $\rho$  with  $\text{Ad } W \circ \rho$ , we get

$$\rho \circ \alpha_{g^\tau} = \alpha_g \circ \rho.$$

Summary: To obtain a Cuntz algebra model for  $\mathcal{C}$ , we need  $\mathfrak{c}^{0,3}(\mathcal{C}) = 0$  and  $\mathfrak{c}^{1,2}(\mathcal{C}) = 0$ .

# Vanishing theorem



The Haagerup subfactor is  $3^{\mathbb{Z}_3}$ .

## Theorem (I)

When a quadratic category  $\mathcal{C}$  of type  $(G, \tau, 1)$  comes from a  $3^G$ -subfactor, then  $\mathfrak{c}^{0,3}(\mathcal{C}) \in H^3(G, \mathbb{T})^{\mathbb{Z}_2}$  and  $\mathfrak{c}^{1,2}(\mathcal{C}) \in H^1(\mathbb{Z}_2, H^2(G, \mathbb{T}))$  vanish.

## Proof.

$$[\alpha_g][\kappa] = [\kappa] \Rightarrow \mathfrak{c}^{0,3}(\mathcal{C}) = 0.$$

$(\kappa, \rho\kappa) \ni T \mapsto V_g \alpha_g(T) \in (\kappa, \rho\kappa)$  gives a projective representation of  $G$  with  $\dim(\kappa, \rho\kappa) = |G| - 1 \Rightarrow \mathfrak{c}^{1,2}(\mathcal{C}) = 0.$  □

## Theorem (I)

Let  $\mathcal{C}$  be a spherical quadratic category with  $(G, \tau, m)$ .

If  $G$  is an odd group and  $m$  is an odd number, then  $G$  is abelian and  $g^\tau = g^{-1}$  for any  $g \in G$ .

Let  $(\pi, V_\pi)$  be an irreducible representation of  $K(\mathcal{C})$ .

Then the formal codegree  $f_\pi$  for  $\pi$  is defined by

$$f_\pi = \sum_{X \in \text{Irr}(\mathcal{C})} \text{Tr}(\pi(X))\pi(\overline{X}).$$

Since  $f_\pi$  commutes with  $\pi(X)$  for every  $X \in \text{Irr}(\mathcal{C})$ , it is a scalar.

## Theorem (Ostrik 2009)

If  $\mathcal{C}$  is spherical, there exists a simple object in the Drinfeld center  $\mathcal{Z}(\mathcal{C})$  whose dimension is  $\dim \mathcal{C} / f_\pi$ .

In particular,  $\dim \mathcal{C} / f_\pi$  is a cyclotomic integer.

## Lemma

If  $G$  and  $m$  are odd, for any non-trivial irreducible representation  $\pi$  of  $G$ ,  $\pi$  and  $\pi^\tau$  are inequivalent.

## Proof.

Suppose that  $\pi$  is a non-trivial irreducible representation of  $G$  with  $\pi \cong \pi^\tau$ . Then  $\pi$  extends to an irreducible representation  $\pi'$  of  $K(\mathcal{C})$  whose formal codegree is  $f_{\pi'} = 2|G|/\dim \pi$ , and

$$\frac{\dim \mathcal{C}}{f_{\pi'}} = \dim \pi + \frac{m|G| \dim \pi \dim \rho}{2}.$$

This is not an algebraic integer. □

## Definition

A **generalized Haagerup category** with a finite abelian group  $G$  is a quadratic category  $\mathcal{C}$  with  $(G, -1, 1)$  satisfying  $\mathfrak{c}^{0,3}(\mathcal{C}) = 0$  and  $\mathfrak{c}^{1,2}(\mathcal{C}) = 0$ .

Caution:

- (1) There exist two quadratic categories of type  $(\mathbb{Z}_3, -1, 1)$  with  $\mathfrak{c}^{0,3}(\mathcal{C}) \neq 0$ .
- (2) To construct the Asaeda-Haagerup subfactors, we need a quadratic category of type  $(\mathbb{Z}_4, -1, 2)$  and  $\mathfrak{c}^{0,3}(\mathcal{C}) = 0$ .

## Theorem

*Generalized Haagerup categories are completely classified by explicit polynomial equations.*

*More precisely, there exists a one-to-one correspondence between the equivalence classes of generalized Haagerup categories and the  $H^2(G, \mathbb{T}) \rtimes \text{Aut}(G)$ -orbits of the gauge equivalence classes of the solutions of the polynomial equations.*

*For a fixed solution, the stabilizer subgroup is isomorphic to the outer automorphism group of the corresponding category.*



# Polynomial equations for odd $G$

Variables:  $A(g, h) \in \mathbb{C}$  and  $\eta \in \mathbb{T}$  with  $\eta^3 = 1$ .

$$\sum_{h \in G} A(h, 0) = -\frac{\bar{\eta}}{d},$$

$$\sum_{h \in G} A(h - g, k) \overline{A(h - g', k)} = \delta_{g, g'} - \frac{\delta_{k, 0}}{d},$$

$$A(k, h) = \overline{A(h, k)},$$

$$A(h, k) = A(-k, h - k)\eta = A(k - h, -h)\bar{\eta},$$

$$\begin{aligned} & \sum_{l \in G} A(x + y, l) A(-x, l + p) A(-y, l + q) \\ &= A(p + x, q + x + y) A(q + y, p + x + y) - \frac{\delta_{x, 0} \delta_{y, 0}}{d}. \end{aligned}$$

# Solutions for the polynomial equations

$G$	# (sols/ $H^2(G, \mathbb{T}) \times \text{Aut}(G)$ )	With Q-system for $\text{id} \oplus \rho$
$\mathbb{Z}_2$	1	1
$\mathbb{Z}_3$	2	1
$\mathbb{Z}_4$	2	1
$\mathbb{Z}_2 \times \mathbb{Z}_2$	1	1
$\mathbb{Z}_5$	2	1
$\mathbb{Z}_6$	4	2
$\mathbb{Z}_7$	$\geq 2$	1
$\mathbb{Z}_8$	$\geq 1$	$\geq 1$
$\mathbb{Z}_4 \times \mathbb{Z}_2$	$\geq 1$	$\geq 1$
$\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$	?	?
$\mathbb{Z}_9$	$\geq 2$	2
$\mathbb{Z}_3 \times \mathbb{Z}_3$	?	0
$\mathbb{Z}_{10}$	?	?
$\mathbb{Z}_{11}$	$\geq 2$	2

# From generalized Haagerup to near-group

## Theorem

Let  $\mathcal{C} \subset \text{End}_0(M)$  be a quadratic category of type  $(G, -1, m)$  with an odd abelian  $G$  and  $\mathfrak{c}^{0,3}(\mathcal{C}) = [\omega]$ .

Let

$$\iota : M \otimes M^{\text{opp}} \hookrightarrow (M \otimes M^{\text{opp}}) \rtimes_{\alpha \otimes \alpha^{\text{opp}}} G.$$

Then  $\iota \circ (\rho \otimes \text{id}) \circ \bar{\iota}$  generates a near-group category with group  $\text{Irr}(D^\omega(G))$  and multiplicity  $m|G|^2$ .

From  $G = \mathbb{Z}_3$  and  $m = 1$ , we get a near group categories for  $\mathbb{Z}_3 \times \mathbb{Z}_3$  or  $\mathbb{Z}_9$  with multiplicity 9.

# Orbifold (de-equivariantization) of near-group categories I

## Theorem

Let  $\mathcal{C} \subset \text{End}(M)$  be a near-group category with  $m = |G|$ .  
If  $H$  is Lagrangian, i.e.  $H = H^\perp$  and  $a|_H = 1$ , then de-equivariantization of  $\mathcal{C}$  by  $H$  is a quadratic category of type  $(G/H, -1, 1)$ .

There is a unique near-group category for  $\mathbb{Z}_3 \times \mathbb{Z}_3$  with  $m = 9$ .  
It has two Lagrangians, giving the Haagerup category and Grossman-Snyder category.

There are two near-group categories for  $\mathbb{Z}_9$ .  
They have Lagrangian  $\mathbb{Z}_3$ , giving  $\mathcal{C}$  with non-trivial  $\mathfrak{c}^{0,3}(\mathcal{C})$ .

## Corollary

There exist exactly 4 quadratic categories type  $(\mathbb{Z}_3, -1, 1)$ .

# Twisted orbifold (de-equivariantization) of near-group categories

## Theorem

Let  $\mathcal{C} \subset \text{End}(M)$  be a near-group category with  $m = |G|$ .

Assume  $G = K \times H$  and  $H = \mathbb{Z}_2^{2l}$ .

Assume  $\exists \omega \in Z^2(H, \mathbb{T})$  such that  $\langle h_1, h_2 \rangle = \omega(h_1, h_2) \overline{\omega(h_2, h_1)}$  is non-degenerate on  $H$ .

Then  $\omega$ -twisted de-equivariantization of  $\mathcal{C}$  by  $H$  is a near-group category with group  $K$  and multiplicity  $2^l |K|$ .

There are two solutions for  $K = \mathbb{Z}_3$ ,  $H = \mathbb{Z}_2 \times \mathbb{Z}_2$  satisfying the above conditions, and they produce 2 near-group categories of  $\mathbb{Z}_3$  with multiplicity 6.