

The Whittaker Plancherel theorem

Nolan R. Wallach

Received: 8 July 2022 / Revised: 23 January 2023, 28 June 2023, 16 August 2023, 23 November 2023 / Accepted: 24 November 2023
Published online: 9 February 2024

© The Mathematical Society of Japan and Springer Nature Japan KK, part of Springer Nature 2024

Communicated by: Toshiyuki Kobayashi

Abstract. The purpose of this article is to give an exposition of a proof of the distributional form of the Whittaker Plancherel Theorem. The proof is an application of Harish-Chandra's Plancherel Theorem for real reductive groups and its exposition can be used as an introduction to Harish-Chandra's Plancherel Theorem. The paper follows the basic method in the author's original approach in his second volume on real reductive groups. An error in the calculation of the Whittaker Transform of a Harish-Chandra wave packet is fixed using a result of Raphaël Beuzart-Plessis.

Keywords and phrases: tempered, representation, isotypic component, Schwartz space, Harish-Chandra Schwartz Space, Whittaker functional, Jacquet integral

Mathematics Subject Classification (2020): Primary 22030; Secondary 22045

N.R. WALLACH

Department of Mathematics, University of California, San Diego, La Jolla, CA 92093, USA

(e-mail: ucsd.nwallach@gmail.com)