

Solutions of heteroclinic type for some classes of semilinear elliptic partial differential equations

By Paul H. RABINOWITZ

Abstract. This paper studies certain classes of equations of the form $-\Delta u = g(x, y, u)$ in an infinite strip (if $n = 2$) or cylinder (if $n > 2$). Variational arguments are used to establish the existence of solutions asymptotic to a pair of x -periodic states.

§1. Introduction

This paper studies the existence of solutions of heteroclinic type for 3 families of semi-linear elliptic partial differential equations. For $n \geq 2$, let \mathcal{D} denote a bounded domain in \mathbb{R}^{n-1} having a smooth boundary, $\partial\mathcal{D}$. Let $\Omega = \mathbb{R} \times \mathcal{D}$, an infinite cylinder if $n > 2$ and an infinite strip if $n = 2$. Points in Ω will be denoted by (x, y) , $x \in \mathbb{R}$, $y \in \mathcal{D}$. Consider the partial differential equation

$$(PDE) \quad -\Delta u = g(x, y, u) \quad (x, y) \in \Omega$$

together with the boundary conditions

$$(1.1) \quad u(x, y) = 0, \quad x \in \mathbb{R}, y \in \partial\mathcal{D}$$

or

$$(1.2) \quad \frac{\partial u}{\partial \nu}(x, y) = 0, \quad x \in \mathbb{R}, y \in \partial\mathcal{D}.$$

1991 *Mathematics Subject Classification.* 35B40, 35J20, 35J65.

This research was sponsored in part by the U. S. Army Research Office under contract #DAAL03-87-K-0043 and the National Science Foundation under grant #MCS-8110556. Any reproduction for the purpose of the U.S. government is permitted.

In (1.2), $\nu = \nu(y)$ denotes the outward pointing normal to $\partial\mathcal{D}$.

Suppose that g is periodic in x and there is a corresponding family, \mathcal{M} , of solutions of (PDE) and (1.1) or (1.2) which are periodic in x and are minimizers of the associated variational problem. The main question of interest here is in the existence of solutions of (PDE) together with (1.1) or (1.2) which approach different members of \mathcal{M} as $x \rightarrow \pm\infty$. If $n = 1$, (PDE) reduces to an ordinary differential equation in x and such solutions are then heteroclinic to a pair of different elements of \mathcal{M} .

The existence of solutions of heteroclinic type will be established for three classes of problems. The first is for the Neumann problem when g is also periodic in u . The second is again for (1.2) for a family of g 's where

$$(1.3) \quad G(x, y, u) = \int_0^u g(x, y, t) dt$$

has a finite number of global maxima u_1, \dots, u_m independently of (x, y) and $G(x, y, u_i) = 0$, $1 \leq i \leq m$. The third case is for the Dirichlet problem (1.1) when G is appropriately coercive in u as $|u| \rightarrow \infty$.

The first type of problem was suggested by some recent work on heteroclinic solutions of reversible Hamiltonian systems. See [1–2] and also the papers [3–4] of Bolotin. We mention also the somewhat related work of Moser [5] on minimal solutions of a variational problem on a torus. The third class of problems was motivated by earlier research of Kirchgässner [6] who studied a problem arising in the theory of water waves—see also Turner [7]:

$$(1.4) \quad -\Delta u = \lambda a(y)u - f(x, y, u, \lambda)$$

for $x, y \in \mathbb{R}$ and $|y| < 1$ with the boundary conditions

$$(1.5) \quad u(x, \pm 1) = 0$$

where $a(y) > 0$, $f(x, y, 0, \lambda) = 0 = f_u(x, y, 0, \lambda)$, and λ is near λ_1 , the smallest eigenvalue of the related linearized eigenvalue problem

$$\begin{cases} \frac{-d^2v}{dy^2} = \lambda a(y)v, & |y| < 1 \\ v(\pm 1) = 0 \end{cases}.$$

Kirchgässner used the Center Manifold Theorem to prove the existence of a small amplitude solution of (1.4)–(1.8) which tends to x -independent states as $|x| \rightarrow \infty$. Further studies of such problems were also made by Kirchgässner and his collaborators. See e.g. [8–9].

The approach taken here is a global variational one that is the analogue for (PDE) and (1.1) or (1.2) of the arguments used in [1–2]. The more complicated cases of g periodic in u will be treated in §2. Then the remaining cases will be sketched in §3.

§2. The periodic case

Periodic in u nonlinearities g for (1.1) will be considered in this section. Suppose that g satisfies

- (g_1) $g \in C^1(\bar{\Omega} \times \mathbb{R}, \mathbb{R})$,
- (g_2) $g(x, y, u)$ is even and 1-periodic in x ,
- (g_3) $G(x, y, u)$ is 1-periodic in u .

It is not important that the periods of g in x and u are the same. This assumption is made merely for notational convenience.

The first step in finding heteroclinic type solutions of (PDE) and (1.2) is to establish the existence of a large class, \mathcal{M} , of solutions of (PDE) and (1.2) which are 1-periodic in x . They will be obtained as minima of a corresponding variational problem. Thus let G be as in (1.3). Let $\Omega_1 = [0, 1] \times \mathcal{D}$ and $|\Omega_1| = \text{volume of } \Omega_1$. Let $E_1 = \{u \in W^{1,2}(\Omega_1) \mid u \text{ is 1-periodic in } x\}$. For $u \in E_1$, set

$$(2.1) \quad I_1(u) = \int_{\Omega_1} \left(\frac{1}{2} |\nabla u|^2 - G(x, y, u) \right) dx dy.$$

Then by (g_1), (g_3), and standard results [10], $I \in C^1(E_1, \mathbb{R})$. Note that by (g_3), for all $k \in \mathbb{Z}$ and $u \in E_1$,

$$(2.2) \quad I_1(u + k) = I_1(u),$$

i.e. I_1 has a natural \mathbb{Z} symmetry.

Let

$$(2.3) \quad c_1 = \inf_{u \in E_1} I_1(u).$$

PROPOSITION 2.4. *There is a $\bar{u} \in E_1$ such that $I_1(\bar{u}) = c_1$.*

PROOF. Let (u_n) be a minimizing sequence for (2.3) with

$$(2.5) \quad I_1(u_m) \leq K.$$

By (2.2), $(u_m + k_m)$ is also a minimizing sequence for (2.3) for any $(k_m) \subset \mathbb{Z}$. Hence it can be assumed that

$$(2.6) \quad 0 \leq [u_m] \equiv \frac{1}{|\Omega_1|} \int_{\Omega_1} u_m dx dy < 1.$$

We claim that (u_m) is bounded in E_1 . The weak lower semicontinuity of I_1 , then implies that along a subsequence, $u_m \rightarrow \bar{u} \in E_1$ weakly in E_1 and $I_1(\bar{u}) = c_1$.

By (2.5) and (g_1) , (g_3) , for some constant K_1 ,

$$(2.7) \quad \|\nabla u_m\|_{L^2(\Omega_1)}^2 \leq K_1 \|u_m\|_{L^1(\Omega_1)} + 2K$$

for all $m \in \mathbb{N}$. Elementary calculus arguments show

$$(2.8) \quad \|u_m\|_{L^1(\Omega_1)} \leq |\Omega_1| |[u_m]| + K_2 \|\nabla u_m\|_{L^2(\Omega_1)}$$

and

$$(2.9) \quad \|u_m\|_{L^2(\Omega_1)} \leq K_3 (|[u_m]| + \|\nabla u\|_{L^2(\Omega_1)}).$$

Combining (2.7)–(2.9) shows (u_m) is bounded in E_1 and Proposition 2.4 is proved. \square

COROLLARY 2.10. *\bar{u} is a classical solution of (PDE) and (1.2) with $\bar{u}(x+1, y) = \bar{u}(x, y)$ for all $(x, y) \in \Omega$.*

PROOF. Since \bar{u} minimizes I_1 on E_1 , $I_1 \in C^1(E_1, \mathbb{R})$, and (g_1) , (g_3) are satisfied, standard arguments imply \bar{u} is a classical solution of (PDE) and (1.2) holds. The periodicity in x follows since $\bar{u} \in E_1$. \square

By (2.2), whenever \bar{u} minimizes I_1 on E_1 , so does $\bar{u} + k$ for all $k \in \mathbb{Z}$.
Let

$$(2.11) \quad \mathcal{M} = \{u \in E_1 \mid I_1(u) = c_1\}.$$

Before constructing solutions of (PDE) of heteroclinic type, some further properties of \mathcal{M} must be obtained. Let

$$(2.12) \quad \bar{c}_1 = \inf_{u \in W^{1,2}(\Omega_1)} I_1(u).$$

PROPOSITION 2.13. $\bar{c}_1 = c_1$.

PROOF. Since $E_1 \subset W^{1,2}(\Omega_1)$, $\bar{c}_1 \leq c_1$. To prove equality, suppose $\bar{c}_1 < c_1$. Then there is a $u \in W^{1,2}(\Omega_1)$ such that $I_1(u) < c_1$. Writing

$$(2.14) \quad \begin{aligned} I_1(u) &= \int_0^{\frac{1}{2}} \int_{\mathcal{D}} \mathcal{L}(u) dx dy + \int_{\frac{1}{2}}^1 \int_{\mathcal{D}} \mathcal{L}(u) dx dy \\ &\equiv \alpha + \beta \end{aligned}$$

where

$$(2.15) \quad \mathcal{L}(u) = \frac{1}{2} |\nabla u|^2 - G(x, y, u),$$

it follows that either α or $\beta < c_1/2$. Suppose e.g. $\alpha < c_1/2$. Define

$$(2.16) \quad \begin{aligned} v(x, y) &= u(x, y) & 0 \leq x \leq \frac{1}{2}, y \in \mathcal{D} \\ &= u(1-x, y) & \frac{1}{2} \leq x \leq 1, y \in \mathcal{D} \end{aligned}$$

and extend v to $\mathbb{R} \times \mathcal{D}$ as a 1-periodic function in x . Then $v \in E_1$. Since G is even and 1-periodic in x , by (g_2) ,

$$(2.17) \quad I_1(v) = 2\alpha < c_1$$

contrary to (2.3) and Proposition 2.4. Thus $\bar{c}_1 = c_1$. \square

PROPOSITION 2.18. *If $u \in W^{1,2}(\Omega_1)$ and $I_1(u) = c_1$, then $u \in \mathcal{M}$ and u is even in x .*

PROOF. Suppose $u \in W^{1,2}(\Omega_1)$ with $I_1(u) = c_1$. Define v as in (2.16) so $v \in E_1$ and is even in x . By (2.3), (2.12), and Proposition 2.13, v and u are critical points of I_1 and therefore both are classical solutions of (PDE) and (1.2) with $v = u$ on $[0, \frac{1}{2}] \times \mathcal{D}$. Let $w = u - v$.

Then w satisfies

$$(2.19) \quad -\Delta w = g(x, y, u) - g(x, y, v) \equiv b(x, y)w$$

where

$$b(x, y) = \frac{g(x, y, u) - g(x, y, v)}{u - v} \quad \text{if} \quad u(x, y) \neq v(x, y)$$

$$= g_u(x, y, u) \quad \text{if} \quad u(x, y) = v(x, y)$$

and

$$(2.20) \quad \frac{\partial w}{\partial \nu}(x, \partial \mathcal{D}) = 0.$$

Note that b is continuous and $w(x, y) = 0$ for $0 \leq x \leq \frac{1}{2}$ and $y \in \mathcal{D}$. But then a local unique continuation theorem for elliptic equations—see e.g. Nirenberg [11]—and a continuation argument imply $w(x, y) \equiv 0$, $x \in [0, 1]$, $y \in \bar{\mathcal{D}}$, i.e. $u \equiv v$ in Ω_1 . \square

A special case of interest is when G is independent of x . Then the elements of \mathcal{M} also possess this property:

PROPOSITION 2.21. *If G is independent of x and $w \in \mathcal{M}$, then w is independent of x .*

PROOF. For each $\theta \in \mathbb{R}$, set

$$(2.22) \quad I_1^\theta(u) = \int_{\theta - \frac{1}{2}}^{\theta + \frac{1}{2}} \int_{\mathcal{D}} \left(\frac{1}{2} |\nabla u|^2 - G(y, u) \right) dx dy.$$

If $u \in E_1$, then $I_1^\theta(u) = I_1(u)$. Therefore

$$(2.23) \quad c_1 = \inf_{E_1} I_1^\theta(u).$$

Since G is independent of x , the argument of Proposition 2.13 shows any minimizer w_1 of I_1^θ is even about $x = \theta$ for each $\theta \in \mathbb{R}$. Hence w is independent of x . But the minimizers of I_1^θ are the minimizers of I_1 . \square

For later purposes, for $k \in \mathbb{N}$, we must also study solutions of (PDE) in

$$(2.24) \quad E_k = \{u \in W^{1,2}(\mathbb{R} \times \mathcal{D}) \mid u \text{ is } k \text{ periodic in } x\}.$$

Define

$$(2.25) \quad I_k(u) = \int_0^k \int_{\mathcal{D}} \mathcal{L}(u) dx dy$$

and

$$(2.26) \quad c_k = \inf_{E_k} I_k(u).$$

PROPOSITION 2.27. $c_k = kc_1$ and if u is a critical point of I_k on E_k , then $u \in \mathcal{M}$.

PROOF. Any $u \in E_1$ can be considered to be an element of E_k with $I_k(u) = kI_1(u)$. Therefore $c_k \leq kc_1$. Let u_k be a critical point of I_k corresponding to c_k . (It exists via the argument of Proposition 2.4.) Suppose $c_k < kc_1$. As in Proposition 2.4, consider the restrictions of u_k to the cylinders $[0, \frac{1}{2}] \times \mathcal{D}$, $[\frac{1}{2}, 1] \times \mathcal{D}$, \dots , $[k - \frac{1}{2}, k] \times \mathcal{D}$. Choose the restriction which makes the smallest contribution to $I_k(u_k)$. This contribution must be less than $c_1/2$. Extending this restriction of u_k evenly and then 1-periodically about an endpoint yields $v \in E_1$ with $I_1(v) < c_1$, a contradiction. Therefore $c_k = kc_1$. The argument of Proposition 2.18 shows $u_k \in \mathcal{M}$. \square

Proposition 2.4 and (g_3) show \mathcal{M} is a sizable set. We assume it is not too large in the following sense:

(\mathcal{M}) \mathcal{M} consists of isolated points.

If (\mathcal{M}) is not satisfied, perturbing the problem slightly produces a new functional for which (\mathcal{M}) holds. E.g. if $\bar{u} \in \mathcal{M}$, replace $G(x, y, u)$ by $\hat{G}(x, y, u) = G(x, y, u) - \epsilon(u - \bar{u}(x, y))^2$ if $(x, y) \in \mathbb{R} \times \bar{\mathcal{D}}$ and $|u - \bar{u}(x, y)| \leq \frac{1}{4}$; $\hat{G}(x, y, u) = G(x, y, u) - \epsilon(u - (\bar{u}(x, y) + 1))^2$ if $(x, y) \in \mathbb{R} \times \bar{\mathcal{D}}$ and $|u - (\bar{u}(x, y) + 1)| \leq \frac{1}{4}$, etc. and extend \hat{G} to the rest of $\mathbb{R} \times \bar{\mathcal{D}} \times \mathbb{R}$ appropriately so that \hat{G} satisfies (g_1) – (g_3) . Then for the associated functional, $\hat{I}_1, \hat{I}_1(\bar{u}) = I_1(\bar{u}) = c_1$ but if $u \notin \bar{u} + \mathbb{Z}$, $\hat{G}(u) < G(u)$ and therefore $\hat{I}_1(u) > I_1(u)$. Therefore the only minima of \hat{I}_1 are $\{\bar{u} + k \mid k \in \mathbb{Z}\}$ and \mathcal{M} is satisfied for the \hat{I}_1 problem.

Our goal now is to show that for any $v \in \mathcal{M}$, there is a solution U of (PDE) such that $\|U - v\|_{L^\infty([n, n+1] \times \mathcal{D})} \rightarrow 0$ as $n \rightarrow -\infty$, i.e. $U \rightarrow v$ uniformly in this sense as $x \rightarrow -\infty$ and similarly $U \rightarrow \mathcal{M} \setminus \{v\}$ as $x \rightarrow \infty$. By (\mathcal{M}) , this means there is a $w \in \mathcal{M} \setminus \{v\}$ such that $U \rightarrow w$ uniformly as $x \rightarrow \infty$. Thus U is a solution of (PDE) of heteroclinic type. A variational argument will be used to prove this result after some further technical preliminaries. Let $B_\rho(v) = \{u \in W^{1,2}(\Omega_1) \mid \|u - v\|_{W^{1,2}(\Omega_1)} < \rho\}$ and $N_\rho(S) = \{u \in W^{1,2}(\Omega_1) \mid \|u - S\|_{W^{1,2}(\Omega_1)} \leq \rho\}$.

PROPOSITION 2.28. *Suppose that g satisfies (g_1) – (g_3) and (\mathcal{M}) holds. Then there is a constant $\rho_0 > 0$ such that if $0 < \rho < \rho_0$,*

- (i) $B_\rho(v) \cap B_\rho(w) = \emptyset$ for all $v \neq w \in \mathcal{M}$.
- (ii) $I_1(u) > c_1$ for all $u \in W^{1,2}(\Omega_1) \setminus \mathcal{M}$.
- (iii) There is an $\alpha(\rho) > 0$ such that

$$I_1(u) \geq c + \alpha(\rho) \quad \text{for all} \quad u \in W^{1,2}(\Omega_1) \setminus N_\rho(\mathcal{M}).$$

PROOF. Let

$$(2.29) \quad \gamma = \inf\{\|v - w\|_{W^{1,2}(\Omega_1)} \mid v \neq w \in \mathcal{M}\}.$$

We claim $\gamma > 0$ and therefore (i) follows with e.g. $\rho_0 = \gamma/2$. To see that $\gamma > 0$, let $u \in \mathcal{M}$ with

$$(2.30) \quad -1 \leq [u] < 2.$$

Then as in the proof of Proposition 2.4, there is a $K = K(c_1)$ such that $\|u\|_{W^{1,2}(\Omega_1)} \leq K$. Since $u \in \mathcal{M}$, it is a classical solution of (PDE) and

(1.2). Therefore by standard elliptic regularity theory arguments, for any $\beta \in (0, 1)$, there is a $K_\beta = K_\beta(c_1)$ such that $\|u\|_{C^{2,\beta}} \leq K_\beta$ where $C^{2,\beta}$ denotes the set of u on $[0, 1] \times \mathcal{D}$ which are 1-periodic in x , C^2 , and whose second derivatives are Hölder continuous of order β . Now if $\gamma = 0$, there are sequences $(v_j) \neq (w_j) \subset \mathcal{M}$ such that

$$(2.31) \quad \|v_j - w_j\|_{W^{1,2}(\Omega_1)} \rightarrow 0$$

as $j \rightarrow \infty$. As in (2.6), it can be assumed that

$$(2.32) \quad 0 \leq [w_j] < 1$$

for all $j \in \mathbb{N}$. By (2.31), $[v_j]$ satisfies (2.30) for large j . Consequently both v_j and w_j converge in C^2 along a subsequence to $u \in \mathcal{M}$. But then u is not an isolated solution of (PDE) and (1.2), contrary to (\mathcal{M}) . Thus $\gamma > 0$ and (i) holds.

Property (ii) follows from the definition of c_1 and Propositions 2.13 and 2.18.

To prove (iii), an indirect argument will again be employed. If (iii) is false, there is a sequence $(u_m) \subset W^{1,2}(\Omega_1) \setminus N_\rho(\mathcal{M})$ such that $0 \leq [u_m] < 1$ and

$$(2.33) \quad I_1(u_m) \rightarrow c_1.$$

As for (i), (2.33) implies (u_m) is bounded in $W^{1,2}(\Omega_1)$. Hence there is a $u \in W^{1,2}(\Omega_1)$ such that, along a subsequence, $u_m \rightarrow u$ weakly in $W^{1,2}(\Omega_1)$. Since I_1 is weakly lower semicontinuous, $I_1(u) = c_1$. Consequently $u \in \mathcal{M}$. Set $\varphi_m = u - u_m$. Note that $\|\varphi_m\|_{W^{1,2}(\Omega_1)} \geq \rho$ and $\varphi_m \rightarrow 0$ weakly in $W^{1,2}(\Omega_1)$ along the subsequence. Therefore

$$(2.34) \quad \begin{aligned} I_1(u_m) &= I_1(u) + \int_{\Omega_1} \left[\frac{1}{2} (|\nabla \varphi_m|^2 + \varphi_m^2) - \nabla u \cdot \nabla \varphi_m \right. \\ &\quad \left. - G(x, y, u - \varphi_m) + G(x, y, u) - \frac{1}{2} \varphi_m^2 \right] dx dy \geq c_1 + \frac{1}{2} \rho^2 \\ &\quad - \int_{\Omega_1} \left[\nabla u \cdot \nabla \varphi_m + G(x, y, u - \varphi_m) \right. \\ &\quad \left. - G(x, y, u) + \frac{1}{2} \varphi_m^2 \right] dx dy. \end{aligned}$$

Since $\varphi_m \rightarrow 0$ weakly in $W^{1,2}(\Omega_1)$, $\varphi_m \rightarrow 0$ in $L^2(\Omega_1)$ along the subsequence and

$$(2.35) \quad \int_{\Omega_1} G(x, y, u - \varphi_m) dx dy \rightarrow \int_{\Omega_1} G(x, y, u) dx dy.$$

Thus the right hand side of (2.34) tends to 0 as $m \rightarrow \infty$, along the subsequence, i.e.

$$(2.36) \quad \lim_{m \rightarrow \infty} I_1(u_m) \geq c_1 + \frac{1}{2}\rho^2$$

contrary to (2.33). Thus there is an $\alpha(\rho) > 0$ as claimed and (iii) holds. \square

REMARK 2.37.

- (i) The argument of (i) of Proposition 2.28 implies that \mathcal{M}/\mathbb{Z} is compact. This fact together with (\mathcal{M}) shows that \mathcal{M}/\mathbb{Z} is a finite set. Therefore setting

$$(2.38) \quad \bar{\gamma} = \inf\{\|u - v\|_{L^2(\Omega_1)} \mid u = v \in \mathcal{M}\},$$

it follows that $\bar{\gamma} > 0$.

- (ii) Choosing $\alpha(\rho)$ still smaller if necessary, the argument of (iii) of Proposition 2.28 shows that $I_2(u) \geq 2c_1 + \alpha(\rho)$ for all $u \in W_{loc}^{1,2}(\mathbb{R} \times \bar{\mathcal{D}}) \setminus N_\rho^2(\mathcal{M})$ where $N_\rho^2(\mathcal{M})$ denotes a uniform neighborhood of \mathcal{M} in $W^{1,2}([0, 2] \times \bar{\mathcal{D}})$.

Let $u \in W_{loc}^{1,1}(\mathbb{R} \times \bar{\mathcal{D}})$ and $k \in \mathbb{Z}$. Set

$$(2.39) \quad P_k u = u|_{[k, k+1] \times \bar{\mathcal{D}}}.$$

Then $P_k u$ can be identified with

$$w(x, y) = P_k u(x + k, y) \in W^{1,2}(\Omega_1).$$

This identification will be clear from the context and will not be made explicitly in what follows.

Now the variational problem which will be used to find solutions of (PDE) of heteroclinic type can be formulated. For $v \in \mathcal{M}$, set

$$(2.40) \quad \Gamma^-(v) = \{U \in W_{loc}^{1,2}(\mathbb{R} \times \mathcal{D}) \mid \|P_k U - v\|_{L^2(\Omega_1)} \rightarrow 0 \\ \text{as } k \rightarrow -\infty \quad \text{and} \quad \|P_k U - (\mathcal{M} \setminus \{v\})\|_{L^2(\Omega_1)} \rightarrow 0 \\ \text{as } k \rightarrow \infty\}.$$

It is clear from the definition that $\Gamma^-(v) \neq \emptyset$. For $k \in \mathbb{Z}$ and $U \in \Gamma^-(v)$, set

$$(2.41) \quad a_k(U) = \int_k^{k+1} \int_{\mathcal{D}} (\mathcal{L}(U) - \mathcal{L}(v)) dx dy.$$

By the definition of \mathcal{M} and Proposition 2.13,

$$(2.42) \quad a_k(U) \geq 0$$

for all such k and U and $a_k(U) = 0$ if and only if $P_k U \in \mathcal{M}$.

Finally define

$$(2.43) \quad J(U) = \sum_{k \in \mathbb{Z}} a_k(U)$$

and set

$$(2.44) \quad c = \inf_{U \in \Gamma^-(v)} J(U).$$

At first glance, it may seem that $J(U)$ equals

$$(2.45) \quad \int_{\mathbb{R} \times \mathcal{D}} (\mathcal{L}(U) - \mathcal{L}(v)) dx dy.$$

However there are functions $U \in \Gamma^-(v)$ such that $J(U) < \infty$ but for which the integral in (2.45) is not conditionally convergent.

Now the main results for this section can be stated

THEOREM 2.46. *Let (g_1) – (g_3) and (\mathcal{M}) be satisfied. Then for each $v \in \mathcal{M}$, there is a $U \in \Gamma^-(v)$ such that $J(U) = c$.*

COROLLARY 2.47. U is a classical solution of (PDE) with $P_k U \rightarrow v$ in $C^2(\Omega_1)$ as $k \rightarrow -\infty$ and $P_k U \rightarrow \mathcal{M} \setminus \{v\}$ in $C^2(\Omega_1)$ as $k \rightarrow \infty$.

PROOF OF THEOREM 2.46. The proof will be divided into four main steps: (A) Construction of a minimizing sequence for (2.44) which converges weakly to some $U \in W_{loc}^{1,2}(\mathbb{R} \times \bar{\mathcal{D}})$; (B) Obtaining the asymptotic behavior for U as $x \rightarrow -\infty$; (C) Obtaining the asymptotic behavior for U as $x \rightarrow \infty$; (D) Showing that U minimizes J on $\Gamma^-(v)$.

(A) A convergent minimizing sequence

For $k \in \mathbb{Z}$ and $W \in \Gamma^-(v)$, set

$$(2.48) \quad \tau_k W(x, y) = W(x - k, y).$$

Then $\tau_k W \in \Gamma^-(v)$ and by (g_1) – (g_2) ,

$$(2.49) \quad J(\tau_k W) = J(W).$$

Now let (W_m) be a minimizing sequence for (2.44). By (2.49), $(\tau_{k(m)} W_m)$ is also a minimizing sequence for any $(k(m)) \subset \mathbb{Z}$. Let

$$(2.50) \quad \bar{\rho} \in (0, \bar{\gamma}/3)$$

where $\bar{\gamma}$ is defined in (2.38). For each m , choose $k(m)$ so that

$$(2.51) \quad \|P_j \tau_{k(m)} W_m - v\|_{L^2(\Omega_1)} \leq \bar{\rho}$$

for all $j < 0$ and

$$(2.52) \quad \|P_0 \tau_{k(m)} W_m - v\|_{L^2(\Omega_1)} > \bar{\rho}.$$

This is possible via the definition of $\Gamma^-(v)$ and (2.38). Without loss of generality, it can be assumed that $k(m) = 0$ for all m . Since $J(W_m) \rightarrow c$, there is an $M > 0$ such that

$$(2.53) \quad J(W_m) \leq M$$

for all $m \in \mathbb{N}$. By (2.51), for any $\ell \in \mathbb{N}$,

$$(2.54) \quad \|W_m\|_{L^2([-\ell, 0] \times \mathcal{D})} \leq \|v\|_{L^2([-\ell, 0] \times \mathcal{D})} + \bar{\rho}\ell.$$

Also by (2.42)–(2.43) and (2.53),

$$(2.55) \quad \frac{1}{2} \|\nabla W_m\|_{L^2([-\ell, 0] \times \mathcal{D})}^2 \leq \max_{(x, y, u) \in [0, 1] \times \bar{\mathcal{D}} \times [0, 1]} |g(x, y, u)| \|W_m\|_{L^1([-\ell, 0] \times \mathcal{D})} + \ell c_1 + M.$$

Now (2.54)–(2.55) and (g_1) , (g_3) yield bounds for (W_m) in $W^{1,2}([-\ell, 0] \times \bar{\mathcal{D}})$. As in (2.55),

$$(2.56) \quad \frac{1}{2} \|\nabla W_m\|_{L^2([0, \ell] \times \mathcal{D})}^2 \leq \max_{(x, y, u) \in [0, 1] \times \bar{\mathcal{D}} \times [0, 1]} |g(x, y, u)| \|W_m\|_{L^1([0, \ell] \times \mathcal{D})} + \ell c_1 + M.$$

By elementary calculus estimates,

$$(2.57) \quad \|W_m\|_{L^1([0, \ell] \times \mathcal{D})} \leq \ell \int_{\mathcal{D}} |W_m(0, y)| dy + \ell \|\nabla W_m\|_{L^1([0, \ell] \times \mathcal{D})}$$

and similarly

$$(2.58) \quad \ell \int_{\mathcal{D}} |W_m(0, y)| dy \leq \|W_m\|_{L^1([-\ell, 0] \times \mathcal{D})} + \ell \|\nabla W_m\|_{L^1([-\ell, 0] \times \mathcal{D})}.$$

Therefore combining (2.56)–(2.58) and the bounds already obtained for (W_m) in $W^{1,2}([-\ell, 0] \times \mathcal{D})$ provides bounds for W_m in $W^{1,2}([-\ell, \ell] \times \mathcal{D})$. Since ℓ is arbitrary, W_m is bounded in $W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$. Consequently there is a $U \in W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$ such that along a subsequence, $W_m \rightarrow U$ weakly in $W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$ and strongly in $L_{\text{loc}}^2(\mathbb{R} \times \mathcal{D})$. Note that by (2.52),

$$(2.59) \quad \|P_k U - v\|_{L^2(\Omega_1)} \leq \bar{\rho}, \quad k < 0$$

and

$$(2.60) \quad \|P_0 U - v\|_{L^2(\Omega_1)} \geq \bar{\rho}.$$

(B) The asymptotic behavior of U as $x \rightarrow -\infty$.

By (2.53), for each $\ell \in \mathbb{N}$,

$$(2.61) \quad \sum_{-\ell}^{\ell} a_p(W_m) \leq M.$$

Therefore by the weak lower semicontinuity of $a_p(\cdot)$,

$$(2.62) \quad \sum_{-\ell}^{\ell} a_p(U) \leq M$$

for each $\ell \in \mathbb{N}$. Hence

$$(2.63) \quad J(U) \leq M.$$

Now (2.63) implies

$$(2.64) \quad a_p(U) \rightarrow 0 \quad \text{as} \quad |p| \rightarrow \infty.$$

It will be shown next that (2.64) leads to (B) and (C), i.e. $U \in \Gamma^-(v)$. Let ρ be positive and together with $\bar{\rho}$ satisfy

$$(2.65) \quad \rho + \bar{\rho} < \bar{\gamma}/2$$

where $\bar{\gamma}$ is as in (2.38) and $\bar{\rho}$ also satisfies (2.50). Choose p_0 so that

$$(2.66) \quad a_p(U) < \frac{\alpha(\rho)}{2}$$

for $|p| > p_0$ where $\alpha(\rho)$ is given by Proposition 2.28. Thus if $|p| > p_0$, $P_p U \in N_\rho(\mathcal{M})$. Therefore there is a $u_p \in \mathcal{M}$ such that

$$(2.67) \quad \|P_p U - u_p\|_{W^{1,2}(\Omega_1)} \leq \rho.$$

Set

$$Q_p U = U|_{[p,p+2] \times \mathcal{D}}.$$

Since by (2.66) again,

$$(2.68) \quad a_p(U) + a_{p+1}(U) < \alpha(\rho),$$

Remark 2.37 (ii) implies that

$$(2.69) \quad \|Q_p U - \mathcal{M}\|_{W^{1,2}([0,2] \times \mathcal{D})} \leq \rho,$$

\mathcal{M} being interpreted as a subset of E_2 . Therefore there is a $\bar{u}_p \in \mathcal{M}$ such that

$$(2.70) \quad \|Q_p U - \bar{u}_p\|_{W^{1,2}([0,2] \times \mathcal{D})} \leq \rho.$$

But

$$(2.71) \quad \begin{aligned} \|Q_p U - \bar{u}_p\|_{W^{1,2}([0,2] \times \mathcal{D})}^2 &= \|P_p U - \bar{u}_p\|_{W^{1,2}(\Omega_1)}^2 \\ &\quad + \|P_{p+1} U - \bar{u}_p\|_{W^{1,2}(\Omega_1)}^2 \end{aligned}$$

so by (2.67) and (2.71),

$$(2.72) \quad \begin{aligned} \|u_p - \bar{u}_p\|_{W^{1,2}(\Omega_1)} &\leq \|u_p - P_p U\|_{W^{1,2}(\Omega_1)} \\ &\quad + \|P_p U - \bar{u}_p\|_{W^{1,2}(\Omega_1)} \leq 2\rho. \end{aligned}$$

Similarly

$$(2.73) \quad \|u_{p+1} - \bar{u}_p\|_{W^{1,2}(\Omega_1)} \leq 2\rho < r.$$

Since $\bar{u}_p \in \mathcal{M}$, the definition of γ implies $u_p = u_{p+1} = \bar{u}_p$ for all $|p| > p_0$. Hence by (2.59), (2.67), and (2.65), for $p < -p_0$,

$$(2.74) \quad \begin{aligned} \|v - u_p\|_{L^2(\Omega_1)} &\leq \|v - P_p U\|_{L^2(\Omega_1)} + \|P_p U - u_p\|_{L^2(\Omega_1)} \\ &\leq \bar{\rho} + \rho < \bar{\gamma}. \end{aligned}$$

Therefore $u_p = v$ for $p < -p_0$ and

$$(2.75) \quad \|P_p U - v\|_{W^{1,2}(\Omega_1)} \leq \rho.$$

To complete (B) requires showing

$$(2.76) \quad \|P_p U - v\|_{W^{1,2}(\Omega_1)} \rightarrow 0$$

as $p \rightarrow -\infty$. But that is now immediate from (2.64) and (iii) of Proposition 2.28.

(C) The asymptotic behavior of U as $x \rightarrow \infty$.

The arguments of (B) have already established the existence of $w \in \mathcal{M}$ such that

$$(2.77) \quad \|P_p U - w\|_{W^{1,2}(\Omega_1)} \leq \rho$$

for all $p > p_0$. Hence (2.64) and (iii) of Proposition 2.28 imply $P_p U \rightarrow w$ as $p \rightarrow \infty$. It remains only to prove that $w \neq v$. A comparison argument will be employed to do so.

Suppose that $w = v$. We claim that there is a $j \geq 0$ and $\beta > 0$ such that

$$(2.78) \quad \|P_j W_m - u\|_{W^{1,2}(\Omega_1)} > \beta$$

for all $u \in \mathcal{M}$ and for all m sufficiently large. Otherwise for each $j \geq 0$, there is a sequence $k_i(j) \rightarrow \infty$ as $i \rightarrow \infty$ and $v_{k_i(j)} \in \mathcal{M}$ such that

$$(2.79) \quad \|P_j W_{k_i(j)} - v_{k_i(j)}\|_{W^{1,2}(\Omega_1)} \rightarrow 0$$

as $i \rightarrow \infty$. Since (W_m) converges weakly to U in $W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$ along a subsequence as $m \rightarrow \infty$, $(P_j W_m)$ is bounded for each j along this subsequence. Therefore only finitely many functions in \mathcal{M} are possible candidates for $v_{k_i(j)}$ in (2.79) via Remark 2.37 (i). Thus without loss of generality, $v_{k_i(j)} = v_j$ independently of i . Again invoking the weak convergence of W_m to U in $W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$ shows

$$(2.80) \quad v_j = P_j U$$

for each $j \geq 0$. By (2.77),

$$(2.81) \quad v_j = w = P_j U$$

for large j . Since $v_j \in \mathcal{M}$, for all $j \geq 0$,

$$(2.82) \quad v_j(0, y) = v_j(1, y) = v_{j+1}(0, y)$$

via (2.80) and

$$(2.83) \quad \frac{\partial v_j}{\partial x}(0, y) = 0 = \frac{\partial v_{j+1}}{\partial x}(0, y).$$

These observations and the unique continuation result used in Proposition 2.18 imply $v_j = v_{j+1}$ for all $j \geq 0$. Hence by (2.81), $v_j = w$ for all $j \geq 0$. In particular

$$(2.84) \quad v_0 = w = P_0 U.$$

But since $v = w$, (2.84) is contrary to (2.60). Thus (2.78) is valid.

Combining (2.78) with (iii) of Proposition 2.28 shows

$$(2.85) \quad a_j(W_m) \geq \alpha(\beta)$$

for all large m . Since $J(W_m) \rightarrow c$, it can be assumed that

$$(2.86) \quad J(W_m) \leq c + \frac{1}{3}\alpha(\beta)$$

for large m . These inequalities will be used to prove that $w \neq v$. Choose

$$(2.87) \quad \delta \in (0, \bar{\gamma}/2)$$

so that

$$(2.88) \quad \max_{u \in \overline{B_{3\delta}(w)}} \int_{\Omega_1} (\mathcal{L}(u) - \mathcal{L}(w)) dx dy \leq \frac{\alpha(\beta)}{3}.$$

With j given via (2.78), since $P_p U \rightarrow w$ in $W^{1,2}(\Omega_1)$ as $p \rightarrow \infty$, $\bar{p} = \bar{p}(\delta) > j$ can be chosen so that for $p \geq \bar{p}$,

$$(2.89) \quad \|P_p U - w\|_{W^{1,2}(\Omega_1)} \leq \frac{\delta}{4}.$$

We claim there is a large $m = m(\delta)$ so that for some $p = p(m) \geq \bar{p}$,

$$(2.90) \quad \|P_p W_m - w\|_{W^{1,2}(\Omega_1)} \leq \delta.$$

Indeed otherwise, for each $p \geq \bar{p}$ and all large m ,

$$(2.91) \quad \|P_p W_m - w\|_{W^{1,2}(\Omega_1)} > \delta.$$

By (2.89),

$$(2.92) \quad \|P_p W_m - w\|_{L^2(\Omega_1)} \leq \|P_p W_m - P_p U\|_{L^2(\Omega_1)} + \frac{\delta}{4}.$$

The first term on the right hand side of (2.92) can be assumed to converge to 0 as $m \rightarrow \infty$. Hence there is an $\bar{m} = \bar{m}(p, \delta)$ such that for $m \geq \bar{m}$,

$$(2.93) \quad \|P_p W_m - w\|_{L^2(\Omega_1)} \leq \frac{\delta}{2}.$$

If $u \in \mathcal{M} \setminus \{w\}$, by (2.87) and (2.93),

$$(2.94) \quad \|P_p W_m - u\|_{W^{1,2}(\Omega_1)} \geq \frac{\delta}{2}.$$

Consequently by (2.91), (2.94), and (iii) of Proposition 2.28,

$$(2.95) \quad a_p(W_m) \geq \alpha \left(\frac{\delta}{2} \right)$$

for all $m \geq \bar{m}$. Choose $\ell = \ell(\delta) \in \mathbb{N}$ so that

$$(2.96) \quad (\ell + 1)\alpha \left(\frac{\delta}{2} \right) > c + \frac{\alpha(\beta)}{3}.$$

Now (2.95) holds for each $p \geq \bar{p}(\delta)$ and $m \geq \bar{m}(p, \delta)$. Hence for $m \geq \max_{\bar{p} \leq p \leq \bar{p} + \ell} \bar{m}(p, \delta)$, by (2.95)–(2.96),

$$(2.97) \quad J(W_m) > c + \frac{\alpha(\beta)}{3},$$

contrary to (2.86). Thus (2.90) holds.

Now finally to complete the proof of (c), for m and p given by (2.90) define

$$\begin{aligned}
 (2.98) \quad U_m(x, y) &= v(x, y) \quad x \leq p, y \in \bar{\mathcal{D}} \\
 &= ((p+1) - x)v(x, y) + (x-p)W_m(x, y) \\
 &\quad p < x \leq p+1, y \in \bar{\mathcal{D}} \\
 &= W_m(x, y) \quad p+1 < x, y \in \bar{\mathcal{D}}.
 \end{aligned}$$

Then $U_m \in \Gamma^-(v)$. It will be shown that $J(U_m) < c$ which violates (2.44) and therefore $v \neq w$. Note that

$$\begin{aligned}
 (2.99) \quad \|P_p U_m - w\|_{W^{1,2}(\Omega_1)} &= \|(p-x)(W_m - w)\|_{W^{1,2}(\Omega_1)} \\
 &\leq 2\|P_p W_m - w\|_{W^{1,2}(\Omega_1)} \leq 2\delta.
 \end{aligned}$$

Hence by (2.88) and (2.99).

$$(2.100) \quad a_p(U_m) \leq \frac{\alpha(\beta)}{3}.$$

Therefore by (2.98), (2.100), (2.85)–(2.86),

$$\begin{aligned}
 (2.101) \quad J(U_m) &= a_p(U_m) + \sum_{p+1}^{\infty} a_k(W_m) \\
 &\leq a_p(U_m) + J(W_m) - a_p(W_m) \\
 &\leq \frac{\alpha(\beta)}{3} + c + \frac{\alpha(\beta)}{3} - \alpha(\beta) = c - \frac{\alpha(\beta)}{3} < c
 \end{aligned}$$

which is impossible. Thus (C) is proved.

(D) $J(U) = c$.

By (B) and (C), $U \in \Gamma^-(v)$. Therefore

$$(2.102) \quad J(U) \geq c.$$

Let $\epsilon > 0$. Then for m sufficiently large and any $\ell \in \mathbb{N}$,

$$(2.103) \quad c + \epsilon \geq J(W_m) \geq \sum_{-\ell}^{\ell} a_p(W_m).$$

Letting $m \rightarrow \infty$, (2.103) shows

$$(2.104) \quad c + \epsilon \geq \sum_{-\ell}^{\ell} a_p(U).$$

Since ℓ is arbitrary, by (2.104),

$$(2.105) \quad c + \epsilon \geq J(U).$$

Hence

$$(2.106) \quad c \geq J(U)$$

and equality must hold.

The proof of Theorem 2.46 is complete. \square

PROOF OF COROLLARY 2.47. Let $\varphi \in C^\infty(\mathbb{R} \times \mathcal{D})$ with ‘compact support’, i.e. φ vanishes for large x and near $\partial\mathcal{D}$. Let $\delta \in \mathbb{R}$. Then $U + \delta\varphi \in \Gamma^-(v)$ and $J(U + \delta\varphi)$ is a C^1 function of δ with a local minimum at $\delta = 0$. Therefore

$$(2.107) \quad J'(U)\varphi \equiv \int_{\mathbb{R} \times \mathcal{D}} (\nabla U \cdot \nabla \varphi - g(x, y, U)\varphi) dx dy$$

for all such φ , i.e. U is a weak solution of (PDE). Standard regularity arguments then show U is a classical solution of (PDE) and satisfies the boundary conditions (1.2). In fact, since $g \in C^1$, $U \in C^{2,\beta}(\mathbb{R} \times \bar{\mathcal{D}})$ for any $\beta \in (0, 1)$. Moreover since $P_k U$ and v are bounded in $C^{2,\beta}([0, 1] \times \bar{\mathcal{D}})$ and $P_k U - v \rightarrow 0$ as $k \rightarrow -\infty$ in $W^{1,2}(\Omega_1)$, by standard interpolation inequalities $P_k U \rightarrow v$ in $C^2(\Omega_1)$ as $k \rightarrow -\infty$. Similarly $P_k U \rightarrow w$ in $C^2(\Omega_1)$ as $k \rightarrow \infty$. \square

REMARK 2.108. (i) It is natural to conjecture at least for the case when $\mathcal{M} = v + \mathbb{Z}$, i.e. \mathcal{M}/\mathbb{Z} is a singleton, that $P_k U \rightarrow v + 1$ or $v - 1$ as $k \rightarrow \infty$. Indeed this has been shown for $n = 1$, the O.D.E. case, in [2]. (ii) The arguments given here extend to the case when $\partial\mathcal{D}$ is a 1-periodic function of x and even in x and in (1.2), ν is the normal to $\partial\Omega$. (iii) If (g_2) is weakened to permit g to behave differently for e.g. $x > 0$ and $x < 0$, one still gets an analogue of Theorem 2.46. See e.g. [2] for an O.D.E. case.

§3. Some related problems

In this section, it will be shown how the results of §2 carry over to some other situations. Since the ideas are so close to those of §2, the exposition will be brief. The first variant of §2 involves (PDE) on a semi-infinite domain. To describe it, consider (PDE) for e.g. $x \leq 0$ with the boundary conditions:

$$(3.1) \quad u(0, y) = \psi(y), \quad y \in \mathcal{D}$$

and

$$(3.2) \quad \frac{\partial u}{\partial \nu}(x, y) = 0 \quad x \leq 0, y \in \partial \mathcal{D}$$

where e.g. ψ is smooth and

$$(3.3) \quad \frac{\partial \psi}{\partial \nu}(y) = 0, \quad y \in \partial \mathcal{D}$$

so (3.1) and (3.2) are compatible.

Let

$$(3.4) \quad \Gamma^- = \{u \in W_{\text{loc}}^{1,2}(\mathbb{R}^- \times \mathcal{D}) \mid u|_{x=0} = \psi \quad \text{and} \\ \|P_k u - \mathcal{M}\|_{W^{1,2}(\Omega_1)} \rightarrow 0 \quad \text{as} \quad k \rightarrow -\infty\}.$$

For $u \in \Gamma^-$ set

$$(3.5) \quad J(u) = \sum_{-\infty}^0 a_p(u)$$

where the a_p are as in (2.41) with any $v \in \mathcal{M}$. Let

$$(3.6) \quad c = \inf_{u \in \Gamma^-} J(u).$$

The analogue of Theorem 2.46 here is:

THEOREM 3.7. *Let (g_1) – (g_3) and (\mathcal{M}) be satisfied. Let ψ be smooth and satisfy (3.3). Then there is a $U \in \Gamma^-$ such that $J(U) = c$. Moreover U is a classical solution of (PDE), (3.1)–(3.2) and for some $v \in \mathcal{M}$,*

$$\|P_k U - v\|_{W^{1,2}(\Omega_1)} \rightarrow 0 \quad \text{as} \quad k \rightarrow -\infty.$$

PROOF. The argument follows the same lines as the proof of Theorem 2.46. If (W_m) is a minimizing sequence for (3.6), the estimates (2.55) and

$$(3.8) \quad \|W_m\|_{L^1([-\ell,0] \times \mathcal{D})} \leq \ell(\|\psi\|_{L^1(\mathcal{D})} + \|\nabla W_m\|_{L^1([-\ell,0] \times \mathcal{D})})$$

lead to $W_{\text{loc}}^{1,2}(\mathbb{R}^- \times \mathcal{D})$ bounds for (W_m) . Continuing to argue essentially as in (A), (C), (D) and Corollary 2.47 yields Theorem 3.7.

To prepare for the next two applications, we reexamine what was done in §2 a bit more abstractly. Thus consider (PDE) on $\mathbb{R} \times \mathcal{D}$ under either the boundary conditions (1.1) or (1.2). Assume that g satisfies (g_1) and a Sobolev growth condition:

$$(g_4) \text{ there are constants } a_1, a_2 \geq 0 \text{ and } s \in [1, \frac{n+2}{n-2}) \text{ such that } |g(x, y, z)| \leq a_1 + a_2|z|^s.$$

If $n = 2$, (g_4) can be weakened.

Let I_1 be as in (2.1) and let E_1 denote the class of $W^{1,2}$ functions on Ω_1 which are 1-periodic in x and also satisfy (1.1) if that boundary condition is operative. By (g_1) and (g_4) , I_1 is bounded on bounded sets and is weakly lower semicontinuous. Define c_1 via (2.3). Assume: (a_1) any minimizing sequence for c_1 is bounded in E_1 . Note that this is *not* true for the setting of §2 unless (g_3) is employed to divide out the \mathbb{Z} symmetry. Given (a_1) , the arguments of §2 show \mathcal{M} defined in (2.11) is nonempty. If (g_2) is also satisfied, then the conclusions of Propositions 2.13, 2.18, and 2.21 are also valid. The same is true of Proposition 2.27 if (a_k) , the analogue of (a_1) for $k \in \mathbb{N}$, holds. Further assuming (\mathcal{M}) , Proposition 2.28 obtains. Suppose: (b) the cardinality of \mathcal{M} is at least 2. Then by Remark 2.37 (i), \mathcal{M} is finite.

For any $v \in \mathcal{M}$, the set $\Gamma^-(v)$ can be introduced as in (2.40), building in the boundary conditions (2.1) if necessary. Define J and c as in (2.43)–(2.44). Then an examination of the proofs of Theorem 2.46 and Corollary 2.47 show they carry over if: (c) a minimizing sequence for c is bounded in $W_{\text{loc}}^{1,2}(\mathbb{R} \times \mathcal{D})$.

To recapitulate, given (g_1) , (g_2) and (g_4) there is an analogue of Theorem 2.46 and Corollary 2.47 provided that (a_k) , (b) (c) and (\mathcal{M}) are satisfied. Next these conditions will be verified for two examples. For the first, suppose that we are dealing with (1.2) and g satisfies

(g_5) $G(x, y, z) \leq 0$ and $= 0$ if and only if z belongs to a finite set \mathcal{F} of cardinality at least two.

(g_6) There are constants $\beta > 0$, $R \geq 0$ such that $G(x, y, z) \geq \beta|z|^2$ for $|z| \geq R$.

Note that by (g_5) and (g_1) , G has local maxima for $x \in \mathbb{R}$, $y \in \bar{\mathcal{D}}$, and $z \in \mathcal{F}$. Therefore $g(x, y, z) = 0$ at such points. \square

PROPOSITION 3.9. *If g satisfies (g_1) – (g_2) , (g_4) – (g_6) , then (a_k) , (b), (c), and (\mathcal{M}) are satisfied.*

PROOF. By (g_1) and (g_5) , $\mathcal{M} = \mathcal{F}$ and $c_1 = 0$. Therefore (b) and (\mathcal{M}) hold. To verify (a_1) (and similarly (a_k)), note that if (u_m) is a minimizing sequence for c_1 , there is a $K > 0$ such that

$$(3.10) \quad \int_{\Omega_1} \left(\frac{1}{2} |\nabla u_m|^2 - G(x, y, u_m) \right) dx dy \leq K$$

so by (g_5) ,

$$(3.11) \quad \|\nabla u_m\|_{L^2(\Omega_1)}^2 \leq 2K.$$

Moreover

$$(3.12) \quad \begin{aligned} \int_{\Omega_1} u_m^2 dx dy &= \int_{\{(x,y) \mid |u_m(x,y)| \leq R\}} u_m^2 dx dy \\ &\quad + \int_{\{(x,y) \mid |u_m(x,y)| > R\}} u_m^2 dx dy \\ &\leq |\Omega_1| R^2 + \beta^{-1} K \end{aligned}$$

via (g_6) and (3.10). Thus (u_m) is bounded in $W^{1,2}(\Omega_1)$. Likewise to verify (c), if (u_m) is a minimizing sequence for c , for any $\ell \in \mathbb{N}$,

$$(3.13) \quad \int_{-\ell}^{\ell} \int_{\mathcal{D}} \left(\frac{1}{2} |\nabla u_m|^2 - G(x, y, u_m) \right) dx dy \leq M + \ell c,$$

and as in (3.10)–(3.12), this leads to a bound for (u_m) in $W_{loc}^{1,2}(\mathbb{R} \times \mathcal{D})$.

For the final application of these ideas, consider (PDE) under (1.1). Then there is a constant $\alpha_1 > 0$ such that

$$(3.14) \quad \|u\|_{L^2(\Omega_1)} \leq \alpha_1 \|\nabla u\|_{L^2(\Omega_1)}$$

for all $u \in E_1$. Suppose that g satisfies (g_1) – (g_2) and

(g_7) There is an $\bar{M} > 0$ such that

$$|g(x, y, z)| \leq \bar{M} \quad \text{for all} \quad (x, y, z) \in \mathbb{R} \times \bar{\mathcal{D}} \times \mathbb{R}.$$

Then a fortiori (g_4) is verified and by (2.7) and (3.14), (a_1) and similarly (a_k) and (c) hold. If also

(g_8) $g(x, y, 0) = 0$

and

(g_9) there is a $\varphi \in E$, such that $I_1(\varphi) < 0$,

then $I_1(0) = 0 > c_1$. Suppose also that

(g_{10}) $g(x, y, z) = -g(x, y, -z)$.

Then $v \in \mathcal{M}$ implies $-v \in \mathcal{M}$ and $-v \neq v$. Thus (b) is satisfied. In general, (\mathcal{M}) may not hold but a modification in the spirit of earlier remarks can be made to replace g by a new function for which (\mathcal{M}) obtains.

Condition (g_9) can be verified for problems of the type considered by Kirchgässner [6] if $\lambda > \lambda_1$. More precisely consider (1.4) together with (1.5) or its higher dimensional analogue (1.1). It is assumed that $a(y) > 0$ on $\bar{\mathcal{D}}$ and is in C^1 . The function f is also assumed to be in C^1 and $f(x, y, z, \lambda) = o(|z|)$ as $|z| \rightarrow 0$ uniformly on bounded λ intervals. Suppose $\lambda > \lambda_1$, the smallest eigenvalue of (1.6) (or its higher dimensional variant). Then it is not difficult to show—see e.g. [10]—that

$$(3.15) \quad I_1(u) = \int_{\Omega_1} \left(\frac{1}{2} |\nabla u|^2 - \frac{\lambda}{2} au^2 \right) dx dy + o(\|u\|_{W^{1,2}(\Omega_1)}^2) \quad \text{as} \quad u \rightarrow 0.$$

In particular for $u = \psi$, a small multiple of the eigenfunction of (1.6) corresponding to λ_1 ,

$$(3.16) \quad I_1(u) = \frac{(\lambda_1 - \lambda)}{2} \int_{\Omega_1} |\nabla u|^2 dx dy + o(\|\nabla u\|_{L^2(\Omega_1)}^2) < 0$$

and (g_9) is verified.

As an example of a class of functions for which (g_1) – (g_2) , (g_7) – (g_{10}) all are satisfied, consider $g(x, y, z) = \lambda a(y)z - h(x, y, z, \lambda)z$ where $\lambda > \lambda_1$, a is as earlier, and h is C^1 in its arguments, even and 1-periodic in x , $h(x, y, 0) = 0$, $h(x, y, -z) = h(x, y, z)$, and there is an $\hat{R} > 0$ such that

$$(3.17) \quad h(x, y, \hat{R}) > \lambda a(y).$$

It is easy to check that (g_1) – (g_2) and (g_8) – (g_{10}) obtain but a priori, (g_7) is not satisfied. Set

$$(3.18) \quad \begin{aligned} \hat{g}(x, y, z) &= g(x, y, z) & |z| \leq \hat{R} \\ &= g(x, y, \hat{R}) & z > \hat{R} \\ &= -g(x, y, -\hat{R}) & z < -\hat{R}. \end{aligned}$$

Then \hat{g} satisfies (g_1) – (g_2) and (g_7) – (g_{10}) . (Actually \hat{g} is Lipschitz continuous rather than C^1 but that is sufficient for the earlier arguments to work.) Therefore there is a solution of heteroclinic type for

$$(3.19) \quad -\Delta u = \hat{g}(x, y, w) \quad x \in \mathbb{R}, y \in \mathcal{D}$$

and (1.1). If this solution satisfies

$$(3.20) \quad \|u\|_{L^\infty(\mathbb{R} \times \mathcal{D})} \leq \hat{R}$$

then u is also a solution of (PDE) via (3.18).

To verify (3.20), let $\hat{\mathcal{M}}$ denote the set of minimizing periodic solutions for (3.19) and (1.1). Then $\hat{\mathcal{M}} \subset \mathcal{M}$. Indeed if $v \in \hat{\mathcal{M}}$, since $v \not\equiv 0$, it has either a positive maximum or negative minimum in Ω_1 , e.g. at (\hat{x}, \hat{y}) . Assuming the former case, if $v(\hat{x}, \hat{y}) \geq \hat{R}$,

$$(3.21) \quad 0 \leq -\Delta v(\hat{x}, \hat{y}) = \hat{g}(\hat{x}, \hat{y}, v) < 0$$

by (3.17)–(3.18). Hence $\hat{R} > v$ in Ω_1 and similarly $v > -\hat{R}$, i.e.

$$(3.22) \quad \|v\|_{L^\infty(\Omega_1)} < \hat{R}.$$

Therefore v satisfies (PDE) and $v \in \mathcal{M}$. Finally if u is a solution of (3.19), since $u \rightarrow \hat{\mathcal{M}}$ as $|x| \rightarrow \infty$, $|u(x, y)| < \hat{R}$ for large $|x|$ via (3.22). If u does not satisfy (3.20), it has a positive maximum or a negative minimum at some $(\hat{x}, \hat{y}) \in \mathbb{R} \times \mathcal{D}$. Then the argument of (3.21) again leads to (3.20) so u is a solution of (PDE).

References

- [1] Rabinowitz, P. H., Heteroclinics for a reversible Hamiltonian system, to appear in *Ergodic Th. and Dynamical Sys.*
- [2] Rabinowitz, P. H., Heteroclinics for a reversible Hamiltonian system, 2, to appear in *Differential and Integral Equations.*
- [3] Bolotin, S. V., The existence of homoclinic motions, *Vestnik Moskovskogo Universiteta, Matematika* **38** (1983), 98–103.
- [4] Bolotin, S. V., Homoclinic orbits to invariant tori of symplectic diffeomorphisms and Hamiltonian systems, preprint.
- [5] Moser, J., Minimal solutions of variational problems on a torus, *Ann. I.H.P. Analyse nonlinéaire* **3** (1986), 229–272.
- [6] Kirchgässner, K., Wave-solutions of reversible systems and applications, *J. Diff. Eq.* **45** (1982), 113–127.
- [7] Turner, R. E. L., Internal waves in fluids with rapidly varying density, *Ann. Sc. Norm. Sup. Pisa, Classe di Scienze, Ser. 4* **8** (1981), 513–573.
- [8] Kirchgässner, K., Nonlinearly resonant surface waves and homoclinic bifurcation, *Adv. Appl. Mech.* **26** (1988), 135–181.
- [9] Mielke, A., Reduction of quasilinear elliptic equations in cylindrical domains with applications, *Math. Mech. App. Sc.* **10** (1988), 51–66.
- [10] Rabinowitz, P. H., *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Reg. Conf. Ser. in Math., No. 65, AMS 1986.
- [11] Nirenberg, L., *Lectures on Linear Partial Differential Equations*, CBMS Reg. Conf. Ser. in Math., No. 17, AMS 1973.

(Received April 20, 1994)

Department of Mathematics
University of Wisconsin-Madison
480 Lincoln Drive
Madison, WI 53706-1388
U.S.A.