

*Topological entropy and topologically
mixing property in symbolic dynamics*

By Li SIMIN

Abstract. In this paper we construct an example to show usually positive topological entropy does not imply mixing property at all in symbolic dynamics.

In ergodic theory, it is well-known that a K -systems, i.e., a measure preserving transformation having positive metric entropy with respect to every nontrivial finite measurable partition, is strongly mixing. But F.Blandchard proved the topological setting of above is not true. He constructed a subshift which has completely positive topological entropy but not topologically mixing. It is not too difficult to check that for a subshift of finite type, it is topologically mixing if and only if it has completely positive topological entropy.

On the other hand, it is well-known [2] that if a continuous map $f : X \rightarrow X$ from an interval or a circle to itself has positive entropy, then some subsystems are topologically mixing, i.e., there exists a closed subset $\Lambda \subset X$ and an integer $m > 0$ such that $f^m(\Lambda) \subset \Lambda$ and $f^m|_{\Lambda}$ is topologically mixing. A. Katok [3] also got analogous results for $C^{1+\alpha}$ ($\alpha > 0$) diffeomorphism of a two dimensional compact manifold.

In zero dimension case, it is easy to check it is also true for a subshift of finite type. However we will construct a subshift which has positive entropy but has no mixing subsystem.

In the construction we use the Denjoy C^1 diffeomorphism of the circle. We will state the construction of the Denjoy diffeomorphism briefly without proof. For the detail see for example [4].

1991 *Mathematics Subject Classification.* 58F20, 54H20.

This research is supported in part by Sumitomo Bank Fellowship.

Consider a countable set $\{I_m = [0, l_m] : m \in \mathbb{Z}, l_m > 0\}$ of closed interval which satisfies $\sum_{m \in \mathbb{Z}} l_m = l < \infty$. Let α be an irrational number and define $\alpha_m, m \in \mathbb{Z}$, by $\alpha_m = m\alpha \pmod{1}$, $0 \leq \alpha_m < 1$. We construct an interval of length of $1+l$ by inserting I_m at $\alpha_m, m \in \mathbb{Z}$, in the interval $[0,1]$. We obtain a circle by identifying the point 0 and 1, and denote it by S^1 . Let α_m and β_m denote the left endpoint and the right endpoint of I_m , respectively.

Construct a C^1 map $f_m : I_m \rightarrow I_{m+1}, f_m(0) = 0, f_m(l_m) = l_{m+1}$ and some other condition (See [4]). Define a map $f : S^1 \rightarrow S^1$ by

$$f(x) = \begin{cases} f_m(x) \in I_{m+1} \subset S^1 & x \in I_m \\ x + \alpha \pmod{1} \in [0, 1] \subset S^1 & x \in [0, 1] \end{cases}$$

It is easily seen f is well defined and it can be made to be a C^1 diffeomorphism. It is called a Denjoy C^1 diffeomorphism of the circle.

For $m \in \mathbb{Z}$, define a half open half closed interval $J_m \subset S^1$ by

$$J_m = \begin{cases} [\beta_{m-1}, \beta_m) \subset S^1 & \beta_{m-1} < \beta_m \\ [\beta_{m-1}, 1] \cup [0, \beta_m) \subset S^1 & \beta_{m-1} > \beta_m \end{cases}$$

Now define a map $\phi : S^1 \rightarrow \Sigma_2 = \{0, 1\}^{\mathbb{Z}}$ by

$$\phi(x)(m) = \begin{cases} 0 & x \notin J_m \\ 1 & x \in J_m \end{cases}$$

for $m \in \mathbb{Z}$ and $x \in S^1$. Give S^1 the usual topology with metric d . Let $X = S^1 - \bigcup_{m \in \mathbb{Z}} \text{Int}(I_m) = [0, 1] \cup \{\beta_m : m \in \mathbb{Z}\}$. X is a closed subset of S^1 and $f(X) = X$. we will show :

- (1) $\phi|_X : X \rightarrow \Sigma_2$ is injective,
- (2) $\phi|_X : X \rightarrow \Sigma_2$ is continuous.

Proof of (1). For $x, y \in [0, 1], x < y$, since $\{\alpha_m : m \in \mathbb{Z}\}$ is dense in $[0, 1]$, there exists an integer $m \in \mathbb{Z}$ and $x \leq \alpha_m < y$. Let $n = \inf\{k > m : a_k \geq y \text{ or } a_k < x\}$. Then $\phi(y)(n) = 1$ but $\phi(x)(n) = 0$. So $\phi(x) \neq \phi(y)$. Similarly $\phi(x) \neq \phi(\beta_m)$ for any $x \in [0, 1]$ and any $m \in \mathbb{Z}$. consequently $\phi|_X$ is injective.

Proof of (2). For any $k > 0, k \in \mathbb{Z}$, let $\delta = \min\{l_m\} - (k + 1) \leq m \leq (k + 1)$. Let $x, y \in X$ and $d(x, y) < \delta$. If for some $m, -k \leq m \leq k, x \in J_m,$

but $y \notin m$, then $\beta_{m-1} \leq x \leq \beta_m$, but $y > \beta_m$ or $y < \alpha_{m-1}$. So $d(x, y) > l_m$ or $d(x, y) > l_{m-1}$. This contradicts with $d(x, y) < \delta$.

Since $\phi|_X$ is an injective continuous map of a compact space to a Hausdorff space, $\phi(X) \subset \Sigma_2$ is closed and $\phi|_X : X \rightarrow \Lambda = \phi(X) \subset \Sigma_2$ is a homeomorphism. Obviously $\phi \circ f(x) = \sigma \circ \phi(x)$. We get a subshift $\sigma : \Lambda \rightarrow \Lambda$ which is conjugate to $f : X \rightarrow X$. Since f acts in X as an irrational rotation. $f|_X$ can not have mixing subsystems. Neither can $\sigma : \Lambda \rightarrow \Lambda$.

Let $\Sigma'_2 = \{a, b\}^{\mathbb{Z}}$ and $\Sigma_4 = \{0a, 0b, 1a, 1b\}^{\mathbb{Z}}$. Define a map $\psi : \Lambda \times \Sigma'_2 \rightarrow \Sigma_4$ by

$$\psi(x, y)(m) = x(m)y(m) \text{ for } m \in \mathbb{Z}, x \in \Lambda, y \in \Sigma'_2$$

Obviously ψ is injective and continuous. Let $W = \psi(\Lambda \times \Sigma'_2) \subset \Sigma_4$. $\sigma \times \sigma : \Lambda \times \Sigma'_2 \rightarrow \Lambda \times \Sigma'_2$ is conjugate to $\sigma : W \rightarrow W$.

Since $\sigma : \Lambda \rightarrow \Lambda$ can not have mixing subsystems, neither can $\sigma \times \sigma : \Lambda \times \Sigma'_2 \rightarrow \Lambda \times \Sigma'_2$. Consequently $\sigma : W \rightarrow W$ does not have mixing subsystems. But $h(\sigma|_W) = h(\sigma \times \sigma|_{\Lambda \times \Sigma'_2}) \geq h(\sigma|_{\Sigma_2}) = \log 2 > 0$

Acknowledgement. The author would like to thank Professor T. Tsuboi for his constant encouragement and instruction.

References

- [1] Blanchard, F., Fully positive topological entropy and topologically mixing, Contemporary Mathematics 135, symbolic dynamics and its application, pp. 95–106.
- [2] Block, L. S. and W. A. Coppel, Dynamics in one dimension, Lecture Notes in Mathematics 1513, Springer-Verlag, 1992.
- [3] Katok, A., Lyapunov exponent, entropy and periodic orbits for diffeomorphism, Publ. Math. IHES 51 (1980), 137–174.
- [4] Tamura, I., Topology of Foliation: An Introduction, 1992.
- [5] Weiss, Subshift of finite type and sofic systems, Monatsh. Math. **77** (1973), 462–474.
- [6] Xiong, J., Chaos caused by a topologically mixing map, Advanced Series in Dynamic Systems (K. Shiraiwa, ed.), vol. 9, pp. 550–572.

(Received November 10, 1994)

Department of Mathematical Sciences
University of Tokyo

Hongo, Tokyo 113
JAPAN