

Boundedness of Fano Threefolds with Log-Terminal Singularities of Given Index

By Alexandr BORISOV

Abstract. We prove that all Fano threefolds with log-terminal singularities of given index belong to finitely many families. This result was previously obtained by the author in the case of unipolar Fano varieties.

1. Introduction

This paper is concerned with Fano varieties with log-terminal singularities. Fano varieties and log-terminal singularities play very important role in modern birational algebraic geometry (cf. [8], [9], [12]).

The main goal of this paper is to prove the following theorem.

THEOREM 1.1. *For any $n \in \mathbb{N}$ the family of Fano threefolds with log-terminal singularities of index n is bounded.*

This theorem is the three-dimensional case of the following conjecture originally proposed by V. Batyrev. It was previously obtained by the author in [4] under the additional assumptions that X is unipolar, i.e. it is \mathbb{Q} -factorial and the Picard number $\rho(X)$ is equal to 1.

CONJECTURE 1.1 (Batyrev). *For any two fixed natural numbers k, n there are just finitely many families of Fano varieties of dimension k with log-terminal singularities of index n .*

Theorem 1.1. has several noteworthy corollaries.

COROLLARY 1.1. *The family of Fano threefolds with rational Gorenstein singularities is bounded.*

1991 *Mathematics Subject Classification.* 14J45.

COROLLARY 1.2. *Suppose X is a Fano threefold with log-terminal singularities. Then the algebraic fundamental group of its smooth locus is finite.*

Also, combined with the recent work of Kollár, Miyaoka, Mori and Takagi it implies boundedness of Fano threefolds with canonical singularities (cf. [18]).

Through the above application and by itself, Theorem 1.1 is a step towards proving the following stronger conjecture, proposed independently by the author (cf. [4]) and V. Alexeev (cf. [2]).

CONJECTURE 1.2 (Borisov-Alexeev). *For any fixed $n \in \mathbb{N}$, $\varepsilon > 0$ there are just finitely many families of n -dimensional Fano varieties with ε -log-terminal singularities.*

This conjecture is important for the so-called log Sarkisov Program (cf. [6]). Some partial results towards it are due to Alexeev and Nikulin (in dimension 2), the author and L. Borisov (in the toric case), and Y. Kawamata (cf. [10]). The boundedness of smooth Fano varieties in any given dimension was proven by Kollár, Miyaoka, and Mori (cf. [16]). Their method (the “rational curve surgery”) was the basis of the method of [4]. Recently, Kollár, Miyaoka, Mori and Takagi proved boundedness of three-dimensional Fano varieties with canonical singularities (using the result of this paper). Also recently, an entirely new approach to the boundedness of Fano varieties was proposed by Ziv Ran (cf. [22], [23]). It is based on studying sheaves of high order differential operators on the plurianticanonical bundles, and has some potential to compete with the rational curve surgery.

Also, H. Tsuji recently announced the proof of Batyrev conjecture in any dimension, under extra assumptions that X is unipolar (cf. [25]). His methods are analytic, and are yet to be independently verified.

In this paper we employ the techniques of [4] together with some new ideas. These new ideas include the following.

1. Using Alexeev Minimal Model program with suitable boundary to find horizontal extremal contractions.
2. Kollár’s effective Base Point Freeness theorem.

3. Using Kawamata's result on the length of extremal curves with suitable boundary to avoid gluing curves in some cases.

The paper is organized as follows. In section 2 we consider the problem in dimension 2. Of course, Batyrev conjecture in dimension 2 was first proven by Nikulin in [21]. But the rational curve surgery provides a very easy proof. Also in section 2 we discuss some reformulations of the problem that follow from Kollár's Effective Base Point Freeness theorem.

In section 3 we prove that, except for a bounded family, all Fano threefolds with log-terminal singularities of given index n contain a covering family of rational curves $\{l\}$ such that $l \cdot (-K_X)$ is bounded in terms of n , and the rationally connected fibration associated to $\{l\}$ has fibers of dimension 2.

In section 4 we complete the proof of Batyrev conjecture in dimension three.

Notations. We utilize the notations of [4]. That is, we will usually identify curves on different birationally equivalent varieties if they coincide in their general points. The identified curves will be usually denoted by the same symbol. The same convention will be used for the two-dimensional subvarieties. If it is necessary to point out that, say, a prime divisor S is considered on a variety X it will be denoted by S_X . Another convention is that $\{l\}$ will denote the family of curves with a general element l and $\{H\}$ will denote the LINEAR system of Weil divisors with a general element H . Whenever we have a family of curves, a general element is reduced and irreducible unless the opposite is explicitly specified. It will be clear in every particular case why these conventions agree with each other.

Additionally, by a constant c or $c(n)$ we will mean some positive constant that only depends on n . The value of c can be different in different parts of the proof.

Acknowledgments. The author is indebted to V. Iskovskikh who introduced him to algebraic geometry and Fano varieties in particular. The author would like to thank V. Alexeev, V. Batyrev, Y. Kawamata, K. Matsuki, S. Mori, M. Reid, and V. Shokurov for their interest in author's birational geometry research over the years. The author is thankful to J. Kollár, J. McKernan and the referee for helpful advice. This research was conducted

while the author was working at Washington University, St. Louis. The author would like to thank the algebraic geometers there (N. M. Kumar, V. Masek, D. Wright) for their support and hospitality.

2. Two-Dimensional Case and Some Easy Equivalences

We begin with the following theorem, which is the two-dimensional case of the conjecture of Batyrev, mentioned above.

THEOREM 2.1. *For any $n \in \mathbb{N}$ the family of log del Pezzo surfaces of index n is bounded.*

PROOF. Suppose X is a log del Pezzo surface of index n . By the result of Miyaoka and Mori (cf. [19]), there exists a covering family $\{l\}$ of rational curves on X such that $l \cdot (-K_X) \leq 4$. If a general l passes through some singularity of X then by Lemma 2.2 of [4], $(-K_X)^2$ is bounded. This is because in dimension 2 for any log-terminal singularity $x_0 \in X$ $mult(x_0) \leq 2 \cdot index(x_0)$.

If a general l is contained in $X \setminus Sing(X)$, consider the minimal resolution (or terminal modification) $\pi : Y \rightarrow X$. There are two cases.

1) Two general points of Y can be connected by a chain of at most two curves from $\{l\}$.

2) Two general curves from $\{l\}$ do not intersect each other. In this case $l^2 = 0$, $-K_X \cdot l = -K_Y \cdot l = 2$.

In the first case $(-K_X)^2 \leq 2!(2 \cdot 4)^2 = 128$. (cf., e.g., [17]).

In the second case, because all log del Pezzo surfaces are rational, there exists a rational curve L on Y such that $L \cdot l > 0$. Applying Mori's bend-and-break procedure to it, we can find a new rational curve L' , such that $L' \cdot l > 0$ and L' does not admit a non-trivial deformation with two fixed points. This means, in particular, that $L' \cdot (-K_Y) \leq 3$. Also, $L' \cdot K_Y \leq L' \cdot K_X$. So $L' \cdot K_Y \leq -1$. By one of the gluing lemmas of Kollár-Miyaoka-Mori (cf. [16]) we can glue together L' and at most 2 copies of l to obtain a new family of rational curves l' , which connects two general points of Y . Therefore $(-K_X)^2 \leq (3 + 2 \cdot 2)^2 = 49$.

So in all cases $(-K_X)^2$ is bounded. The family of such log del Pezzo surfaces is bounded by the results of Kollár (cf. [13]). \square

REMARK. Instead of applying the gluing lemma above, one can use a simple Riemann-Roch argument. We chose the above proof only for its similarity with our proof of the three-dimensional case.

The following theorem is a corollary of the results of Kollár.

THEOREM 2.2. *Suppose X is a Fano variety of dimension k with log-terminal singularities of index n . Then we have the following.*

1. *For some natural number N that only depends on k , the Cartier divisor $-N \cdot n \cdot K_X$ is very ample. For $k = 3$ one can choose $N = 4320$.*
2. *For every k the following two statements are equivalent.*
 - a) *The family of such varieties is bounded.*
 - b) *The “degree” $(-K_X)^k$ is bounded.*

PROOF. The first assertion is a direct corollary of the Effective Base Point Freeness theorem of Kollár (cf. [13], 1.1, 1.2). For the second assertion we only need to prove that (b) implies (a). If (b) is satisfied then by the result of Kollár and Matsusaka (cf. [15]) all coefficients of the Hilbert polynomial of $(-nK_X)$ are bounded. Together with the first assertion this implies the result. \square

The following theorem shows that instead of bounding $(-K_X)^k$ one can also bound the dimension of the space of global sections of big enough multiple of $(-nK_X)$.

THEOREM 2.3. *Suppose X , n , k are as above. Then there is a constant A that only depends on k with the following property.*

For all $l \geq 2k$, $(-nK_X)^k \leq A \cdot h^0(l \cdot (-nK_X))$

PROOF. By the Kawamata-Viehweg vanishing theorem (cf. [12]), for all $i \geq 0$ $h^0(-inK_X) = \chi(-inK_X)$.

So $f(i) = h^0(-inK_X)$ is a polynomial of degree k for $i \geq 0$. It obviously has the following five properties.

1. $f(0) = 1$
2. $f(l) = h^0(l \cdot (-nK_X))$

3. $f(i) \geq 0$ for all $i \geq 0$
4. If for some $i \leq l$, $f(l - i) > 0$, then $f(i) \leq f(l)$
5. The highest degree term of f is $\frac{(-nK_X)^k}{k!}x^k$

Because f is not identically zero, it has at most k integer roots. So for any fixed $l \geq 2k$ one can find a sequence of k integers $1 \leq i_1 < i_2 < \dots < i_k \leq 2k$ such that for all j , $f(l - i_j) > 0$. By the property (4) this implies that $f(i_j) \leq f(l)$. Also $f(0) = 1 \leq f(l)$, because otherwise f would vanish at points i_j and l . The only way it could happen without getting $(k + 1)$ zeroes is if $i_k = 2k = l$. But in this case $f(k) = 0$ and f vanishes for at least half of the $(2k - 2)$ numbers $1, 2, \dots, (k - 1), (k + 1), \dots, (2k - 2), (2k - 1)$. So we get $(k + 1)$ zeroes anyway.

There are just finitely many possibilities for the sequence $\{i_j\}$. For each of them, $f(x)$ is determined by its values at 0 and i_j , $j = 1, 2, \dots, k$. And we can use the above inequalities and Lagrange interpolation formula at the above points to estimate the highest coefficient of $f(x)$ above by some multiple of $f(l)$. Combining these estimates with the property (5) proves the theorem. \square

REMARK. The constant $2k$ in the above theorem can probably be improved. See, in particular, [4], Lemma 2.1. It would be more important, however, to drop the dependence on n . The problem is, of course, that this would require some Riemann-Roch theorem for Weil divisors, and $h^0(-iK_X)$ is in general not a polynomial for $i \geq 0$.

By the above theorems, to prove the Batyrev conjecture in dimension 3 it is enough to obtain a bound on the self-intersection or the dimension of the space of global sections of the very ample divisor $H = -4320nK_X$. This is going to be our goal. Many of our constructions will be carried out under the assumption that $h^0(H)$ or H^3 is sufficiently large.

3. Enlarging the Miyaoka-Mori Family

In the remainder of the paper we will use the notion of rationally connected fibration associated to a covering family of rational curves. We refer to [7] for the construction (cf. also [16]).

THEOREM 3.1. *Suppose X is a 3-dimensional Fano variety with log-terminal singularities of fixed index n . Then for some constant c either $(-K_X)^3 \leq c$ or there exists a covering family of rational curves $\{l\}$ on X with the following properties.*

- 1) *The degree $l \cdot H \leq c$*
- 2) *The rationally connected fibration associated to $\{l\}$ has fibers of dimension 2.*

PROOF. Suppose X is as above. By Miyaoka-Mori theorem (cf. [19]) there is a covering family $\{l\}$ of rational curves on X such that $l \cdot (-K_X) \leq 6$.

Consider the associated rationally connected fibration. If its image is a point, $(-K_X)^3$ is bounded (cf., e.g. [4]). If its image is a curve then we are done. So we just need to consider the case when the associated rationally connected fibration has image of dimension 2. In this case the family $\{l\}$ is the family of fibers. If a general l passes through the singularities of X , one can use the methods of [4], sections 5,6, to construct a new family whose rationally connected fibration has fibers of dimension at least 2, and degree is bounded in terms of n . So we can assume that $\{l\}$ doesn't pass through $Sing(X)$.

In this case $\{l\}$ is a free family on $X \setminus Sing(X)$ in the terminology of Nadel ([20]). Consider the associated rationally connected fibration on the terminal modification $\pi : Y \rightarrow X$. That is, we have a Zariski open subset U of Y and a proper morphism $\varphi : U \rightarrow Z$. Take a "general" curve C on Z . Consider the divisor D which is a Zariski closure of $\varphi^{-1}(C)$. By the definition of Y , D is \mathbb{Q} -Cartier, $k \cdot D$ is Cartier for some $k \in \mathbb{N}$. Consider an ample \mathbb{Q} -Cartier divisor M on Y which is obtained by subtracting some exceptional divisors of π from $\pi^*(-K_X)$.

By construction $D \cdot l = 0$ and $(K_X + M) \cdot l = 0$. For big enough and divisible enough m , $mM - kD$ is very ample. Let's denote it by F . Then one can apply the Alexeev Minimal Model Program (cf. [1]) to $(Y, K_Y + \frac{1}{m}|F|)$.

Let us denote the end result of this program by Y_1 . From the construction there exists a one-dimensional Zariski closed subset S of Y_1 such that the restriction of $(\pi_{Y_1}^Y)^{-1}$ to $U = Y_1 \setminus S$ is an isomorphism.

We have several possibilities for Y_1 . Suppose first that Y_1 is a minimal model. Then $K_{Y_1} + \frac{1}{m}|F|_{Y_1}$ is numerically effective on Y_1 . By intersecting two general hyperplane sections we can choose a covering family of (not necessarily rational) curves $\{C\}$ such that $C \subset U$ and $C_Y \cdot D > 0$. Then we

get a contradiction as follows.

$$0 \leq (K_{Y_1} + \frac{1}{m}|F|_{Y_1}) \cdot C = (K_Y + \frac{1}{m}|F|) \cdot C_Y \leq -\frac{1}{m}(kD) \cdot C_Y < 0$$

So the Y_1 is not a minimal model, and thus we have a Mori fibration $Y_1 \rightarrow Z$.

If $\dim Z = 0$ then we can immediately bound $(-K_X)^3$. Indeed, for any positive divisible enough p we have the following inequalities.

$$h^0(-pK_X) \leq h^0(-pK_Y) \leq h^0(-pK_{Y_1})$$

By the Kawamata-Vieweg vanishing and Riemann-Roch, $h^0(-pK_{Y_1})$ grows like $\frac{1}{3!}(-K_{Y_1})^3 \cdot p^3$ and $h^0(-pK_X)$ grows like $\frac{1}{3!}(-K_X)^3 \cdot p^3$. By the result of Kawamata [10] (cf. also [18]) $(-K_{Y_1})^3$ is bounded. Thus $(-K_X)^3$ is bounded.

If $\dim Z = 1$ then the fibers are smooth del Pezzo surfaces. Because they are rational and form a bounded family, we can choose a covering family of curves $\{L\}$ on Y_1 such that a general L belongs to a general fiber, connects two general points of the fiber, and has bounded intersection with the anticanonical class. We can then pull it back to Y and down to X . This family would obviously satisfy our requirements. Note also that this would be a coextremal ray on Y in the terminology of Batyrev [3].

If $\dim Z = 2$ then we claim that the family of fibers of the fibration is not $\{l\}$. Suppose it is $\{l\}$. If at any step of the program we contract a prime divisor intersecting $\{l\}$ then $\{l\}$ would cease to be free and would never become free again. Thus no such contraction ever occurred. But this means that

$$(K_{Y_1} + \frac{1}{m}F_{Y_1}) \cdot l = (K_Y + \frac{1}{m}F_Y) \cdot l$$

The left hand side of the above equality is negative, while the right hand side is zero by the definition of F and because l doesn't pass through the singularities of X .

So the family of fibers $\{L\}$ is not $\{l\}$. It is a free family on Y_1 and $-K_{Y_1} \cdot L = 2$. By pulling it back to Y we get a free family on Y with the same intersection with $-K_Y$. We can then glue it with $\{l\}$ (see [17]) to get the desired covering family of rational curves. \square

4. Completion of the Proof

First of all, let's define some number $M = M(n)$ which will play a crucial role in the proof of the main theorem.

LEMMA 4.1. *For every n there exists some constant $M \geq 4320n$ such that for any del Pezzo surface with log-terminal singularities of index dividing n , and any divisor $D \in |-4320nK_S|$, the pair $(S, \frac{1}{M}D)$ is a log-terminal pair.*

PROOF. It follows essentially from the boundedness of del Pezzo surfaces of given index. First of all, the multiplicity of any of the irreducible components of D can not be too big because $H \cdot D$ is bounded for some very ample H on X (which can be actually chosen to be some fixed multiple of $(-nK_S)$). So for big enough M , $(S, \frac{1}{M}D)$ is a log pair, i.e. $\frac{1}{M}D$ is a boundary. Moreover, on the minimal resolution $\pi : Y \rightarrow S$ the exceptional divisors have simple normal crossings. For every such exceptional divisor E there is a covering family of curves $\{L\}$ on Y that intersect E , such that $L \cdot (-K_S)$ is bounded. This implies that for every divisor D as above, the coefficient of E in π^*D is bounded. So for each Y we have only finitely many possibilities for the linear system that the strict preimage D' of D belongs to. Altogether we have a bounded family of pairs (Y, π^*D) . For each pair there exists the minimal number $M(Y, \pi^*D)$ such that the pair $(S, \frac{1}{M}D)$ is log-terminal. This M can be found by successive blow-ups of Y . Clearly, it is Zariski semi-continuous as a function of the pair (Y, π^*D) , so it is uniformly bounded. \square

REMARK. The bound in the above lemma is not explicit. It would be very interesting to find an explicit bound for M in terms of n .

In the remainder of this section we consider the Fano threefolds X with log-terminal singularities of index n , which are equipped with the family of rational curves $\{l\}$ from the statement of Theorem 3.1. Let us denote the family of fibers of its rationally connected fibration by $\{S\}$. This is a LINEAR system of divisors (cf. [4]).

We have the following easy proposition.

PROPOSITION 4.1. *In the above situation and notations, there exist constants c_1 and c_2 depending only on the degree $H \cdot l$ such that for a general S*

$$(H|_S)^2 \leq c_1, \text{ and } h^0(H|_S) \leq c_2.$$

PROOF. Possibly gluing $\{l\}$ with itself once, we may assume that two general points of S are connected by some curve l . Suppose $l \cdot H = m$. By Lemma 2.2 of [4], $(H|_S)^2 \leq m^2$. To bound $h^0(H|_S)$, consider $(m+1)$ curves l and $(m+1)$ points on each of them. If a section of $H^0(H|_S)$ vanishes at all these points, it has to vanish at all these curves. This implies that it is identically zero on S , by intersecting with a general l , because $l \cdot l \geq 1$. So $h^0(H|_S) \leq (m+1)^2$.

THEOREM 4.1. *In the above situation and notations, suppose that the base locus of $|S_X|$ contains a curve. Then H^3 is bounded.*

PROOF. The divisor H is very ample on X . Consider the family of (probably not rational) curves $\{C\} = H|_S$. By the above proposition, $C \cdot H$ is bounded. Because X has log-terminal singularities of index n , and the base locus of $|S_X|$ contains a curve, we can find a point x_0 in this base locus, with multiplicity of the local ring bounded by $2n$. Then the curves C passing through this point are not contained in any closed subvariety of X . So H^3 is bounded (cf., e.g. [4], Lemma 2.2.) \square

REMARK. If X is \mathbb{Q} -factorial, and $\rho(X) = 1$, the conditions of the above theorem are always satisfied. So in this case one can use the above theorem instead of the argument of [4], section 4, to get a bound for $(-K_X)^3$, which is polynomial in n .

For the remainder of this section we will assume that the base locus of S_X contains no curves. If X is \mathbb{Q} -factorial, this implies that it is empty, because it is equal to the intersection of two general elements of the one-dimensional linear system S_X . In general, it could consist of finitely many points.

By the Log Minimal Model Program (cf., e.g. [14], 6.16; also [24]) there exists a small partial resolution of singularities $\pi : Y \rightarrow X$ such that Y is \mathbb{Q} -factorial, and S is π -nef. (If X is \mathbb{Q} -factorial then $Y = X$). Because S_Y is π -nef, the base locus of S_Y contains no exceptional curves of π , so

it is empty. Therefore, $|S_Y|$ is a free linear system, two general S_Y do not intersect. The variety Y has log-terminal singularities of index n . The anticanonical class $-K_Y = \pi^*(-K_X)$. So by trivial adjunction a general S_Y is a del Pezzo surface with log-terminal singularities of index dividing n .

By the Cone Theorem on Y , we have at least one of the following.

1. One of the extremal curves of π , say L , intersects S_Y .
2. There is an extremal contraction on Y whose fibers intersect S_Y .

In the second case we could have a fibration, a divisorial contraction, or a small contraction. In the fibration case, we can get a covering family of curves on the fibers with bounded intersection with $-K_Y$, whose rationally connected fibration is different from $|S|$. So we can glue together this family and $\{l\}$ to get a new family, which connects two general points of X and has bounded intersection with H . This implies boundedness of H^3 , so we are left with the following three possibilities.

1. There is a divisorial contraction on Y whose fibers intersect S .
2. There is an exceptional curve L on Y , such that $L \cdot K_Y < 0$, and L intersects S .
3. One of the extremal curves of π , say L , intersects S_Y . In this case $L \cdot K_Y = 0$.

The first case is the simplest. It is treated in the following proposition.

PROPOSITION 4.2. *Suppose X and Y are as above, and we have a divisorial contraction on Y . Then $h^0(H)$ is bounded.*

PROOF. Suppose E is the exceptional divisor of the contraction. As in [4], section 6, we can find a covering family $\{L\}$ on Y such that $-3 \leq K_{Y_1} \cdot L \leq 0$ and $-E \cdot L \leq 3$. (The variety Y in our case has log-terminal singularities rather than terminal, but the Subadjunction still works).

This implies that $L \cdot H$ is bounded. Suppose $L \cdot H = d$. By Proposition 4.1, $h^0(H|_S)$ is bounded. Suppose it is equal to e . Also fix a constant M from Lemma 4.1. Then $h^0(H) - h^0(H_Y - (3M + d + 1)S_Y) \leq (3M + d + 1)e$. If $h^0(H)$ is big enough, then there exists some $D \in |H_Y - (3M + d + 1)S_Y|$. Let us write $D = a \cdot E + F$, where $E \notin \text{Supp} F$. By Lemma 4.1 ($S, \frac{1}{M}D|_S$)

is a log pair. This means that $\frac{1}{M}D|_S$ is a boundary. Therefore $a \leq M$. The following chain of inequalities provides a contradiction.

$$\begin{aligned} d = L \cdot H &= (3M + d + 1)L \cdot S + L \cdot D \geq (3M + d + 1)L \cdot S + aL \cdot E \geq \\ &\geq (3M + d + 1)L \cdot S - 3M \geq (d + 1) \square \end{aligned}$$

Note that in this case we did not use the full strength of Lemma 4.1. In particular in this case we could actually choose M to be polynomial in n . We will however need the full strength of the lemma for the remaining two cases. We will treat them together in the following proposition.

PROPOSITION 4.3. *Suppose X and Y are as above, and we have a small contraction on Y with some fiber L intersecting S . Then $h^0(H)$ is bounded.*

PROOF. Suppose L is as above. Recall again a number M from Lemma 4.1. By Proposition 4.1, if $h^0(H_X) = h^0(H_Y)$ is big enough then $h^0(H_Y - (3M)S_Y) > 0$. Take some $F \in |H_Y - (3M)S_Y|$ and consider the divisor D which consists of all the components of F containing L .

By the definition of M the log pair $(Y, \frac{1}{M}D)$ is log-terminal in the neighborhood of the general point of L . Consider the log-terminal modification (Y_1, B_1) of (Y, B) , where $B = \frac{1}{M}D$ (cf. [14], 6.16, d=1). The log adjunction formula for the morphism $\pi_Y^{Y_1}$ is the following.

$$K_{Y_1} + B_1 = (\pi_Y^{Y_1})^*(K_Y + B) + \sum \alpha_i A_i,$$

where A_i are exceptional divisors of $\pi_Y^{Y_1}$ and all $\alpha_i \leq 0$ because $K_{Y_1} + B_1$ is relatively nef.

Because $(Y, \frac{1}{M}D)$ is log-terminal in the neighborhood of the general point of L , none of the divisors A_i lies above L . So we can consider the curve L_1 on Y_1 which is the closure of the pullback of the general point of L . Obviously, L_1 is an extremal curve on Y_1 , and $L_1 \cdot (K_{Y_1} + B_1) \leq L \cdot (K_Y + B)$. By Kawamata's theorem on the length of an extremal curve (cf. [11]), $-2 \leq L_1 \cdot (K_{Y_1} + B_1)$.

So we have the following chain of inequalities.

$$-2 \leq L_1 \cdot (K_{Y_1} + B_1) \leq L \cdot (K_Y + B) \leq L \cdot (K_Y + \frac{1}{M}F) =$$

$$= L \cdot (K_Y + \frac{1}{M}(H_Y - 3M \cdot S)) = L \cdot K_Y \cdot (1 - \frac{4320n}{M}) - 3L \cdot S \leq -3$$

The above contradiction completes the proof. \square

References

- [1] Alexeev, V., General elephants of Q-Fano 3-folds, *Compositio Math.* **91** (1994), 91–116.
- [2] Alexeev, V., Boundedness and K^2 for log surfaces, *Internat. J. Math.* **5** (1994), no. 6, 779–810.
- [3] Batyrev, V. V., The cone of effective divisors of threefolds, *Contemporary Math.* **131** (1989), Part 3, 337–352.
- [4] Borisov, A., Boundedness theorem for Fano log-threefolds, *J. Algebraic Geom.* **5** (1996), no. 1, 119–133.
- [5] Borisov, A. and L. Borisov, Singular toric Fano varieties, *Acad. Sci. USSR Sb. Math.* **75** (1993), no. 1, 277–283.
- [6] Bruno, A. and K. Matsuki, Log Sarkisov Program, *Internat. J. Math.* **8** (1997), no. 4, 451–494.
- [7] Campana, F., Une version géométrique généralisée du théorème du produit de Nadel, *Bull. Soc. Math. France* **119** (1991), no. 4, 479–493.
- [8] Clemens, H., Kollár, J. and S. Mori, Higher dimensional complex geometry, *Astérisque* **166** (1988).
- [9] Iskovskikh, V. A. and Yu. G. Prokhorov, Fano varieties, *Algebraic Geometry, V, 1–247, Encyclopaedia Math. Sci., 47*, Springer, Berlin, 1999.
- [10] Kawamata, Y., Boundedness of Q-Fano threefolds, *Contemporary Math.* **131** (1989), Part 3, 439–445.
- [11] Kawamata, Y., On the length of an extremal rational curve, *Invent. Math.* **105** (1991), no. 3, 609–611.
- [12] Kawamata, Y., Matsuda, K. and K. Matsuki, Introduction to the Minimal Model Program, “Algebraic geometry, Sendai”, *Adv. Stud. Pure Math.* **10** (1987), 283–360.
- [13] Kollár, J., Effective base point freeness, *Math. Ann.* **296** (1993), no. 4, 595–605.
- [14] Kollár, J., et al., Flips and abundance for algebraic threefolds, *Astérisque* **211** (1992).
- [15] Kollár, J. and T. Matsusaka, Riemann-Roch type inequalities, *Amer. J. Math.* **105** (1983), no. 1, 229–252.
- [16] Kollár, J., Miyaoka, Y. and S. Mori, Rational connectedness and boundedness of Fano manifolds, *J. Differential Geom.* **36** (1992), no. 3, 765–779.
- [17] Kollár, J., Miyaoka, Y. and S. Mori, Rationally connected varieties, *J. Algebraic Geom.* **1** (1992), 429–448.

- [18] Kollár, J., Miyaoka, Y., Mori, S. and H. Takagi, Boundedness of Canonical \mathbb{Q} -Fano 3-Folds, preprint(2000). RIMS preprint series, # 1273. Web address: <http://www.kurims.kyoto-u.ac.jp/kenkyubu/paper/>.
- [19] Miyaoka, Y. and S. Mori, A numerical criterion for uniruledness, *Ann. of Math.* **124** (1986), 65–69.
- [20] Nadel, A. M., The boundedness of degree of Fano manifold with Picard number one, *J. Amer. Math. Soc.* **4** (1991), no. 4, 681–692.
- [21] Nikulin, V. V., Del Pezzo surfaces with log terminal singularities, *Mat. Sb.* **180** (1989), no. 2, 226–243.
- [22] Ziv Ran, On semipositivity of sheaves of differential operators and the degree of a unipolar \mathbb{Q} -Fano variety, preprint. Web address: <http://front.math.ucdavis.edu/math.AG/9811022>.
- [23] Ziv Ran, Herb Clemens. A new method in Fano geometry, preprint. Web address: <http://front.math.ucdavis.edu/math.AG/0001120>.
- [24] Shokurov, V. V., 3-fold log flips, *Izv. Ross. Akad. Nauk Ser Mat.* **56** (1992), 105–203; *Russian Acad. Sci. Izv. Math.* **40** (1993), 95–202.
- [25] Tsuji, H., Boundedness of \mathbb{Q} -Fano varieties of Picard number one, preprint. Web address: <http://front.math.ucdavis.edu/math.AG/9903043>.

(Received July 25, 2000)
(Revised January 31, 2001)

Department of Mathematics
Penn State University
University Park
PA 16802, USA
E-mail: borisov@math.psu.edu