

A Note on Semistable Barsotti-Tate Groups

By Fabien TRIHAN

Abstract. We show that the Dieudonné crystal associated to a Barsotti-Tate group with potentially semistable reduction over a smooth curve is overconvergent. As a corollary, we obtain the rationality of the L -function associated to this group.

1. Introduction

Let U/\mathbb{F}_p be a smooth curve and G/U a Barsotti-Tate group. Assume G/U has potentially semistable reduction (see 4.2 for a precise definition). We show that the Dieudonné crystal as defined in [1] is overconvergent in the sense of Berthelot. As a corollary we get the rationality of the L -function associated to G/U . In the third section we study the local situation, that is, semistable Barsotti-Tate groups over a complete discrete valuation field of equal characteristic p . Using Extension groups in the category of Dieudonné crystals and their interpretation in terms of syntomic cohomology (as defined in [13]) we prove that the Dieudonné crystal associated to such group extends to a log Dieudonné crystal over the ring of integers. Using the gluing properties of overconvergent F -isocrystals over smooth curves proved in [14], we deduce from section three the overconvergence of the Dieudonné crystal associated to G/U and the rationality of its L -function in the last section. We end both sections three and four by some open questions.

2. Acknowledgments

This project started in 2003 at the University of Kyoto following three days talk with Professor Kato. I am very grateful to him. I also would like to express my gratitude to O. Brinon, L. Hesselholt, I. Longhi, M.-H. Nicole, A. Shiho and T. Tsuji for very fruitful conversations. The author would like also to thank the referee for his careful reading and for his suggestion to improve the redaction of the key lemma 3.8. This paper has been written

2000 *Mathematics Subject Classification.* 14F30.

during a stay at the University of Tokyo. I thank again T. Saito for his great hospitality. The author has been supported by the Royal Society.

3. Semistable Barsotti-Tate Groups and Extensions

In this section, we extend the Dieudonné crystal of a semistable Barsotti-Tate group over a complete discrete valuation field of equal characteristic $p > 0$ to a log Dieudonné crystal.

3.1. Let k be a perfect field of characteristic p endowed with its Frobenius σ , $W := W(k)$ the ring of Witt vectors and $K = \text{Frac}(W)$. We denote $\eta := \text{Spec}(k((t)))$. Let G_η/η a Barsotti-Tate group. Following [5]:

3.2 DEFINITION. The Barsotti-Tate group G_η/η is called semistable if there exists a filtration:

$$0 \subset G_\eta^\mu \subset G_\eta^f \subset G_\eta$$

by Barsotti-Tate groups such that the following conditions hold:

1. G_η^f and G_η/G_η^μ extend to Barsotti-Tate groups G_1 and G_2 over $k[[t]]$. In this case, the composed map

$$G_\eta^f \hookrightarrow G_\eta \rightarrow G_\eta/G_\eta^\mu$$

extends to a map $G_1 \rightarrow G_2$.

2. $G_1^\mu := \text{Ker}(G_1 \rightarrow G_2)$ and $G_2^{\acute{e}t} := \text{coker}(G_1 \rightarrow G_2)$ are Barsotti-Tate groups over $k[[t]]$.
3. G_1^μ is of multiplicative type and $G_2^{\acute{e}t}$ is étale.

3.3 REMARK. It has been shown in [5], 2.5, that an abelian variety A over η has semistable reduction if and only if its associated Barsotti-Tate group $G_\eta := \varinjlim_n A[p^n]$ is semistable.

3.4. Let S be a fine log-scheme over $\text{Spec}(k)$ endowed with the trivial log-structure. We denote the absolute Frobenius of S by σ_S , lying above σ . We work on the log crystalline site with the étale topology, denoted

$\mathit{Crys}(S/W)$ ([7]). An object of $\mathit{Crys}(S/W)$ is a pair (S', P) , where S' is an étale scheme over S , P is a p.d.-thickening of S' over W with respect to the p.d.-structure of (p) , and we are given an isomorphism between the inverse image of the log structure of P on S' and the inverse image of the log structure of S on S' . Morphisms of $\mathit{Crys}(S/W)$ are defined in the evident way. The topology of $\mathit{Crys}(S/W)$ is given by the étale topology of each P . In the applications of this paper, S is mainly one of the followings :

1. S/k is a proper smooth curve with the log structure on S associated to the divisor $S \setminus U$, for some open subset U .
2. $S = U$ with the trivial log structure.
3. $S = \mathrm{Spec}(k[[t]])$ with the log structure associated to the closed point.
4. $S = \eta$ with the trivial log structure.

3.5. A crystal E on $\mathit{Crys}(S/W)$ is called a Dieudonné crystal if it is a finite locally free crystal endowed with linear operators $F : \sigma_S^* E \rightarrow E$ and $V : E \rightarrow \sigma_S^* E$ called respectively Frobenius and Verschiebung such that $FV = p$ and $VF = p$. If (D, F_D, V_D) is a Dieudonné crystal on $\mathit{Crys}(S/W)$, its $\mathcal{O}_{S/W}$ -dual D^\vee is endowed with a structure of Dieudonné crystal such that $F_{D^\vee} = (V_D)^\vee$ and $V_{D^\vee} = (F_D)^\vee$.

3.6. Let G be a Barsotti-Tate group over S . By the crystalline Dieudonné theory (see for example [1], [2], [4]), the Dieudonné crystal $\mathbb{D}(G)$ on $\mathit{Crys}(S/W)$ is defined by forgetting the log structures of objects of $\mathit{Crys}(S/W)$ (\mathbb{D} is a contravariant functor). More precisely, let π denote the canonical morphism from S to S_{triv} , the scheme S endowed with the trivial log-structure. Then $\pi^* \mathbb{D}(G)$ is a Dieudonné crystal on $\mathit{Crys}(S/W)$ that we still denote $\mathbb{D}(G)$. The $\mathcal{O}_{S/W}$ -dual of $\mathbb{D}(G)$ will be denoted by $D(G)$, so that $D(\cdot)$ becomes a covariant functor. We will furthermore, denote by $\mathbf{1} := D(\mathbb{Q}_p/\mathbb{Z}_p)$ the Dieudonné crystal $(\mathcal{O}_{S/W}, F = p, V = id)$ and by $\mathbf{1}(1) := D(\mu_{p^\infty})$ the Dieudonné crystal $(\mathcal{O}_{S/W}, F = id, V = p)$. The Dieudonné crystals $\mathbf{1}$ and $\mathbf{1}(1)$ are dual to each other.

3.7. We recall the construction of the syntomic cohomology as defined in [13] in the case $S = \mathrm{Spec}(k[[t]])$ with the log structure associated to

the closed point. Let D be a Dieudonné crystal over S/W . The syntomic complex \mathcal{S}_D is the total complex associated to the bicomplex

$$\begin{array}{ccccc} & D^0 & \xrightarrow{\nabla} & D \otimes \Omega_{\mathcal{Y}}^1 & \\ \mathbf{1} - F_1 & \downarrow & & \downarrow & \mathbf{1} - F_2 \\ & D & \xrightarrow{\nabla} & D \otimes \Omega_{\mathcal{Y}}^1 & \end{array}$$

We explain the notations: $\mathcal{Y} = Spf(W[[t]])$ is endowed with the log-structure associated to $\mathbb{N} \rightarrow W[[t]]$ sending n to t^n . It is a log smooth formal lifting of S and we denote $\sigma_{\mathcal{Y}}$ a lifting of the Frobenius of S sending the variable t to t^p . By abuse of notation, we still denote (D, ∇, F_D, V_D) the realization of the Dieudonné crystal D at the p.d. thickening $(S \subset \mathcal{Y})$ endowed with its connection, Frobenius and Verschiebung. Consider the composed map

$$D \xrightarrow{\iota} \sigma_{\mathcal{Y}}^* D \rightarrow \sigma_{\mathcal{Y}}^* D / V_D(D)$$

where ι is the map sending $x \rightarrow 1 \otimes x$. Set $Lie(D)$ to be the image of the above map. Then $Lie(D)$ is a locally free \mathcal{O}_S -module (see [13], 5.3) and we denote D^0 , the kernel of the surjective map $D \rightarrow Lie(D)$. Finally, we explain the Frobenius operators. The map $F_1 : D^0 \rightarrow D$ is constructed as follows: the composed map

$$\tilde{F}_1 : D^0 \xrightarrow{\mathbf{1}} D \xrightarrow{\iota} \sigma_{\mathcal{Y}}^* D \xrightarrow{F_D} D,$$

is in $p \cdot D$ (see [13], 5.8.1) and we set $F_1 := p^{-1} \tilde{F}_1$. On the other side, remark that $\sigma_{\mathcal{Y}}(\Omega_{\mathcal{Y}}^1) \subset p \cdot \Omega_{\mathcal{Y}}^1$ so that we can define a map

$$F_2 := F_D \circ \iota \otimes p^{-1} \sigma_{\mathcal{Y}}.$$

3.8 PROPOSITION. *Assume k is algebraically closed and let $S = \text{Spec}(k[[t]])$ endowed with the log structure associated to the closed point. Then, we have:*

$$H^i(S, \mathcal{S}_{\mathbf{1}(1)}) = H^i(\eta, \mathcal{S}_{\mathbf{1}(1)}) = \begin{cases} \widehat{k((t))}^{\times}, & i = 1 \\ 0, & \text{otherwise,} \end{cases}$$

where $\hat{M} = \varprojlim_n M/M^{p^n}$ for any multiplicative group M .

PROOF. First, we prove the claim for $H^i(\eta, \mathcal{S}_{\mathbf{1}(1)})$. By [13], 5.10, we have

$$H^i(\eta, \mathcal{S}_{\mathbf{1}(1)}) = H_{fl}^i(\eta, T_p \mathbf{G}_m).$$

Since $k((t))$ is a C_1 -field, we have

$$H_{fl}^i(\eta, \mathbf{G}_m) = \begin{cases} k((t))^\times & , i = 1 \\ 0, & \text{otherwise.} \end{cases}$$

By using the short exact sequence

$$0 \rightarrow \mu_{p^n} \rightarrow \mathbf{G}_m \xrightarrow{p^n} \mathbf{G}_m \rightarrow 0$$

on the flat site, we see that

$$H_{fl}^i(\eta, T_p \mathbf{G}_m) = \begin{cases} \widehat{k((t))}^\times & , i = 1 \\ 0, & \text{otherwise.} \end{cases}$$

So the claim for $H^i(\eta, \mathcal{S}_{\mathbf{1}(1)})$ is proved.

Next we prove the claim for $H^i(S, \mathcal{S}_{\mathbf{1}(1)})$. In the case of the crystal $D = \mathbf{1}(1)$ the short exact sequence

$$0 \rightarrow D^0 \rightarrow D \rightarrow Lie(D) \rightarrow 0$$

is induced by the canonical short exact sequence in the crystalline site:

$$0 \rightarrow \mathcal{I}_{S/W} \rightarrow \mathcal{O}_{S/W} \rightarrow \mathbf{G}_a \rightarrow 0$$

which induces on the pd-thickening $S \subset \mathcal{Y}$ the short exact sequence:

$$0 \rightarrow p.W[[t]] \rightarrow W[[t]] \rightarrow k[[t]] \rightarrow 0.$$

Hence, the syntomic complex of $\mathbf{1}(1)$ over S is the total complex associated to the bicomplex

$$\begin{array}{ccccc} & p.W[[t]] & \xrightarrow{d} & W[[t]] \frac{dt}{t} & \\ \mathbf{1} - F_1 & \downarrow & & \downarrow & \mathbf{1} - F_2 \\ & W[[t]] & \xrightarrow{d} & W[[t]] \frac{dt}{t} & \end{array}$$

where $d : W[[t]] \rightarrow W[[t]] \frac{dt}{t}$ is the map sending an element $\sum_i a_i t^i$ to $(\sum_i i a_i t^i) \frac{dt}{t}$, F_1 is the map sending an element $p \cdot \sum_i a_i t^i$ to $\sum_i \sigma(a_i) t^{pi}$ and F_2 the map sending an element $(\sum_i a_i t^i) \frac{dt}{t}$ to $(\sum_i \sigma(a_i) t^{pi}) \frac{dt}{t}$. Hence, $\mathcal{S}_{1(1)}$ is the complex concentrated in degree 0, 1, 2:

$$[pW[[t]] \xrightarrow{d, 1-F_1} W[[t]] \frac{dt}{t} \oplus W[[t]] \xrightarrow{1-F_2, -d} W[[t]] \frac{dt}{t}].$$

Remark that this complex is isomorphic to the complex

$$[W[[t]] \xrightarrow{p d, p-\sigma} W[[t]] \frac{dt}{t} \oplus W[[t]] \xrightarrow{1-F_2, -d} W[[t]] \frac{dt}{t}].$$

We compute the H^0 : By definition $H^0 = \text{Ker}(d) \cap \text{Ker}(1 - F_1)$. Since $\text{Ker}(d) = pW$, H^0 is equal to the set of element $p \cdot a \in pW$ such that $pa - \sigma(a) = 0$. Since the p -adic valuation $v(\sigma(a))$ is equal to $v(a)$, the previous equality gives $a = 0$.

We compute the H^2 : to show that this is zero, we just need to show that the map $\pi := (1 - F_2, -d)$ is surjective. But for any $\sum_i c_i t^i \frac{dt}{t} \in W[[t]] \frac{dt}{t}$, the element $(\sum_i b_i t^i \frac{dt}{t}, 0)$, with $b_i = c_i + \sigma(b_{i/p})$ if p divide i and $b_i = c_i$ else is an antecedent of $\sum_i c_i t^i \frac{dt}{t}$ by π .

We now turn to the computation of $H^1 := \text{Ker}(\pi) / \text{Im}(d, 1 - F_1)$. The group $\text{Ker}(\pi)$ is the set of elements $(\sum_i a_i t^i \frac{dt}{t}, \sum_i b_i t^i)$ such that $a_0 \in \mathbb{Z}_p$ and for n , any positive integer with p -adic valuation r , $a_n = nb_n + (n/p)\sigma(b_{n/p}) + \dots + (n/p^r)\sigma^r(b_{n/p^r})$. We get this way an isomorphism

$$\text{Ker}(\pi) \simeq \mathbb{Z}_p \frac{dt}{t} \oplus W[[t]]$$

by sending $(\sum_i a_i t^i \frac{dt}{t}, \sum_i b_i t^i)$ to $(a_0, \sum_i b_i t^i)$, which induces an isomorphism

$$\text{Im}(d, 1 - F_1) \simeq 0 \oplus \text{Im}(1 - F_1),$$

since the elements in $\text{Im}(d)$ have no constant terms.

We get

$$H^1 = \mathbb{Z}_p \frac{dt}{t} \oplus W[[t]] / \text{Im}(1 - F_1).$$

On the other hand, $k((t))^\times \simeq t^\mathbb{Z} \times k^\times \times (1 + tk[[t]])$ and $(k((t))^\times)^{p^n} = k((t^{p^n}))^\times \simeq t^{p^n \mathbb{Z}} \times k^\times \times (1 + t^{p^n} k[[t^{p^n}]])$. So, we are reduced to identify

$W[[t]]/Im(1 - F_1)$ and $\varprojlim_n (1 + tk[[t]]/(1 + t^{p^n} k[[t^{p^n}]])$. We first prove that the lefthand side is p -adically complete. By, [15], chapter 8, it is enough to prove that $I = Im(1 - F_1)$ is closed and in particular complete. Let $(f_m(t) = \sum_i b_i^{(m)} t^i)_{m \in \mathbb{N}}$ a sequence of elements in I converging to $f(t) = \sum_i b_i t^i \in W[[t]]$. We want to show that $f(t)$ is in fact in I . Since $f_m(t) \in I$, for any m , there exists some sequence $(a_i^{(m)})_i \in W^{\mathbb{N}}$ such that $b_i^{(m)} = pa_i^{(m)} - \sigma(a_{\frac{i}{p}}^{(m)})$ if p divides i and $b_i^{(m)} = pa_i^{(m)}$ else. We construct by induction on the p -adic valuation of i , a sequence $(a_i)_i \in W^{\mathbb{N}}$ such that $(1 - F_1)(\sum_i pa_i t^i) = f(t)$. For $v_p(i) = 0$, that is when p does not divide i , $pa_i^{(m)}$ converges when m goes to infinity to b_i so that $(a_i^{(m)})$ converges to an element $a_i \in W$ such that $b_i = pa_i$. Assume now that for any i such that $v_p(i) \leq r$, $(a_i^{(m)})$ converges to an element $a_i \in W$. Then, if $v_p(i) = r + 1$, we have $b_i^{(m)} = pa_i^{(m)} - \sigma(a_{\frac{i}{p}}^{(m)})$, with $(b_i^{(m)})_m$ converging to an element b_i and by induction hypothesis, $(\sigma(a_{\frac{i}{p}}^{(m)}))_m$ converging to an element $\sigma(a_{\frac{i}{p}})$ and so we deduce that $(a_i^{(m)})$ converges to an element $a_i \in W$.

Let $D = \mathbf{1}(1)$. We compute now $H^1(S, \mathcal{S}_D)/p^n$: we have a short exact sequence

$$0 \rightarrow \mathcal{S}_D \xrightarrow{\times p^n} \mathcal{S}_D \rightarrow \mathcal{S}_{D,n} \rightarrow 0,$$

which induces an exact sequence

$$H^1(S, \mathcal{S}_D) \xrightarrow{\times p^n} H^1(S, \mathcal{S}_D) \rightarrow H^1(S, \mathcal{S}_{D,n}) \rightarrow H^2(S, \mathcal{S}_D).$$

Since we already have proved that $H^2(S, \mathcal{S}_D) = 0$, we deduce for any n the isomorphisms

$$H^1(S, \mathcal{S}_D)/p^n \simeq H^1(S, \mathcal{S}_{D,n}).$$

By [13], 5.14.6, we also have

$$H^1(\eta, \mathcal{S}_D)/p^n \simeq H^1(\eta, \mathcal{S}_{D,n}).$$

Again, by using the short exact sequence:

$$0 \rightarrow \mathcal{S}_{D,1} \rightarrow \mathcal{S}_{D,n+1} \xrightarrow{\times p} \mathcal{S}_{D,n} \rightarrow 0$$

and the 5-lemma, we are reduced by induction to prove that

$$H^1(S, \mathcal{S}_{D,1}) \simeq H^1(\eta, \mathcal{S}_{D,1}).$$

Using the second description of the syntomic complex, we have the quasi-isomorphisms:

$$\begin{aligned} \mathcal{S}_{\mathbf{1}(1),S} \otimes \mathbb{Z}/p &\simeq [k[[t]] \xrightarrow{0, -\sigma} k[[t]] \frac{dt}{t} \oplus k[[t]] \xrightarrow{\pi_S} k[[t]] \frac{dt}{t}], \\ \mathcal{S}_{\mathbf{1}(1),\eta} \otimes \mathbb{Z}/p &\simeq [k((t)) \xrightarrow{0, -\sigma} k((t)) \frac{dt}{t} \oplus k((t)) \xrightarrow{\pi_\eta} k((t)) \frac{dt}{t}] \end{aligned}$$

and the map $H^1(S, \mathcal{S}_{\mathbf{1}(1)})/p \rightarrow H^1(\eta, \mathcal{S}_{\mathbf{1}(1)})/p$ is induced by the natural inclusion

$$k[[t]] \frac{dt}{t} \oplus k[[t]] \hookrightarrow k((t)) \frac{dt}{t} \oplus k((t)).$$

Now, we compute $H^1(\eta, \mathcal{S}_{\mathbf{1}(1)})/p$. For any element $(\sum_i a_i t^i \frac{dt}{t}, \sum_i b_i t^i) \in \text{Ker}(\pi_\eta)$ we find the same conditions that $a_0 \in \mathbb{F}_p$ and for n , any positive integer with p -adic valuation r , $a_n = nb_n + (n/p)\sigma(b_{n/p}) + \dots + (n/p^r)\sigma^r(b_{n/p^r})$. For negative integers and working modulo $\text{Im}(\sigma) = k((t^p))$, we claim that only the b_j 's with $b_{-j} = 0$ for any j prime to p , gives a solution. Namely, for such j we have $a_{-j} = -jb_{-j}$ but then $a_{-jp^k} = \sigma(-jb_{-j})$ for any positive integer k . But since $\sum_i a_i t^i \in k((t))$, we must have $a_{-jp^k} = 0$ for k big enough. Therefore, the canonical inclusion

$$k[[t]] \frac{dt}{t} \oplus k[[t]] \hookrightarrow k((t)) \frac{dt}{t} \oplus k((t))$$

induces the identity map

$$H^1(S, \mathcal{S}_D)/p = \mathbb{F}_p \frac{dt}{t} \oplus k[[t]]/k[[t^p]] \rightarrow \mathbb{F}_p \frac{dt}{t} \oplus k[[t]]/k[[t^p]] = H^1(\eta, \mathcal{S}_D)/p.$$

Hence, we proved the canonical isomorphism $H^1(S, \mathcal{S}_D) \simeq H^1(\eta, \mathcal{S}_D)$ and so the proof of the proposition is finished. \square

3.9. Let D_1, D_2 some Dieudonné crystals over S/W . We will denote $\text{Ext}_{S/W}(D_1, D_2)$ (or $\text{Ext}(D_1, D_2)$ if there is no ambiguity) the isomorphism classes of extensions

$$0 \rightarrow D_2 \rightarrow ? \rightarrow D_1 \rightarrow 0$$

in the category of Dieudonné crystals over S/W . Any commutative diagram

$$\begin{array}{ccc} S' & \xrightarrow{f} & S \\ \downarrow & & \downarrow \\ \text{Spec}(W) & \xrightarrow{g} & \text{Spec}(W) \end{array}$$

induces in the crystalline topos a functor $f^* : (S'/W)_{crys} \rightarrow (S/W)_{crys}$ allowing to define for any Dieudonné crystals D_1 and D_2 over S/W a canonical map

$$f^* : Ext^1(D_1, D_2) \rightarrow Ext^1(f^* D_1, f^* D_2),$$

sending the isomorphism class of an extension:

$$0 \rightarrow D_2 \rightarrow ? \rightarrow D_1 \rightarrow 0$$

to the isomorphism class of the extension

$$0 \rightarrow f^* D_2 \rightarrow f^* ? \rightarrow f^* D_1 \rightarrow 0.$$

(The exactness of this sequence follows from the local freeness of D_1 .)

3.10. Let G_η/η a semistable Barsotti-Tate group and denote as in 3.2 $G_\eta^f, G_\eta^\mu, G_1, G_1^\mu, G_2$ and $G_2^{ét}$ its associated Barsotti-Tate groups. We denote $S := \text{Spec}(k[[t]])$ endowed with the log-structure induced by its closed point. We also denote $j : \eta \rightarrow \text{Spec}(k[[t]])$ the open immersion. Then there is a commutative diagram of exact sequences:

$$\begin{array}{ccccc} Ext(D(G_2^{ét}), D(G_1^\mu)) & \xrightarrow{f_{log}} & Ext(D(G_2^{ét}), D(G_1)) & \xrightarrow{g_{log}} & Ext(D(G_2^{ét}), D(G_1/G_1^\mu)) \\ h_1 \downarrow & & h_2 \downarrow & & h_3 \downarrow \\ Ext(D(G_\eta/G_\eta^f), D(G_\eta^\mu)) & \xrightarrow{f} & Ext(D(G_\eta/G_\eta^f), D(G_\eta^f)) & \xrightarrow{g} & Ext(D(G_\eta/G_\eta^f), D(G_\eta^f/G_\eta^\mu)) \end{array}$$

where the horizontal maps are defined by applying the functor $\mathbb{R}Hom(D(G_2^{ét}), \cdot)$ and $\mathbb{R}Hom(D(G_\eta/G_\eta^f), \cdot)$ to the short exact sequences:

$$0 \rightarrow D(G_1^\mu) \rightarrow D(G_1) \rightarrow D(G_1/G_1^\mu) \rightarrow 0,$$

and

$$0 \rightarrow D(G_\eta^\mu) \rightarrow D(G_\eta^f) \rightarrow D(G_\eta^f/G_\eta^\mu) \rightarrow 0$$

of Dieudonné crystals over $(S/W)_{crys}$ and $(\eta/W)_{crys}$ respectively. The vertical maps are induced by the functor $j^* : (S/W)_{crys} \rightarrow (\eta/W)_{crys}$.

3.11 LEMMA. *Assume k is algebraically closed. Then the map g_{\log} is surjective.*

PROOF. Since k is algebraically closed, $G_2^{\acute{e}t} \simeq (\mathbb{Q}_p/\mathbb{Z}_p)^a$ and we can reduce to the case $a = 1$, that is to the case $D(G_2^{\acute{e}t}) = \mathbf{1}$. By [13], 5.9, $Ext(\mathbf{1}, D(G_1)) \simeq H^1(k[[t]], \mathcal{S}_{D(G_1)})$. Similarly, we have

$$Ext(\mathbf{1}, D(G_1/G_1^\mu)) \simeq H^1(k[[t]], \mathcal{S}_{D(G_1/G_1^\mu)})$$

so that the cokernel of g_{\log} is $H^2(k[[t]], \mathcal{S}_{D(G_1^\mu)})$. Again, since k is algebraically closed, we can reduce to the case $D(G_1^\mu) = \mathbf{1}(1)$ and the assertion results from 3.8. \square

3.12 LEMMA. *Assume that k is algebraically closed, then h_1 is an isomorphism.*

PROOF. We are reduced to prove that

$$Ext_{S/W}(\mathbf{1}, \mathbf{1}(1)) \simeq Ext_{\eta/W}(\mathbf{1}, \mathbf{1}(1)).$$

Using [13], 5.9 and 5.10, it is enough to prove that the map

$$H^1(S, \mathcal{S}_{\mathbf{1}(1)}) \rightarrow H^1(\eta, \mathcal{S}_{\mathbf{1}(1)})$$

is an isomorphism but this has already been proved in 3.8. \square

3.13 THEOREM. *Assume k is algebraically closed.*

Let $\alpha \in Ext(D(G_\eta/G_\eta^f), D(G_\eta^f))$ be the isomorphism class of the extension:

$$0 \rightarrow D(G_\eta^f) \rightarrow D(G_\eta) \rightarrow D(G_\eta/G_\eta^f) \rightarrow 0.$$

There exists a short exact sequence of Dieudonné crystals over S/W :

$$0 \rightarrow D(G_1) \rightarrow D_{\log} \rightarrow D(G_2^{\acute{e}t}) \rightarrow 0,$$

such that its isomorphism class β is sent by h_2 to α .

As a corollary, we get:

3.14 COROLLARY. *Let $G_\eta/\eta := k((t))$ be a semistable Barsotti-Tate group. Then its Dieudonné crystal $D(G_\eta)$ extends to a Dieudonné crystal*

D_{log} over S , the scheme $\text{Spec}(k[[t]])$ endowed with the log-structure induced by its closed point.

We now prove the theorem:

PROOF. Let $\gamma \in \text{Ext}(D(G_2^{ét}), D(G_1/G_1^\mu))$ be the isomorphism class of the extension:

$$0 \rightarrow D(G_1/G_1^\mu) \rightarrow D(G_2) \rightarrow D(G_2^{ét}) \rightarrow 0$$

such that we have $g(\alpha) = h_3(\gamma)$. Since g_{log} is surjective, there exists $\tilde{\gamma} \in \text{Ext}(D(G_2^{ét}), D(G_1))$ such that $g_{log}(\tilde{\gamma}) = \gamma$. Since $g(\alpha - h_2(\tilde{\alpha})) = 0$, there exists some $\delta \in \text{Ext}(D(G_\eta/G_\eta^f), D(G_\eta^\mu))$, corresponding by 3.12 to a unique $\tilde{\delta} \in \text{Ext}(D(G_2^{ét}), D(G_1^\mu))$, such that $f(\delta) = \alpha - h_2(\tilde{\alpha})$. Then $\beta := f_{log}(\tilde{\delta}) + \tilde{\gamma}$ is sent by h_2 to α . \square

3.15 DEFINITION. Let G_η/η be a Barsotti-Tate group. We say that it is overconvergent if its associated Dieudonné isocrystal, corresponding to a (φ, ∇) over

$$\mathcal{E} = \{a = \sum_{-\infty}^{+\infty} a_i x^i \mid a_i \in K, \sup_i |a_i| < \infty, |a_i| \rightarrow 0 (i \rightarrow -\infty)\}$$

(see [14]) admits a lattice as (φ, ∇) -module over

$$\mathcal{E}^\dagger = \{a \in \mathcal{E} \mid |a_i| r^i \rightarrow 0 (i \rightarrow -\infty) \text{ for a certain } r, 0 < r < 1\}.$$

As a corollary of 3.14, we have:

3.16 COROLLARY. Any semistable Barsotti-Tate group G_η/η is overconvergent.

3.17 REMARK.

1. Any Barsotti-Tate group coming from an abelian variety is overconvergent: the abelian variety has potentially semistable reduction and in consequence it has been shown in [13] that the Dieudonné crystal of the abelian variety (which coincides with the Dieudonné crystal of the associated Barsotti-Tate group) comes from a log Dieudonné crystal after taking some finite étale base change.

2. Barsotti-Tate groups associated to p -adic representations of the absolute Galois group of η with infinite monodromy are not overconvergent ([18]).
3. We have shown that if G_η/η is semistable then it is overconvergent. Reciprocally, if G_η/η is overconvergent, can we conclude that it is potentially semistable? Since G_η/η is overconvergent its associated isocrystal will be quasi-unipotent by the local p -adic monodromy theorem of André-Kedlaya-Mebkhout. So we know that it will come from some log-Dieudonné crystal after considering some finite étale base change. The previous question can thus be rephrased as: Is there a log Dieudonné functor from the category (still to be defined) of log p -divisible groups to the category of log Dieudonné modules over $k[[t]]$ and if yes, is this functor an equivalence of categories (as this is the case without log-structure by [4])?
4. Recall (see [19]) that G_η/η is endowed with a unique Frobenius slope filtration, whose quotients are isoclinic Barsotti-Tate groups. Assume each quotients to be overconvergent. Does it imply that G_η/η is overconvergent?

4. Semistable Barsotti-Tate Groups over Smooth Curves

4.1. In this section, we consider a dense open subset U of a proper smooth connected curve C/\mathbb{F}_p . For any closed point $x \in C$, we denote $\eta_x := \text{Spec}(k(x)((t)))$ and $K_x := \text{Frac}(W(k(x)))$. For any \mathbb{F}_p -scheme T , we will denote $\bar{T} := T \times_{\mathbb{F}_p} \bar{\mathbb{F}}_p$.

4.2 DEFINITION. A Barsotti-Tate group G/U is called semistable if at any closed point $x \in Z := C \setminus U$, $G_{\eta_x} := G \times_U \eta_x/\eta_x$ is semistable. We say that a Barsotti-Tate group G/U is potentially semistable if and only if there exists some finite Galois covering $U' \rightarrow U$ such that $G' := G \times_U U'/U'$ is semistable.

4.3. Let G/U a potentially semistable Barsotti-Tate group. We associate to G/U the following L -function:

$$L(U, G, t) := \prod_{x \in U} \det(1 - t^{\deg(x)} F_x, D(G_x))^{-1},$$

where $(D(G_x), F_x)$ is the F -isocrystal over $k(x)$ deduced from $D(G)$ by restriction and $\deg(x) := [k(x) : \mathbb{F}_p]$.

We are going to show that the Dieudonné crystal associated to a potentially semistable Barsotti-Tate group is overconvergent and that its associated L function is a rational function. We will need the following lemma:

4.4 LEMMA. ([6])

Let E be a convergent F -isocrystal over U . Let $\pi : \bar{U} \rightarrow U$ the canonical étale covering. Assume that π^*E is overconvergent, then E is overconvergent.

4.5 THEOREM. Let G/U a potentially semistable Barsotti-Tate group. Then its associated Dieudonné crystal $D(G)$ over U has a structure of overconvergent F -isocrystal $D(G)^\dagger$ over U .

PROOF. By the previous lemma we can assume $U = \bar{U}$ and by finite étale descent we can assume that G/U is semistable. For any closed point $x \in Z$, we denote $\eta_x := \text{Spec}(\text{Frac}(\hat{\mathcal{O}}_{C,x}))$ and $S_x := \text{Spec}(\hat{\mathcal{O}}_{C,x})$ endowed with the log-structure induced by its closed point. By 3.14, the Dieudonné crystal $D(\mathcal{G} \times_U \eta_x)$ extends to a Dieudonné crystal D_{\log} over S_x . Hence, the assertion follows from [14], proposition 4. \square

4.6 COROLLARY. Let G/U a potentially semistable Barsotti-Tate group. Then its L -function $L(G, U, t)$ is a rational function in t . More precisely, we have:

$$L(G, U, t) = \prod_{i=0}^2 \det(1 - tF, H_{\text{rig},c}^i(U, D(G)^\dagger))^{(-1)^{i+1}}.$$

PROOF. By 4.5 $D(G)$ has a structure of overconvergent F -isocrystal and the formula results from [17], theorem 1.2. Finally, the rationality results from the finiteness of the cohomological groups $H_{\text{rig},c}^i(U, D(G)^\dagger)$ which follows from [14], corollary 8 and the Poincaré duality of rigid cohomology. \square

4.7 REMARK. Let G_F/F be a Barsotti-Tate group, where F is the function field of C .

1. It is a priori not always possible to extend G_F/F to some Barsotti-Tate group G over some dense open subset U of C (but this is the case when G_F/F is the Barsotti-Tate group associated to an abelian variety). For example, consider the étale case. Then, we can replace Barsotti-Tate groups by p -adic representations. We can find an example of a p -adic representation of $\text{Gal}(\bar{F}/F)$ that ramifies at infinitely many places and thus don't factorize through any fundamental group of some dense open subset U of C . To construct such representation, it is enough to construct a \mathbb{Z}_p -extension K of F that ramifies at infinitely many places (it exists: see for example [11]). Take any extension L/F with Galois group $(\mathbb{Z}/p)^\times$. Then the extension $K.L/F$ has a Galois group isomorphic to \mathbb{Z}_p^\times and the natural projection $\text{Gal}(\bar{F}/F) \rightarrow \text{Gal}(K.L/F)$ gives an example of one-dimensional p -adic representation of $\text{Gal}(\bar{F}/F)$ that ramifies at infinitely many places.
2. If $G_1, G_2/U$ are two Barsotti-Tate groups with G_F/F as generic fiber, are G_1 and G_2 isomorphic (or at least isogenous)? See [5] for some evidences on this question. If the answer to this question is yes and G_F/F extends to some Barsotti-Tate group G/U , then we can define the Hasse-Weil L -function of G_F/F as $L(G_F, t) := L(U, G, t)$.

References

- [1] Berthelot, P., Breen, L. and W. Messing, Théorie de Dieudonné cristalline. II, Lecture Notes in Math., vol. 930, Springer-Verlag, New-york and Berlin, 1982.
- [2] Berthelot, P. and W. Messing, Théorie de Dieudonné cristalline. III, The Grothendieck Festschrift, Vol. I, Progr. Math., 86, Birkhauser Boston, Boston, MA, 1990.
- [3] Cartier, P., Groupes formels associés aux anneaux de Witt généralisés, C. R. Acad. Sci. Paris, Sér. **AB** (1967), 265.
- [4] de Jong, A. J., Crystalline Dieudonné module theory via formal and rigid geometry, Inst. Hautes études Sci. Publ. Math. **82** (1995), 5–96.
- [5] de Jong, A. J., Homomorphisms of Barsotti-Tate groups and crystals in positive characteristic, Invent. Math. **134** No. 2, (1998), 301–333; erratum *ibid.* **138** No. 1, (1999), 225.

- [6] Etesse, J.-Y., Descente étale des F -isocristaux surconvergers et rationalité des fonctions L de schmas abéliens, *Ann. Sci. cole Norm. Sup. (4)* **35** (2002), no. 4, 575–603.
- [7] Kato, K., *Logarithmic structures of Fontaine-Illusie*, Algebraic analysis, geometry and number theory, the Johns Hopkins Univ. Press (1989), 191–224.
- [8] Kato, K., *Logarithmic Dieudonné theory*, preprint, 1992.
- [9] Etesse, J.-Y. and B. Le Stum, Fonctions L associées aux F -isocristaux surconvergers II: Zéros et pôles unités, *Invent. math.* **127** (1997), 1–31.
- [10] Grothendieck, A., *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris), 3. Société Mathématique de France, Paris, 2003. xviii+327 pp.
- [11] Gold, R. and H. Kisilevsky, On geometric Z_p -extensions of function fields, *Manuscripta Math.* **62** (1988), no. 2, 145–161.
- [12] Hesselholt, L., *Lecture notes on Witt vectors*, preprint 2005, <http://www-math.mit.edu/~larsh/papers/s03/wittsurvey.pdf>.
- [13] Kato, K. and F. Trihan, On the conjectures of Birch and Swinnerton-Dyer in characteristic $p > 0$, *Invent. Math.* **153** (2003), no. 3, 537–592.
- [14] Matsuda, S. and F. Trihan, Image directe supérieure et unipotence. (French) [Higher direct image and unipotency], *J. Reine Angew. Math.* **569** (2004), 47–54.
- [15] Matsumura, H., *Commutative algebra*, Mathematics Lecture Note Series. New York: W. A. Benjamin, Inc. xii, 262 p. (1970).
- [16] Mumford, D., *Lectures on curves on an algebraic surface*, Annals of Mathematics Studies, vol. 59, Princeton University Press, Princeton, N.J., 1966.
- [17] Trihan, F., Fonction L unité d’un groupe de Barsotti-Tate, *Manuscripta Math.* **96** (1998), no. 4, 397–419.
- [18] Tsuzuki, N., Finite local monodromy of overconvergent unit-root F -isocrystals on a curve, *Amer. J. Math.* **120** (1998), no. 6, 1165–1190.
- [19] Zink, T., On the slope filtration, *Duke Math. J.* **109** (2001), no. 1, 79–95.

(Received September 16, 2008)

(Revised October 1, 2008)

School of Mathematics
University of Nottingham
University Park
NG7 2RD, Nottingham