

*A Limit Theorem on Maximum Value of Hedging with a Homogeneous Filtered Value Measure**

By Yuji UMEZAWA

Abstract. We study a hedging problem for an European contingent claim in a certain incomplete market model by using a homogeneous filtered value measure. We consider the minimal hedging risk in discrete time model and its continuous limit. As a result, we show that this limit is described by the unique viscosity solution of a kind of Hamilton-Jacobi-Bellman equation.

1. Introduction

The problem of hedging the risk of contingent claims in incomplete markets is important in financial mathematics. While it is still possible to stay on the safe side by using the classical superhedging strategy, the cost is often too high and one cannot actually do such hedging in many practical situations. In such situations, one of the most adequate way is to hedge the exposure from these claims in a way that the resulting risk remains within the prescribed limits. That is, one tries to compose the hedging strategies by using available assets in order to minimize the hedging error which is measured by some risk measure.

One of the right way to measure such risk is to use coherent risk measures, or more general convex risk measures. The concept of coherent risk measure is proposed by Artzner et al.[1], to assess the risk of such financial positions by an axiomatic approach. Föllmer and Schied[14] defined the class of convex risk measures by relaxing the axiom of homogeneity. Cheridito et al.[6] introduced coherent/convex risk measures for stochastic processes. Many authors such as Artzner et al.[2], Detlefsen, Scandolo[13], and Kusuoka, Morimoto[16] introduced the concept of dynamic risk measure in discrete settings. Delbaen[12] proposed the concept of dynamic risk

*This research is supported by the 21 century COE program at Graduate School of Mathematical Sciences, the University of Tokyo.

2010 *Mathematics Subject Classification.* Primary 91B30, Secondary 60F99, 91B24.

measure in a continuous setting. On hedging by using coherent or convex risk measures, many authors such as Barrieu and El Karoui[4][5], Cherny and Madan[9], Nakano[18], Roorda[19], and Umezawa[21] studied hedging strategies by using a coherent/convex risk measure in single-period settings. Recently, Cherny[8] gives results in discrete multi-period setting.

In this paper, we study a hedging for an European contingent claim by using a homogeneous filtered value measure in a multi-period setting. We first consider the minimizing problem of hedging risk in discrete multi-period time market model. The discussion of this part is similar to that of Cherny[8]. Next we consider its continuous limit. As a result, we prove that this limit is describe by the unique viscosity solution of some Hamilton-Jacobi-Bellman equation.

The structure of this paper is as follows. In the rest of this section, we introduce the notion of homogeneous filtered value measure[16], and state the claim of our main theorem. In Section 2, we consider a discrete time model and show a kind of Bellman's principle. In Section 4, we give the proof of our main theorem. In Appendix, we give a brief introduction and results on the theory of viscosity solution.

1.1. Definition

Let \mathcal{L} be the set of all probability measures on the Borel space $(\mathbf{R}, \mathcal{B}(\mathbf{R}))$, \mathcal{L}^∞ the set of $\nu \in \mathcal{L}$ which satisfies $\nu(\mathbf{R} \setminus [-M, M]) = 0$ for some $M > 0$, and \mathcal{M} the set of all probability measures on $([0, 1], \mathcal{B}[0, 1])$. For $\nu \in \mathcal{L}$, we denote the distribution function of ν by F_ν , i.e., $F_\nu(z) = \nu((-\infty, z])$, $z \in \mathbf{R}$. We define $Z : [0, 1] \times \mathcal{L} \rightarrow \mathbf{R}$ by $Z(x, \nu) = \inf\{z; F_\nu(z) > x\}$, $x \in [0, 1)$, $\nu \in \mathcal{L}$. For $\alpha \in (0, 1]$, We define $\eta_\alpha : \mathcal{L}^\infty \rightarrow \mathbf{R}$ by

$$(1) \quad \eta_\alpha(\nu) = \alpha^{-1} \int_0^\alpha Z(x, \nu) dx, \quad \nu \in \mathcal{L}^\infty,$$

and $\eta_0 : \mathcal{L}^\infty \rightarrow \mathbf{R}$ by

$$(2) \quad \eta_0(\nu) = \inf\{z \in \mathbf{R} \mid \nu((-\infty, z]) > 0\}, \quad \nu \in \mathcal{L}^\infty.$$

DEFINITION 1.1. We say that the mapping $\eta : \mathcal{L}^\infty \rightarrow \mathbf{R}$ is a mild value measure if there exists a subset $\mathcal{M}_0 \subset \mathcal{M}$ such that

$$\eta(\nu) = \inf_{m \in \mathcal{M}_0} \int_0^1 \eta_\alpha(\nu) m(d\alpha), \quad \nu \in \mathcal{L}^\infty.$$

DEFINITION 1.2. Let $\eta : \mathcal{L}^\infty \rightarrow \mathbf{R}$ be a mild value measure and (Ω, \mathcal{F}, P) a standard probability space.

(1) For any $X \in L^\infty(\Omega, \mathcal{F}, P)$ and any sub σ -algebra $\mathcal{G} \subset \mathcal{F}$, we define a \mathcal{G} -measurable random variable $\eta(X|\mathcal{G})$ by $\eta(X|\mathcal{G}) = \eta(P(X \in dx|\mathcal{G}))$, where $P(X \in dx|\mathcal{G})$ is a regular conditional probability law of X given a sub σ -algebra \mathcal{G} .

(2) For any $X \in L^\infty(\Omega, \mathcal{F}, P)$ and a filtration $\{\mathcal{F}_k\}_{k=0}^n$, we inductively define a $\{\mathcal{F}_k\}$ -adapted process $\{U_k\}_{k=0}^n$ by

$$\begin{aligned}
 U_n &= \eta(X|\mathcal{F}_n), \\
 (3) \quad U_{k-1} &= \eta(U_k|\mathcal{F}_{k-1}), \quad k = n, n-1, \dots, 1.
 \end{aligned}$$

We call U_k , $k = 0, 1, \dots, n$ a homogeneous filtered value measure for $X \in L^\infty$ at k . Also we denote U_0 by $\eta(X|\{\mathcal{F}_k\}_{k=0}^n)$.

The basic properties of a mild value measure and a homogeneous filtered value measure are shown in [16].

1.2. Main theorem

Let $\mathcal{Z} = \{z_1, z_2, \dots, z_N\} \subset \mathbf{R}^M$, equipped with the discrete topology, and $\mathcal{B}(\mathcal{Z})$ the Borel algebra with respect to this. We consider a probability \hat{P} on $(\mathcal{Z}, \mathcal{B}(\mathcal{Z}))$ define by $\hat{P}[\{z_j\}] = p_j$, $j = 1, 2, \dots, N$, where $p_j > 0$, $j = 1, 2, \dots, N$ and $\sum_{j=1}^N p_j = 1$. Also we consider M -dimensional random variables \hat{Z} and $\hat{Y}^{(n)}$, $n \in \mathbf{N}$ on $(\mathcal{Z}, \mathcal{B}(\mathcal{Z}))$ defined by

$$\begin{aligned}
 (4) \quad \hat{Z}(z) &= (\hat{Z}_1(z), \hat{Z}_2(z), \dots, \hat{Z}_M(z)) = z, \quad z \in \mathcal{Z}, \\
 \hat{Y}^{(n)}(z) &= (\hat{Y}_1^{(n)}(z), \hat{Y}_2^{(n)}(z), \dots, \hat{Y}_M^{(n)}(z)), \\
 (5) \quad \hat{Y}_i^{(n)}(z) &= \exp(\hat{Z}_i(z) \sqrt{\frac{T}{n}} + b_i \frac{T}{n}), \quad i = 1, 2, \dots, M, \quad n \in \mathbf{N},
 \end{aligned}$$

where b_i , $i = 1, 2, \dots, M$ and T are positive numbers. We denote by $\hat{\mathcal{P}}$ the set of all probability measures on $(\mathcal{Z}, \mathcal{B}(\mathcal{Z}))$. We identify $\hat{\mathcal{P}}$ as the hyperplane $\{(q_1, q_2, \dots, q_N) \mid q_j \geq 0, j = 1, 2, \dots, N, \sum_{j=1}^N q_j = 1\}$ on \mathbf{R}^N .

Let (Ω, \mathcal{F}, P) be the direct product probability space of countably infinite copies of $(\mathcal{Z}, \mathcal{B}(\mathcal{Z}), \hat{P})$. We consider M -dimensional random variables

Z_k and $Y_k^{(n)} = (Y_{1,k}^{(n)}, \dots, Y_{M,k}^{(n)})$, $k, n \in \mathbf{N}$ on (Ω, \mathcal{F}) defined by

$$(6) \quad Z_k(\omega) = \hat{Z}(\omega_k),$$

$$(7) \quad Y_k^{(n)}(\omega) = \hat{Y}^{(n)}(\omega_k),$$

where $\omega = (\omega_1, \omega_2, \dots) \in \Omega$. We give a filtration $\{\mathcal{F}_k\}_{k=0,1,2,\dots}$ on (Ω, \mathcal{F}) by

$$(8) \quad \begin{aligned} \mathcal{F}_0 &= \sigma(\emptyset, \Omega), \\ \mathcal{F}_k &= \sigma(Z_1, Z_2, \dots, Z_k), \quad k = 1, 2, \dots \end{aligned}$$

Note that the process $(Y_k^{(n)})$ is $\{\mathcal{F}_k\}$ -adapted, and $Y_k^{(n)}$, $k = 1, 2, \dots$ is independent of \mathcal{F}_l , $l = 1, 2, \dots, k - 1$ for each $n \in \mathbf{N}$. We denote by \mathcal{P} the set of all probability measures which is absolutely continuous with respect to P .

We consider an $\{\mathcal{F}_k\}$ -adapted M -dimensional process $\{S_k^{(n)}\}_{k=1,2,\dots,n}$ for each $n \in \mathbf{N}$ defined by

$$(9) \quad \begin{aligned} S_k^{(n)} &= (S_{1,k}^{(n)}, S_{2,k}^{(n)}, \dots, S_{M,k}^{(n)}), \quad k = 1, 2, \dots, n \\ S_{i,k}^{(n)} &= S_{i,0} \prod_{l=1}^k Y_{i,l}^{(n)}, \quad i = 1, 2, \dots, M. \end{aligned}$$

where $S_0 = (S_{1,0}, S_{2,0}, \dots, S_{M,0}) \in (0, \infty)^M$ is a constant vector. $S_k^{(n)}$ is interpreted as the price vector of M risky assets at time k . We call a $\{\mathcal{F}_k\}$ -predictable M dimensional process $\xi = (\xi_k)_{k=1,2,\dots}$ a self-financing strategy, and denote by \mathcal{SF} the set of all self-financing strategy. Then we define a random variable $V_k^{(n)}(v, \xi)$ for $v \geq 0$, and $\xi \in \mathcal{SF}$ by $V_0^{(n)}(v, \xi) = v$ and

$$(10) \quad V_k^{(n)}(v, \xi) = v + \sum_{l=1}^k \xi_l \cdot (S_l^{(n)} - S_{l-1}^{(n)}).$$

$V_k^{(n)}(v, \xi)$ represents the discount value of self-financing portfolio (v, ξ) at time k .

Hereafter we use the following type of mild value measure:

$$(11) \quad \eta(\nu) = \int_0^1 \eta_\alpha(\nu) \mu(d\alpha), \quad \nu \in \mathcal{L}^\infty, \mu \in \mathcal{M},$$

and fix it. We define $\hat{\mathcal{Q}}$ by

$$\hat{\mathcal{Q}} = \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{X}] \geq 0, \text{ for all random variables } \hat{X} \text{ on } (\mathcal{Z}, \mathcal{B}(\mathcal{Z})) \text{ such that } \eta(\nu_{\hat{X}}) \geq 0 \text{ holds.}\},$$

where $\nu_{\hat{X}}$ is the probability distribution on \hat{X} .

Let $\hat{C}([0, \infty)^M : \mathbf{R})$ be the set of functions $g : [0, \infty)^M \rightarrow \mathbf{R}$ such that

$$(12) \quad \begin{aligned} |g(x) - g(x')| &\leq K|x - x'|, \quad x, x' \in [0, \infty)^M, \\ |g(x)| &\leq K(1 + |x|^{2m}), \quad x \in [0, \infty)^M, \end{aligned}$$

hold for some $K > 0, m \in \mathbf{N}$. Take $f \in \hat{C}([0, \infty)^M : \mathbf{R})$ and fix it.

We assume the following in what follows.

ASSUMPTION 1.3. $M + 1 < N$.

REMARK. This assumption indicates that the market model which we consider here is incomplete.

ASSUMPTION 1.4. *the set $\hat{\mathcal{Q}} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Z}_i] = 0\}$ contains at least one inner point, where we consider the relative topology of the usual topology on \mathbf{R}^N to the hyperplane $\hat{\mathcal{P}}$.*

Let us define $\Gamma \in \mathbf{R}^{M \times M}$ by

$$\Gamma = \{\gamma = (\gamma_{ii'})_{i,i'=1,2,\dots,M} \mid \gamma_{ii'} = E^{\hat{Q}}[\hat{Z}_i \hat{Z}_{i'}], \text{ for some } \hat{Q} \in \hat{\mathcal{Q}} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Z}_i] = 0\}\}.$$

Note that $\gamma \in \Gamma$ is nonnegative definite. Also we can easily see that Γ is compact with respect to the usual topology on $\mathbf{R}^{M \times M}$.

Our main theorem is the following.

THEOREM 1.5. *We have $\lim_{n \rightarrow \infty} \sup_{\xi \in \mathcal{SF}} \eta(V_n^{(n)}(v, \xi) + f(S_n^{(n)})) = v + U(0, S_0)$, where $U : [0, T] \times [0, \infty)^M \rightarrow \mathbf{R}$ is the unique viscosity solution of the following Hamilton-Jacobi-Bellman equation:*

$$(13) \quad \begin{aligned} \frac{\partial U}{\partial t} + \inf_{\gamma=(\gamma_{ii'})_{i,i' \in \Gamma}} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{ii'} x_i x_{i'} \frac{\partial^2 U}{\partial x_i \partial x_{i'}} &= 0, \\ U(T, x) &= f(x), \quad x \in [0, \infty)^M, \end{aligned}$$

satisfying $U(t, \cdot) \in \hat{C}([0, \infty)^M : \mathbf{R})$ for each $t \in [0, T]$.

We give a proof of Theorem 1.5 in Section 4.

Example 1.6. We consider the case where $M = 1, N = 3$, and $\eta = \eta_\alpha$, $\alpha \in (0, 1)$. We also assume that $E^{\hat{P}}[\hat{Z}] = \sum_{j=1}^3 p_j z_j = 0$. In this case, we have $\hat{Q} = \{\hat{Q} \in \hat{P} \mid \hat{q}_j = \hat{Q}[\hat{Z} = z_j] \leq p_j/\alpha, j = 1, 2, 3\}$.

We see that the condition “ $\hat{Q} \in \hat{Q} \cap \{\hat{Q} \in \hat{P} \mid E^{\hat{Q}}[\hat{Z}] = 0\}$ ” is equivalent to the following:

$$\hat{q}_1 \in [\underline{q}, \bar{q}], \quad \hat{q}_2 = -\frac{z_3}{z_2 - z_3} - \frac{z_1 - z_3}{z_2 - z_3} \hat{q}_1, \quad \hat{q}_3 = \frac{z_2}{z_2 - z_3} + \frac{z_1 - z_2}{z_2 - z_3} \hat{q}_1,$$

where

$$\underline{q} = 0 \vee \frac{-\alpha z_3 - p_2(z_2 - z_3)}{\alpha(z_1 - z_3)} \vee \frac{-z_2}{z_1 - z_2},$$

$$\bar{q} = \frac{p_1}{\alpha} \wedge \frac{-z_3}{z_1 - z_3} \wedge \frac{-\alpha z_2 + p_3(z_2 - z_3)}{\alpha(z_1 - z_2)}.$$

Also we have $\Gamma = [\underline{q}(z_1 - z_2)(z_1 - z_3) - z_2 z_3, \bar{q}(z_1 - z_2)(z_1 - z_3) - z_2 z_3]$. We can easily see that $\underline{q} < \bar{q}$. Hence Assumption 1.4 holds.

2. Discrete model

Throughout this paper, we use the following notation:

$$(14) \quad xy = (x_1 y_1, x_2 y_2, \dots, x_M y_M) \in \mathbf{R}^M,$$

for $x = (x_1, x_2, \dots, x_M), y = (y_1, y_2, \dots, y_M) \in \mathbf{R}^M$.

We consider some maximization problem on hedging with a homogeneous filtered value measure in a discrete time market model. The following setting is parallel to that of Section 1. Let (Ω, \mathcal{F}, P) be a standard probability space with a filtration $\{\mathcal{F}_k\}_{k=0,1,\dots,n}$. Also, let be $Y_k = (Y_{1,k}, \dots, Y_{M,k}), k = 1, 2, \dots, n$ be identically distributed M -dimensional random variables such that Y_k is $\{\mathcal{F}_k\}$ -measurable and independent of $\mathcal{F}_l, l = 1, 2, \dots, k - 1$. We assume that $Y_{i,k}(\omega) > 0, \omega \in \Omega, \underline{Y} = \min_{i=1,2,\dots,M} \text{ess.inf}_\omega Y_{i,1}(\omega) > 0$, and $\bar{Y} = \max_{i=1,2,\dots,M} \text{ess.sup}_\omega Y_{i,1}(\omega) < \infty$. We denote by \mathcal{P} the set of all probability measures on (Ω, \mathcal{F}) which are absolutely continuous with respect to P .

We define M dimensional $\{\mathcal{F}_k\}$ -adapted process $\{S_k\}_{k=1,2,\dots,n}$ by

$$\begin{aligned}
 S_k &= (S_{1,k}, S_{2,k}, \dots, S_{M,k}), \\
 (15) \quad S_{i,k} &= S_{i,0} \prod_{l=1}^k Y_{i,l}, \quad i = 1, 2, \dots, M,
 \end{aligned}$$

where $S_0 = (S_{1,0}, S_{2,0}, \dots, S_{M,0}) \in (0, \infty)^M$ is a constant vector.

We call an M dimensional $\{\mathcal{F}_k\}$ -predictable process $\xi = (\xi_k)_{k=1,2,\dots}$ a self-financing strategy, and denote by \mathcal{SF} the set of all self-financing strategies. Then we define a random variable $V_k(v, \xi)$ for $v \in (0, \infty)$, $\xi \in \mathcal{SF}$ by

$$V_0(v, \xi) = v \text{ and } V_k(v, \xi) = v + \sum_{l=1}^k \xi_l \cdot (S_l - S_{l-1}), \quad k \in \mathbf{N}.$$

As in section 1, we use the following type of mild value measure:

$$(16) \quad \eta(\nu) = \int_0^1 \eta_\alpha(\nu) \mu(d\alpha), \quad \nu \in \mathcal{L}^\infty, \quad \mu \in \mathcal{M},$$

and fix it. Let $\Phi_\mu : [0, 1] \rightarrow [0, 1]$, $\mu \in \mathcal{M}$ be a mapping defined by

$$\begin{aligned}
 (17) \quad \Phi_\mu(x) &= \int_{[1-x, 1]} \left(1 - \frac{1-x}{\alpha}\right) \mu(d\alpha), \quad x \in [0, 1], \\
 \Phi_\mu(1) &= 1.
 \end{aligned}$$

We can easily see that $0 \leq \Phi_\mu(x) \leq x$, since $\{1 - (1-x/\alpha)\} \leq \{1 - (1-x)\} = x$ holds for any $\alpha \in (0, 1]$. We also define

$$\begin{aligned}
 (18) \quad \mathcal{Q} &= \{Q \in \mathcal{P} \mid E^Q[X] \geq 0 \text{ for all } X \in L^\infty(\Omega, \mathcal{F}_1, P) \\
 &\quad \text{such that } \eta(\nu_X) \geq 0 \text{ hold}\}.
 \end{aligned}$$

We assume the following.

ASSUMPTION 2.1. $\mathcal{Q} \cap \bigcap_{i=1}^M \{Q \in \mathcal{P} \mid E^Q[Y_{i,1}] = 1\} \neq \emptyset$.

Let us denote $\eta(\nu_X)$ by $\bar{\eta}(X)$ for $X \in L^\infty(\Omega, \mathcal{F}, P)$, where ν_X is the distribution of X . Obviously we have $\bar{\eta}(X) = \inf_{Q \in \mathcal{Q}} E^Q[X]$, $X \in L^\infty(\Omega, \mathcal{F}, P)$. Also we see that $\mathcal{Q} = \{Q \in \mathcal{P} \mid Q[A] \geq \Phi_\mu(P[A]), A \in \mathcal{F}\}$ by Theorem 4.73 in [15]. We denote

$$(19) \quad \psi_g(x, y) = \bar{\eta}((xy) \cdot (Y_1 - \mathbf{1}) + g(xY_1)), \quad \mathbf{1} = (1, 1, \dots, 1) \in \mathbf{R}^M,$$

for $x \in [0, \infty)^M$, $y \in \mathbf{R}^M$, and $g \in \hat{C}([0, \infty)^M; \mathbf{R})$. Since

$$\begin{aligned} & |\{(xy) \cdot (Y_1 - \mathbf{1}) + g(xY_1)\} - \{(x'y') \cdot (Y_1 - \mathbf{1}) + g(x'Y_1)\}| \\ & \leq K\{(\bar{Y} + 1)|xy - x'y'| + \bar{Y}|x - x'|\}, \end{aligned}$$

$x, x' \in [0, \infty)^M$, $y, y' \in \mathbf{R}^M$, for some $K > 0$, and $\bar{\eta}$ is monotone, i.e., $\bar{\eta}(X) \leq \bar{\eta}(X')$, for $X, X' \in L^\infty(\Omega)$ such that $X \leq X'$, then we have

$$|\psi_g(x, y) - \psi_g(x', y')| \leq K\{(\bar{Y} + 1)|xy - x'y'| + \bar{Y}|x - x'|\}.$$

Hence $(x, y) \mapsto \psi_g(x, y)$ is continuous. Next we define

$$(20) \quad \phi_g(x) = \sup_{y \in \mathbf{R}^M} \psi_g(x, y).$$

Note that $\phi_g(x) < +\infty$, $x \in [0, \infty)^M$. Indeed, there exists $\bar{Q} \in \mathcal{Q} \cap \bigcap_{i=1}^M \{Q \in \mathcal{P} \mid E^Q[Y_{i,1}] = 1\}$ by Assumption 2.1, and then we see that $\phi_g(x) \leq E^{\bar{Q}}[g(xY_1)] \leq K(1 + |x|^{2m})$, for some $K > 0$. Also we have

$$\begin{aligned} \phi_g(x) &= \sup_{y \in \mathbf{R}^M} \inf_{Q \in \mathcal{Q}} E^Q[(xy) \cdot (Y_1 - \mathbf{1}) + g(xY_1)] \\ &= \inf_{Q \in \mathcal{Q}} \sup_{y \in \mathbf{R}^M} E^Q[(xy) \cdot (Y_1 - \mathbf{1}) + g(xY_1)] \\ &= \inf_{Q \in \mathcal{Q} \cap \bigcap_{i=1}^M \{Q \in \mathcal{P} \mid E^Q[Y_{i,1}] = 1\}} E^Q[g(xY_1)], \end{aligned}$$

by Takahashi's Minimax Theorem in [20]. Since $|g(xY_1) - g(x'Y_1)| \leq K\bar{Y}|x - x'|$ and $|g(xY_1)| \leq K(1 + \bar{Y}^{2m}|x|^{2m})$ holds for $x, x' \in [0, \infty)^M$ and $g \in \hat{C}([0, \infty)^M; \mathbf{R})$, where $K > 0$ is a constant which is independent of x, x', Q , the mapping $\phi_g : [0, \infty)^M \rightarrow \mathbf{R}$ belongs to $\hat{C}([0, \infty)^M; \mathbf{R})$. Then we can define an operator H on $\hat{C}([0, \infty)^M; \mathbf{R})$ by $Hg = \phi_g$. Also we inductively define $H^k g = H(H^{k-1}g)$, $k = 1, 2, \dots$ and $H^0 g = g$ for $g \in \hat{C}([0, \infty)^M; \mathbf{R})$.

Now we fix $n \in \mathbf{N}$. We define random variables $L_k(v, \xi)$, $k = 0, 1, \dots, n$, $v \in (0, \infty)$, and $\xi \in \mathcal{SF}$ inductively by $L_n(v, \xi) = V_n(v, \xi) + f(S_n)$ and

$$(21) \quad L_{k-1}(v, \xi) = \eta(L_k(v, \xi) | \mathcal{F}_{k-1}), \quad k = n, n-1, \dots, 1,$$

where $f \in \hat{C}([0, \infty)^M; \mathbf{R})$. Obviously $L_k(v, \xi)$, $k = 0, 1, \dots, n$ equals the homogeneous value measure for $V_n(v, \xi) + f(S_n)$ at k .

We prove the following theorem, which is our main result in this section.

THEOREM 2.2 (Bellman's Principle). *We have*

$$\begin{aligned} & \sup_{\xi' \in \mathcal{SF}} L_k(v, (\xi_1, \xi_2, \dots, \xi_k, \xi'_{k+1}, \xi'_{k+2}, \dots, \xi'_n)) \\ & = V_k(v, \xi) + H^{n-k} f(S_k), \quad k = 0, 1, \dots, n, \end{aligned}$$

for any $v \in (0, \infty)$ and $\xi \in \mathcal{SF}$.

We show a lemma to prove Theorem 2.2.

LEMMA 2.3. *For any $g \in \hat{C}([0, \infty)^M : \mathbf{R})$ and $\varepsilon > 0$, there exists a Borel measurable function $\gamma^\varepsilon(x) = \gamma_g^\varepsilon(x) = (\gamma_1^\varepsilon(x), \gamma_2^\varepsilon(x), \dots, \gamma_M^\varepsilon(x))$ on $[0, \infty)^M$ such that $Hg(x) - \varepsilon \leq \bar{\eta}((x\gamma^\varepsilon(x)) \cdot (Y_1 - \mathbf{1}) + g(xY_1))$, $x \in [0, \infty)^M$ holds.*

PROOF. First we define a multivalued mapping $\Gamma^\varepsilon : ([0, \infty)^M, \mathcal{B}[0, \infty)^M) \Rightarrow (\mathbf{R}^M, \mathcal{B}(\mathbf{R}^M))$ by $\Gamma^\varepsilon(x) = \{y \in \mathbf{R}^M \mid Hg(x) - \varepsilon \leq \psi_g(x, y)\}$. Obviously the set on the right side is nonempty, so it is sufficient to show that this multivalued mapping is measurable, i.e.,

$$(22) \quad \Gamma^{-w, \varepsilon}(A) = \{x \in [0, \infty)^M \mid \Gamma^\varepsilon(x) \cap A \neq \emptyset\} \in \mathcal{B}[0, \infty)^M,$$

for any closed set $A \subset \mathbf{R}^M$. If Γ^ε is measurable, there exists a measurable selection $\gamma^\varepsilon(x) \in \Gamma^\varepsilon(x)$ and this mapping satisfies the condition. Also, we may assume that A is compact. Indeed, if $\Gamma^{-w, \varepsilon}(A') \in \mathcal{B}[0, \infty)^M$ for any compact set A' , we see that $\Gamma^{-w, \varepsilon}(A) = \bigcup_{m=1}^{\infty} \Gamma^{-w, \varepsilon}(A \cap [-m, m]^M) \in \mathcal{B}[0, \infty)^M$.

We show that $\Gamma^{-w, \varepsilon}(A)$ is closed for any compact set A . Take a sequence $(x_m)_{m \in \mathbf{N}}$ of $\Gamma^{-w, \varepsilon}(A)$ such that $\lim_{m \rightarrow \infty} x_m = x \in [0, \infty)^M$. Then there exists a $y_m \in \Gamma^\varepsilon(x_m) \cap A$ for each m . Since A is compact, we can choose a subsequence $(y_{m(l)})_{l \in \mathbf{N}}$ of $(y_m)_{m \in \mathbf{N}}$ such that $y_{m(l)}$ converges to some $y \in A$ as $l \rightarrow \infty$. Taking $\lim_{l \rightarrow \infty}$ in both sides of the equation $Hg(x_{m(l)}) - \varepsilon \leq \psi_g(x_{m(l)}, y_{m(l)})$, we have $Hg(x) - \varepsilon \leq \psi_g(x, y)$ because ψ_g and Hg are continuous. This implies that $y \in \Gamma^\varepsilon(x) \cap A$ and $x \in \Gamma^{-w, \varepsilon}(A)$. \square

Now we give the proof of Theorem 2.2. First we show that

$$L_k(v, \xi) \leq V_k(v, \xi) + H^{n-k} f(S_k),$$

by mathematical induction on k . Obviously, the claim holds when $k = n$. Suppose that the claim holds for some k , Then we have

$$\begin{aligned} L_{k-1}(v, \xi) &= \eta(L_k(v, \xi) | \mathcal{F}_{k-1}) \\ &\leq \eta(V_k(v, \xi) + H^{n-k} f(S_k) | \mathcal{F}_{k-1}) \\ &= V_{k-1}(v, \xi) + \eta((S_{k-1} \xi_k) \cdot (Y_k - \mathbf{1}) + H^{n-k} f(S_{k-1} Y_k) | \mathcal{F}_{k-1}) \\ &\leq V_{k-1}(v, \xi) + H^{n-(k-1)} f(S_{k-1}). \end{aligned}$$

Hence we have the claim.

Next we show that there exists $\xi^\varepsilon \in \mathcal{SF}$ for $\varepsilon > 0$ such that

$$\begin{aligned} &V_k(v, \xi) + H^{n-k} f(S_k) - \frac{k\varepsilon}{n} \\ &\leq L_k(v, (\xi_1, \xi_2, \dots, \xi_k, \xi_{k+1}^\varepsilon, \xi_{k+2}^\varepsilon, \dots, \xi_n^\varepsilon)), \quad k = n, n-1, \dots, 1 \end{aligned}$$

for any $\xi \in \mathcal{SF}$. Applying Lemma 2.3 for $g = f$, we see that there exists some Borel measurable function γ^ε on $[0, \infty)^M$ such that

$$(Hf)(x) - \frac{\varepsilon}{n} \leq \bar{\eta}((x\gamma^\varepsilon(x)) \cdot (Y_1 - \mathbf{1}) + f(xY_1)),$$

for $x \in [0, \infty)^M$. Then, $\xi_n^\varepsilon = \gamma^\varepsilon(S_{n-1})$ is an $\{\mathcal{F}_{n-1}\}$ -measurable random variable such that

$$\begin{aligned} &V_{n-1}(v, \xi) + (Hf)(S_{n-1}) - \frac{\varepsilon}{n} \\ &\leq V_{n-1}(v, \xi) + \bar{\eta}((x\gamma^\varepsilon(x)) \cdot (Y_1 - \mathbf{1}) + f(xY_1)) |_{x=S_{n-1}} \\ &\leq V_{n-1}(v, \xi) + \eta((S_{n-1} \xi_n^\varepsilon) \cdot (Y_n - \mathbf{1}) + f(S_{n-1} Y_n) | \mathcal{F}_{n-1}) \\ &\leq \eta(V_n(v, \xi) + f(S_n) | \mathcal{F}_{n-1}) \\ &\leq L_{n-1}(v, \xi_1, \xi_2, \dots, \xi_{n-1}, \xi_n^\varepsilon). \end{aligned}$$

Using induction, we can construct ξ^ε such that the condition holds by the same way. This completes the proof of Theorem 2.2.

3. Proof of Theorem 1.5

3.1. Preparations

Let (Ω, \mathcal{F}, P) be the probability space defined in Section 1.2. For $k \in \mathbf{N}$, we denote by \mathcal{L}_k the set of measurable mappings $\rho : \Omega \times \mathcal{Z} \rightarrow \mathbf{R}$ which satisfy the following:

$$\begin{aligned} &\rho(\cdot, z) \text{ is } \{\mathcal{F}_{k-1}\} \text{ measurable for each } z \in \mathcal{Z}, \\ &\sum_{j=1}^N \rho(\omega, z_j) p_j = 1, \quad \omega \in \Omega, \\ &\sum_{j \in J} \rho(\cdot, z_j) p_j \geq \Phi_\mu\left(\sum_{j \in J} p_j\right), \quad J \subset \{1, 2, \dots, M\}, \quad z_j \in \mathcal{Z}, \quad j \in J. \end{aligned}$$

Also we denote by $\mathcal{Q}^{(n)}$, $n \in \mathbf{N}$ the set of probability measures $Q \in \mathcal{P}$ which satisfy the following:

$$\begin{aligned} E\left[\frac{dQ}{dP} \mid \mathcal{F}_n\right] &= \prod_{k=1}^n \rho_k(\cdot, Z_k), \quad \rho_k \in \mathcal{L}_k, \\ \int_{\Omega} \rho_k(\omega', Z_k(\omega)) Y_{i,k}^{(n)}(\omega) P(d\omega) &= 1, \quad \omega' \in \Omega, \quad k = 1, 2, \dots, n. \end{aligned}$$

Let us denote $X_{i,k}^{(n)} = \prod_{l=1}^k Y_{i,l}^{(n)}$ for $n \in \mathbf{N}$, $i = 1, 2, \dots, M$, and $k = 1, 2, \dots, n$. We also denote $X_k^{(n)} = (X_{1,k}^{(n)}, X_{2,k}^{(n)} \dots X_{M,k}^{(n)})$, $k = 1, 2, \dots, n$. We define $a_{ij}^{(n)} = \exp(z_{ij} \sqrt{T/n} + b_i T/n)$, $i = 1, 2, \dots, M$, $j = 1, 2, \dots, N$, where $z_j = (z_{1j}, z_{2j}, \dots, z_{Mj}) \in \mathcal{Z}$.

Our purpose in this subsection is to prove the following.

LEMMA 3.1.

(1) $(H^{(n)})^k g(x) = \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(x X_k^{(n)})]$, $n \in \mathbf{N}$, $g \in \hat{C}([0, \infty)^M : \mathbf{R})$,

where $H^{(n)}$, $n \in \mathbf{N}$ are operators on $\hat{C}([0, \infty)^M : \mathbf{R})$ that correspond to H in Section 3.

(2) $\sup_{n \in \mathbf{N}} \sup_{Q \in \mathcal{Q}^{(n)}} E^Q[\max_{k=1,2,\dots,n} |X_k^{(n)}|^{2m}] < \infty$, $m \in \mathbf{N}$.

(3) There exists a positive number L , which only depends on M , such that

$E^Q[|X_{k+l}^{(n)} - X_k^{(n)}|^4] \leq L(lT/n)^2$ holds for any $k, l = 0, 1, \dots, n$, $k + l \leq n$, and $Q \in \mathcal{Q}^{(n)}$.

PROOF. We show assertion (1). We see by Theorem 4.73 in [15] that $\bar{\eta}(X) = \inf_{Q \in \bar{\mathcal{Q}}_1^{(n)}} E^Q[X]$, for $X \in L^\infty(\Omega, \mathcal{F}_1, P)$, where

$$(23) \quad \begin{aligned} \bar{\mathcal{Q}}_1^{(n)} &= \{Q \in \mathcal{P} \mid Q[A] \geq \Phi_\mu(P[A]), A \in \mathcal{F}_1\} \\ &= \{Q \in \mathcal{P} \mid E[\frac{dQ}{dP}|\mathcal{F}_1] = \rho_1(\cdot, Z_1) \in \mathcal{L}_1\} \end{aligned}$$

Then we have $H^{(n)}g(x) = \inf_{Q \in \mathcal{Q}_1^{(n)}} E^Q[g(xY_1^{(n)})]$, where $\mathcal{Q}_1^{(n)} = \bar{\mathcal{Q}}_1^{(n)} \cap \bigcap_{i=1}^M \{Q \in \mathcal{P} \mid E[\rho_1(\omega', Z_1)Y_{i,1}^{(n)}] = 1, \omega' \in \Omega\}$, by a way similar to that in Section 2. We can easily see that $\mathcal{Q}^{(n)} \subset \mathcal{Q}_1^{(n)}$. We show the inverse implement. Take $Q \in \mathcal{Q}_1^{(n)}$ and define $\tilde{Q} \in \mathcal{P}^{(n)}$ by $E[d\tilde{Q}/dP|\mathcal{F}_n] = \prod_{k=1}^n \tilde{\rho}_k(\cdot, Z_k)$, where $\tilde{\rho}_k(\omega, z) = E[\frac{dQ}{dP}|\mathcal{F}_1](\omega)$, $\omega \in \Omega$, $z \in \mathcal{Z}$, $k = 1, 2, \dots, n$. Then we see that $\tilde{Q} \in \mathcal{Q}^{(n)}$ and $E^Q[g(xY_1^{(n)})] = E^{\tilde{Q}}[g(xY_1^{(n)})]$. This implies that $H^{(n)}g(x) = \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(xY_1^{(n)})]$.

Suppose that $(H^{(n)})^l g(x) = \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(xX_l^{(n)})]$, $l \leq k$, $g \in \hat{C}([0, \infty)^M)$ holds for some $k \in \{1, 2, \dots, n-1\}$. First we show that

$$(24) \quad (H^{(n)})^{k+1} g(x) \leq \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(xX_{k+1}^{(n)})].$$

Take $Q \in \mathcal{Q}^{(n)}$ and $\rho_l \in \mathcal{L}_l$, $l \leq n$ such that $E[dQ/dP|\mathcal{F}_n] = \prod_{l=1}^n \rho_l(\cdot, Z_l)$ holds. We define $Q^{\tilde{\omega}} \in \mathcal{P}$ for each $\tilde{\omega} \in \Omega$ by $E[dQ^{\tilde{\omega}}/dP|\mathcal{F}_n] = \prod_{l=1}^n \rho'_l(\cdot, Z_l)$, where $\rho'_l(\omega, z) = \rho'_{k+1}(\tilde{\omega}, z)$, $\omega \in \Omega$, $z \in \mathcal{Z}$. Then we see that $Q^{\tilde{\omega}} \in \mathcal{Q}^{(n)}$ and

$$\begin{aligned} E^Q[g(xX_{k+1}^{(n)})|\mathcal{F}_k](\tilde{\omega}) &= E^Q[g(yY_{k+1}^{(n)})|\mathcal{F}_k](\tilde{\omega})|_{y=xX_k^{(n)}} \\ &= E^{Q^{\tilde{\omega}}}[g(yY_1^{(n)})]|_{y=xX_k^{(n)}} \geq H^{(n)}g(xX_k^{(n)}). \end{aligned}$$

Also we have

$$\begin{aligned} E^Q[g(xX_{k+1}^{(n)})] &= E^Q[E^Q[g(xX_{k+1}^{(n)})|\mathcal{F}_k]] \\ &\geq E^Q[H^{(n)}g(xX_k^{(n)})] \geq (H^{(n)})^k H^{(n)}g(x) = (H^{(n)})^{k+1} g(x). \end{aligned}$$

This implies that (24) holds.

Next we show that

$$(25) \quad (H^{(n)})^{k+1}g(x) \geq \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(xX_{k+1}^{(n)})].$$

Take $\varepsilon > 0$. Then we see by assumption that there exist $\bar{Q}^0, \bar{Q}^j \in \mathcal{Q}^{(n)}$ $j = 1, 2, \dots, N$ such that

$$(26) \quad \begin{aligned} (H^{(n)})^{k+1}g(x) &\geq E^{\bar{Q}^0}[(H^{(n)})^k g(xY_1^{(n)})] - \frac{\varepsilon}{2}, \\ (H^{(n)})^k g(xa_j^{(n)}) &\geq E^{\bar{Q}^j}[g(xa_j^{(n)}X_k^{(n)})] - \frac{\varepsilon}{2}. \end{aligned}$$

Take $\rho_l^0, \rho_l^j \in \mathcal{L}_l$, $l = 1, 2, \dots, n$, $j = 1, 2, \dots, M$, such that $E[\frac{d\bar{Q}^0}{dP}|\mathcal{F}_n] = \prod_{l=1}^n \rho_l^0(\cdot, Z_l)$, $E[\frac{d\bar{Q}^j}{dP}|\mathcal{F}_n] = \prod_{l=1}^n \bar{\rho}_l^j(\cdot, Z_l)$ hold. Then we define $\bar{Q} \in \mathcal{P}$ by $E[\frac{d\bar{Q}}{dP}|\mathcal{F}_n] = \prod_{l=1}^n \bar{\rho}_k(\cdot, Z_k)$, where

$$(27) \quad \begin{aligned} \bar{\rho}_1 &= \bar{\rho}_1^0, \\ \bar{\rho}_l &= \sum_{j=1}^N \bar{\rho}_{l-1}^j 1_{\{Z_1=z_j\}}, \quad l = 2, 3, \dots, n. \end{aligned}$$

We see that $\bar{Q} \in \mathcal{Q}^{(n)}$ and

$$(28) \quad (H^{(n)})^{k+1}g(x) \geq E^{\bar{Q}}[g(xX_{k+1}^{(n)})] - \varepsilon \geq \inf_{Q \in \mathcal{Q}^{(n)}} E^Q[g(xX_{k+1}^{(n)})] - \varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, we have (25). This shows the assertion (1).

Next we show the assertion (2). Since $X_{i,n}^{(n)}$ is martingale under $Q \in \mathcal{Q}^{(n)}$, it is sufficient to show that $\sup_{n \in \mathbb{N}} \sup_{Q \in \mathcal{Q}^{(n)}} E^Q[|X_{i,n}^{(n)}|^{2m}] < \infty$, for each $i = 1, 2, \dots, M$. Take $Q \in \mathcal{Q}^{(n)}$ and fix $\omega \in \Omega$. We denote $q_{\omega,k,j} = Q[Z_k = z_j | \mathcal{F}_{k-1}](\omega)$, $k = 1, 2, \dots, n$, $j = 1, 2, \dots, N$. We see that

$$1 = E^Q[Y_{i,k}^{(n)} | \mathcal{F}_{k-1}](\omega) = \sum_{j=1}^N q_{\omega,k,j} (a_{il}^{(n)})$$

$$\begin{aligned}
 &= 1 + E^Q[Z_{i,k}|\mathcal{F}_{k-1}](\omega)\sqrt{\frac{T}{n}} + b_i\frac{T}{n} \\
 &\quad + \sum_{j=1}^N q_{\omega,k,j}\{\exp(z_{ij}\sqrt{\frac{T}{n}} + b_i\frac{T}{n}) - (1 + z_{ij}\sqrt{\frac{T}{n}} + b_i\frac{T}{n})\}.
 \end{aligned}$$

Then we have

$$\begin{aligned}
 &E^Q[|Y_{i,k}^{(n)}|^{2m}|\mathcal{F}_{k-1}](\omega) \\
 &= 1 + 2mE^Q[Z_{i,k}|\mathcal{F}_{k-1}](\omega)\sqrt{\frac{T}{n}} + 2mb_i\frac{T}{n} \\
 &\quad + \sum_{j=1}^N q_{\omega,k,j}\{\exp(2mz_{ij}\sqrt{\frac{T}{n}} + 2mb_i\frac{T}{n}) - (1 + 2mz_{ij}\sqrt{\frac{T}{n}} + 2mb_i\frac{T}{n})\},
 \end{aligned}$$

and

$$\begin{aligned}
 &E^Q[|Y_{i,k}^{(n)}|^{2m}|\mathcal{F}_{k-1}](\omega) \\
 &\leq 1 + \sum_{j=1}^N \{2m|\exp(z_{ij}\sqrt{\frac{T}{n}} + b_i\frac{T}{n}) - (1 + z_{ij}\sqrt{\frac{T}{n}} + b_i\frac{T}{n})| \\
 &\quad + |\exp(2mz_{ij}\sqrt{\frac{T}{n}} + 2mb_i\frac{T}{n}) - (1 + 2mz_{ij}\sqrt{\frac{T}{n}} + 2mb_i\frac{T}{n})|\}.
 \end{aligned}$$

for $k = 1, 2, \dots, n$. We denote by $b^{(n)}$ the right term of this inequality. Note that $b^{(n)}$ is independent of k, ω and Q . We see that $\lim_{n \rightarrow \infty} n(b^{(n)} - 1)$ exists and $|n(b^{(n)} - 1)| \leq b, n \in \mathbf{N}$, for some $b > 0$. Then we have

$$\begin{aligned}
 E^Q[|X_{i,n}^{(n)}|^{2m}] &\leq E^Q[|X_{i,n-1}^{(n)}|^{2m} E^Q[|Y_{i,n}^{(n)}|^{2m}|\mathcal{F}_{n-1}]] \\
 &\leq (1 + \frac{b}{n})E^Q[|X_{i,n-1}^{(n)}|^{2m}] \leq \dots \leq (1 + \frac{b}{n})^n, \quad Q \in \mathcal{Q}^{(n)}.
 \end{aligned}$$

This implies that $\limsup_{n \rightarrow \infty} \sup_{Q \in \mathcal{Q}^{(n)}} E^Q[|X_{i,n}^{(n)}|^{2m}] \leq e^b$. Hence we have the assertion.

We show that assertion (3) holds. We see that there exists some $c > 0$ such that $E^Q[|Y_{i,k+1}^{(n)} - 1|^4|\mathcal{F}_k](\omega) \leq c(\frac{T}{n})^2$, for each $i \leq M, k \leq n - 1, Q$ and ω by an argument similar to that of (2). Then using the result of (2), we have $E^Q[|X_{i,k+1}^{(n)} - X_{i,k}^{(n)}|^4] \leq c(T/n)^2 E^Q[|X_{i,k}^{(n)}|^4] \leq c'(T/n)^2$ for some $c' > 0$. Hence we have assertion (3) by Burkholder's Inequality. \square

3.2. Proposition on limit of value

Let $\bar{\mathcal{P}}$ be the set of probability measures \bar{P} on $(C[0, \infty)^M, \mathcal{B}[C[0, \infty)^M])$ such that the following is satisfied:

The coordinate function $w(\cdot) = (w_1(\cdot), \dots, w_M(\cdot))$ is a positive martingale with respect to $\{\mathcal{B}_t\}_{t \in [0, T]}$ under \bar{P} , where we define $\{\mathcal{B}_t\}_{t \in [0, T]}$ by $\mathcal{B}_t = \sigma(w(u); u \leq t), \quad 0 \leq t \leq T$.

$$\bar{P}(w_i(0) = 1, \quad i = 1, 2, \dots, M) = 1.$$

$\langle w_i, w_{i'} \rangle, \quad i, i' \in \{1, 2, \dots, M\}$ are absolutely continuous with respect to Lebesgue measure, and

$$\left(\frac{1}{w_i(u)w_{i'}(u)} \frac{d\langle w_i, w_{i'} \rangle}{dt}(u) \right)_{i, i' = 1, 2, \dots, M} \in \Gamma, \quad u \in [0, T], \quad \bar{P} \text{ a.s.}$$

Our purpose in this subsection is to prove the following.

PROPOSITION 3.2. *Take $g \in \hat{C}([0, \infty)^M : \mathbf{R})$ and an arbitrary subsequence (\bar{n}) of $(n)_{n \in \mathbf{N}}$.*

(1) *There exist a subsequence $(\bar{n}(k))_{k \in \mathbf{N}}$ of (\bar{n}) and a continuous mapping $W_g : [0, T] \times [0, \infty)^M \rightarrow \mathbf{R}$ such that*

$$W_g(t, x) = \lim_{k \rightarrow \infty} (H^{(\bar{n}(k))})^{\bar{n}(k) - [\bar{n}(k)t/T]} g(x),$$

for any $(t, x) \in [0, T] \times [0, \infty)^M$, where $[x]$ represents the greatest integer that is not greater than x . Also this convergence is uniform on any compact subsets on $[0, T] \times [0, \infty)^M$.

(2) $W_g(t, \cdot)$ belongs to $\hat{C}([0, \infty)^M : \mathbf{R})$ for each $t \in [0, T]$.

(3) There exists $\bar{P}_{t,x} \in \bar{\mathcal{P}}$ such that $W_g(t, x) = E^{\bar{P}_{t,x}}[g(xw(T-t))]$ holds for any $(t, x) \in [0, T] \times [0, \infty)^M$,

(4) $W_g(t, x) = \lim_{k \rightarrow \infty} (H^{(\bar{n}(k))})^{[\frac{\bar{n}(k)t'}{T}]} W_g(t+t', \cdot)(x)$, for $t, t' \in [0, T]$ with $t+t' \in [0, T]$ and $x \in [0, \infty)^M$.

(5) W_g is a viscosity subsolution of Hamilton-Jacobi-Bellman equation (13).

We need two lemmas to prove Lemma 3.2. Let us define $X(t, \omega; n) = (X_1(t, \omega; n), X_2(t, \omega; n), \dots, X_M(t, \omega; n))$, $t \in [0, T]$, $\omega \in \Omega$, $n \in \mathbf{N}$ by

$$\begin{aligned}
 X(t, \omega; n) &\equiv \frac{nt - T[nt/T]}{T} X_{[nt/T]+1}^{(n)} \\
 &\quad + \frac{T([nt/T] + 1) - nt}{T} X_{[nt/T]}^{(n)}, \quad t \neq T, \\
 (29) \quad X(T, \omega; n) &\equiv X_n^{(n)},
 \end{aligned}$$

i.e., $X(t; n)$ is the linear interpolation of $X^{(n)}$.

LEMMA 3.3. *Let $g \in \hat{C}([0, \infty)^M : \mathbf{R})$. Then we have*

$$|(H^{(n)})^{k'} g(x) - (H^{(n)})^k g(x)| \leq \bar{K} |x| \sqrt{\frac{T}{n}} |k' - k|, \quad k, k' \in \{0, 1, \dots, n\},$$

for some $\bar{K} > 0$ which does not depend on $n \in \mathbf{N}$.

PROOF. We may assume that $k' > k$. Let $K > 0$ be a constant such that $|g(x)| \leq K(1 + |x|^{2m})$, $|g(x) - g(x')| \leq K|x - x'|$, $x, x' \in [0, \infty)^M$ holds. By virtue of Lemma 3.1(3), we have

$$\begin{aligned}
 |E^Q[g(xX_{k'}^{(n)})] - E^Q[g(xX_k^{(n)})]| &\leq E^Q[|g(xX_{k'}^{(n)}) - g(xX_k^{(n)})|] \\
 &\leq K|x|E^Q[|X_{k'}^{(n)} - X_k^{(n)}|^4]^{\frac{1}{4}} \leq \bar{K}|x| \sqrt{\frac{T}{n}} |k' - k|
 \end{aligned}$$

for $Q \in \mathcal{Q}^{(n)}$. Hence we have the assertion. \square

LEMMA 3.4. *Let $(Q^{(n)})_{n \in \mathbf{N}}$, $Q^{(n)} \in \mathcal{Q}^{(n)}$ be an arbitrary sequence. We define a probability measure $P^{(n)}$ on $(C[0, \infty)^M, \mathcal{B}[C[0, \infty)^M])$ by $P^{(n)} = Q^{(n)} \circ X(\cdot; n)^{-1}$ for each $n \in \mathbf{N}$. Then the sequence $(P^{(n)})_{n \in \mathbf{N}}$ is tight. Moreover any cluster point of $(P^{(n)})_{n \in \mathbf{N}}$ belongs to $\bar{\mathcal{P}}$.*

PROOF. We can easily see that

$$\begin{aligned}
 E^{P^{(n)}}[|w(t) - w(t')|^4] &= E^{Q^{(n)}}[|X(t'; n) - X(t; n)|^4] \\
 &\leq K|t' - t|^2, \quad t, t' \in [0, T],
 \end{aligned}$$

for some $K > 0$ by Lemma 3.1. Hence $(P^{(n)})_{n \in \mathbf{N}}$ is tight.

Let \bar{P} be a cluster point of $(P^{(n)})_{n \in \mathbf{N}}$. Obviously we see that the coordinate function $w(\cdot) = (w_1(\cdot), w_2(\cdot), \dots, w_M(\cdot))$ is a positive martingale with respect to $\{\mathcal{B}_t\}_{t \in [0, T]}$ under \bar{P} , and $\bar{P}(w_i(0) = 1, i = 1, 2, \dots, M) = 1$.

For each $n \in \mathbf{N}$ and $\tilde{\omega} \in \Omega$, we define a probability measure $\hat{Q}_{\tilde{\omega}, k}^{(n)} \in \hat{\mathcal{Q}}$, $k = 0, 1, \dots, n - 1$ by $\hat{Q}_{\tilde{\omega}, k}^{(n)}[\hat{A}] = E^{\hat{P}}[\rho_k^{(n)}(\tilde{\omega}, \hat{Z})1_{\hat{A}}]$, where $E[dQ^{(n)}/dP|\mathcal{F}_n] = \prod_{l=1}^n \rho_l^{(n)}(\cdot, Z_l)$. Let $(g_{ii'})_{i, i'=1, 2, \dots, M} \in C([0, T] \times C[0, \infty)^M)^{M \times M}$ be a matrix valued function such that each $g_{ii'}$, $i, i' = 1, 2, \dots, M$ is bounded $\{\mathcal{B}_t\}$ -adapted function, and $(g_{ii'}(u, w)) \in \mathbf{R}^{M \times M}$ is nonnegative definite for all $(u, w) \in [0, T] \times C[0, \infty)^M$. For $n \in \mathbf{N}, k = 1, 2, \dots, n$ we have

$$\begin{aligned} & E^{P^{(n)}} \left[\sum_{i, i'=1}^M g_{ii'}\left(\frac{kT}{n}\right) \left(w_i\left(\frac{kT}{n} + \frac{T}{n}\right) - w_i\left(\frac{kT}{n}\right) \right) \left(w_{i'}\left(\frac{kT}{n} + \frac{T}{n}\right) - w_{i'}\left(\frac{kT}{n}\right) \right) \right] \\ &= E^{Q^{(n)}} \left[\sum_{i, i'=1}^M g_{ii'}\left(\frac{kT}{n}, X(\cdot; n)\right) E^{Q^{(n)}} \left[(X_{i, k+1}^{(n)} - X_{i, k}^{(n)}) (X_{i', k+1}^{(n)} - X_{i', k}^{(n)}) | \mathcal{F}_k \right] \right] \\ &= E^{Q^{(n)}} \left[\sum_{i, i'=1}^M g_{ii'}\left(\frac{kT}{n}, X(\cdot; n)\right) X_{i, k} X_{i', k} \frac{E^{\hat{Q}_{\tilde{\omega}, k}^{(n)}}[(\hat{Y}_i^{(n)} - 1)(\hat{Y}_{i'}^{(n)} - 1)] \frac{T}{n}}{T/n} \right] \\ &\leq E^{P^{(n)}} \left[\max_{\hat{Q}} \sum_{i, i'=1}^M g_{ii'}\left(\frac{kT}{n}\right) w_i\left(\frac{kT}{n}\right) w_{i'}\left(\frac{kT}{n}\right) \frac{E^{\hat{Q}}[(Y_i^{(n)} - 1)(Y_{i'}^{(n)} - 1)] \frac{T}{n}}{T/n} \right], \end{aligned}$$

where \hat{Q} runs $\hat{\mathcal{Q}} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Y}_i^{(n)}] = 1\}$. Then we see that

$$\begin{aligned} & \int_s^t \sum_{i, i'=1}^M g_{ii'}(u, w) d\langle w_i, w_{i'} \rangle_u \\ & \leq \int_s^t \left(\max_{\gamma \in \Gamma} \sum_{i, i'=1}^M \gamma_{ii'} g_{ii'}(u, w) w_i(u) w_{i'}(u) \right) du, \end{aligned}$$

for $t, s \in [0, T]$. Also we have

$$\int_s^t \sum_{i, i'=1}^M g_{ii'}(u, w) d\langle w_i, w_{i'} \rangle_u$$

$$\geq \int_s^t \left(\min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \gamma_{ii'} g_{ii'}(u, w) w_i(u) w_{i'}(u) \right) du,$$

for $t, s \in [0, T]$, by the same way. Hence we deduce that $\bar{P} \in \bar{\mathcal{P}}$. This completes the proof. \square

Now we show Proposition 3.2. Let $K > 0$ be a positive number such that $|g(x) - g(x')| \leq K|x - x'|$, $x, x' \in [0, \infty)^M$ and $|g(x)| \leq K(1 + |x|^{2m})$, $x \in [0, \infty)^M$ hold. We denote $W_g^{(n)}(t, x) = (H^{(n)})^{n - [nt/T]} g(x)$. Then using Lemma 3.3, we see that

$$\begin{aligned} & |W_g^{(n)}(t, x) - W_g^{(n)}(t, x')| \leq K|x - x'|, \\ & |W_g^{(n)}(t, x)| \leq K(1 + |x|^{2m}), \\ (30) \quad & |W_g^{(n)}(t, x) - W_g^{(n)}(t', x)| \leq K|x| \sqrt{\frac{T}{n} \left| \left[\frac{nt}{T} \right] - \left[\frac{nt'}{T} \right] \right|}, \end{aligned}$$

for some $K > 0$ which is independent of $n \in \mathbf{N}$.

(1): We see that the family $\{W_g^{(\bar{n})}(t, \cdot)\}_{n \in \mathbf{N}, t \in [0, T]} \subset \hat{C}([0, \infty)^M : \mathbf{R})$ is uniformly bounded and equicontinuous on any compact set of $[0, \infty)^M$. Then using Ascoli-Arzelà's theorem we see that there exists a continuous function $W_{1,g,t}$ on $[0, 1]^M$ for each $t \in [0, T]$ and a subsequence (\bar{n}_1) of (\bar{n}) , which does not depend on t , such that

$$(31) \quad \sup_{x \in [0, 1]^M} |W_g^{(\bar{n}_1)}(t, x) - W_{1,g,t}(x)| \rightarrow 0, \quad \bar{n}^1 \rightarrow \infty.$$

Also we see by the same way that there exists a continuous function $W_{2,g,t}$ on $[0, 2]^M$ for each $t \in [0, T]$ and a subsequence (\bar{n}_2) of (\bar{n}_1) , which does not depend on t , such that

$$(32) \quad \sup_{x \in [0, 2]^M} |W_g^{(\bar{n}_2)}(t, x) - W_{2,g,t}(x)| \rightarrow 0, \quad \bar{n}^2 \rightarrow \infty.$$

Obviously $W_{1,g,t} = W_{2,g,t}$ on $[0, 1]^M$, for each $t \in [0, T]$.

Then for each $l \in \mathbf{N}$, we can inductively define continuous functions $W_{l+1,g,t}$ on $[0, l + 1]^M$ for $t \in [0, T]$ and subsequences (\bar{n}_{l+1}) of (\bar{n}_l) , which does not depend on t , such that $W_{1,g,t} = W_{2,g,t} = \dots = W_{l+1,g,t}$ on $[0, l]^M$.

We also define continuous functions $W_{g,t}$ on $[0, \infty)^M$ for $t \in [0, T]$ by $W_{g,t}(x) = W_{l,g,t}(x)$ on $[0, l]^M$. Obviously $W_{g,t}$ is well-defined, and we see that the sequence $(\bar{n}(k))$, $\bar{n}(k) = \bar{n}_k(k)$, $k \in \mathbf{N}$ satisfies $\lim_{k \rightarrow \infty} W_g^{(\bar{n}(k))}(t, x) = W_{g,t}(x)$ for $(t, x) \in [0, T] \times [0, \infty)^M$. Using (30), we have

$$(33) \quad |W_{g,t}(x) - W_{g,t'}(x')| \leq K|x - x'| + K|x|\sqrt{|t - t'|}$$

for $(t, x), (t', x') \in [0, T] \times [0, \infty)^M$. This implies that the mapping $(t, x) \rightarrow W_{g,t}(x)$ is continuous. Then we can define a continuous mapping $W_g : [0, T] \times [0, \infty)^M$ by $W_g(t, x) = W_{g,t}(x)$. Obviously, W_g and $(\bar{n}(k))_{k \in \mathbf{N}}$ satisfy the assertion.

(2): We can easily show the assertion by letting $k \rightarrow \infty$ on both sides of the inequalities (30).

(3): Fix $(t, x) \in [0, T] \times [0, \infty)^M$. Hereafter we simply write $(n)_{n \in \mathbf{N}}$ instead of $(\bar{n}(k))_{k \in \mathbf{N}}$ for convention. We see by Lemma 3.1 that there exists $Q_l^{(n)} \in \mathcal{Q}^{(n)}$ for each n, l such that $W_g^{(n)}(t, x) \geq E^{Q_l^{(n)}}[g(xX_{n-\lfloor nt/T \rfloor}^{(n)})] \geq W_g^{(n)}(t, x) - 1/l$. Let $P_l^{(n)} = Q_l^{(n)} \circ X(\cdot; n)^{-1}$. Since $(P_l^{(n)})_{n, l \in \mathbf{N}}$ is tight by Lemma 3.4, then there exist a cluster point $P_{t,x} \in \bar{\mathcal{P}}$. Obviously $P_{t,x}$ satisfies the condition.

(4): Fix $t, t' \in [0, T]$, $x \in [0, \infty)^M$. First we claim that

$$(34) \quad |(H^{(n)})^{\lfloor nt'/T \rfloor} W_g^{(n)}(t + t', \cdot)(x) - (H^{(n)})^{\lfloor nt'/T \rfloor} W_g(t + t', \cdot)(x)| \rightarrow 0,$$

as $n \rightarrow \infty$. Fix $\varepsilon > 0$. Since the convergence $W_g^{(n)}(t + t', \cdot) \rightarrow W_g(t + t', \cdot)$, $n \rightarrow \infty$ is uniform on $\prod_{i=1}^M [0, x_i R]$, for each $R > 0$, there exists $n(R) \in \mathbf{N}$ such that

$$\sup_{y \in \prod_{i=1}^M [0, x_i R]} |W_g^{(n)}(t + t', y) - W_g(t + t', y)| < \varepsilon/2,$$

for $n > n(R)$. We denote

$$(35) \quad F^{(n)} = W_g^{(n)}(t + t', \cdot)(xX_{\lfloor nt'/T \rfloor}^{(n)}) - W_g(t + t', \cdot)(xX_{\lfloor nt'/T \rfloor}^{(n)}),$$

for $n \in \mathbf{N}$. Then for each $Q^{(n)} \in \mathcal{Q}^{(n)}$, $n > n(R)$ we have

$$\begin{aligned} & |E^{Q^{(n)}}[F^{(n)}]| \\ & \leq E^{Q^{(n)}}[|F^{(n)}|1_{\{|X_{[nt'/T]}^{(n)}| \leq R\}}] + E^{Q^{(n)}}[|F^{(n)}|1_{\{|X_{[nt'/T]}^{(n)}| > R\}}] \\ & \leq \frac{\varepsilon}{2} + E^{Q^{(n)}}[|F^{(n)}|^2]^{\frac{1}{2}} Q^{(n)}[|X_{[nt'/T]}^{(n)}| > R]^{\frac{1}{2}} \\ & \leq \frac{\varepsilon}{2} + \frac{K}{R} E^{Q^{(n)}}[1 + |X_{[nt'/T]}^{(n)}|^{2m}]^{\frac{1}{2}} E^{Q^{(n)}}[|X_{[nt'/T]}^{(n)}|^2]^{\frac{1}{2}} \\ & \leq \frac{\varepsilon}{2} + \frac{K'}{R}, \end{aligned}$$

where K, K' are positive numbers which do not depend on R, n , and $Q^{(n)}$. Let $R = \varepsilon/2K'$ and $n_0 = n(\varepsilon/2K')$. Then we have $|E^{Q^{(n)}}[F^{(n)}]| < \varepsilon$ and

$$|(H^{(n)})^{[nt'/T]}W_g^{(n)}(t + t', \cdot)(x) - (H^{(n)})^{[nt'/T]}W_g(t + t', \cdot)(x)| \leq \varepsilon,$$

for $n > n_0$. Hence we have the claim.

Also we see that $|W_g^{(n)}(t, x) - W_g(t, x)| \rightarrow 0$, as $n \rightarrow \infty$ and,

$$\begin{aligned} & |(H^{(n)})^{[\frac{nt'}{T}]}W_g^{(n)}(t + t', \cdot)(x) - W_g^{(n)}(t, x)| \\ & = |(H^{(n)})^{n - [n(t+t')/T] + [nt'/T]}g(x) - (H^n)^{n - [nt'/T]}g(x)| \\ & \leq K|x|\sqrt{\frac{T}{n}|\lceil \frac{nt'}{T} \rceil + \lceil \frac{nt'}{T} \rceil - \lceil \frac{n(t+t')}{T} \rceil|} \rightarrow 0, \end{aligned}$$

as $n \rightarrow \infty$. We have the assertion from these results.

(5): Take $(\bar{t}, \bar{x}) \in [0, T] \times [0, \infty)^M$. Let $\hat{U} \in C^\infty([0, T] \times [0, \infty)^M)$ be a function such that $\hat{U}(\bar{t}, \bar{x}) = W_g(\bar{t}, \bar{x})$ and $\hat{U} \geq W_g$ on some neighbourhood V of (\bar{t}, \bar{x}) hold. We can easily see by (2) that $W_g(t, x) \leq K(1 + |x|^{2m})$ for some $K > 0$, $m \in \mathcal{M}$. Then we may assume that $\hat{U}(t, x) \geq W_g(t, x)$, and $|\hat{U}(t, x)| + |(\partial\hat{U}/\partial t)(t, x)| + \sum_{i,i'} |x_i x_{i'} (\partial^2\hat{U}/\partial x_i \partial x_{i'})(t, x)| \leq K(1 + |x|^{2m})$, for any $(t, x) \in [0, T] \times [0, \infty)^M$. Let $\bar{\gamma} = (\bar{\gamma}_{i,i'}) \in \Gamma$ be the element such that

$$\begin{aligned} & \sum_{i,i'=1}^M \frac{1}{2} \bar{\gamma}_{i,i'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}) = \min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{i,i'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}) \\ & = \min_{\hat{Q}} \sum_{i,i'=1}^M \frac{1}{2} E^{\hat{Q}}[\hat{Z}_i \hat{Z}_{i'}] \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}), \end{aligned}$$

where \hat{Q} runs $\hat{Q} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Z}_i] = 0\}$.

Also, let $\hat{Q}^{(n)} \in \hat{Q} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Y}_i^{(n)}] = 1\}$, $n \in \mathbf{N}$ be measures that attain the minimal of

$$\min_{\hat{Q}} \sum_{i,i'=1}^M \frac{1}{2} \frac{E^{\hat{Q}}[(\hat{Y}_i^{(n)} - 1)(\hat{Y}_{i'}^{(n)} - 1)]}{T/n} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}), \quad n \in \mathbf{N},$$

where \hat{Q} runs $\hat{Q} \cap \bigcap_{i=1}^M \{\hat{Q} \in \hat{\mathcal{P}} \mid E^{\hat{Q}}[\hat{Y}_i^{(n)}] = 1\}$ for each $n \in \mathbf{N}$. Note that we can naturally regard $\hat{Q}^{(n)}$ as a probability measure on (Ω, \mathcal{F}_1) for each $n \in \mathbf{N}$. Take $\rho^{(n)} \in \mathcal{L}_1$ such that $d\hat{Q}^{(n)}/dP = \rho^{(n)}(\cdot, Z_1)$ holds. We define $\bar{Q}^{(n)} \in \mathcal{Q}^{(n)}$ by $E[\frac{d\bar{Q}^{(n)}}{dP} | \mathcal{F}_n] = \prod_{k=1}^n \bar{\rho}_k^{(n)}(\cdot, Z_k)$, where $\bar{\rho}_k^{(n)} = \rho^{(n)}$, $k = 1, 2, \dots, n$, and $\bar{P}^{(n)} = \bar{Q}^{(n)} \circ X(\cdot; n)^{-1}$ for $n \in \mathbf{N}$. We see that there exists a cluster point $\bar{P} \in \mathcal{P}$ of $\{\bar{P}^{(n)}\}_{n \in \mathbf{N}}$ and $d\langle w_i, w_{i'} \rangle_u = \bar{\gamma}_{i,i'} w_i(u) w_{i'}(u) du$, \bar{P} -a.s., by an argument similar to that of the proof of Lemma 3.4. Then we have

$$\begin{aligned} \hat{U}(\bar{t}, \bar{x}) &= W_g(\bar{t}, \bar{x}) = \lim_{k \rightarrow \infty} (H^{(n)})^{[nh/T]} W_g(\bar{t} + h, \cdot)(\bar{x}) \\ &\leq \limsup_{k \rightarrow \infty} E^{\bar{P}^{(n)}} [W_g(\bar{t} + h, \bar{x} w(\frac{T}{n} [\frac{nh}{T}]))] \\ &= E^{\bar{P}} [W_g(\bar{t} + h, \bar{x} w(h))] \leq E^{\bar{P}} [\hat{U}(\bar{t} + h, \bar{x} w(h))]. \end{aligned}$$

Then we see by Ito's formula that

$$(36) \quad 0 \leq E^{\bar{P}} \left[\int_0^h \left(\frac{\partial \hat{U}}{\partial t} + \sum_{i,i'=1}^M \frac{1}{2} \bar{\gamma}_{i,i'} x_i x_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}} \right) (\bar{t} + u, \bar{x} w(u)) du \right].$$

Dividing both sides by $h > 0$ and letting $h \rightarrow \infty$, we have

$$\begin{aligned} 0 &\leq \left(\frac{\partial \hat{U}}{\partial t} + \sum_{i,i'=1}^M \frac{1}{2} \bar{\gamma}_{i,i'} x_i x_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}} \right) (\bar{t}, \bar{x}) \\ &= \left(\frac{\partial \hat{U}}{\partial t} + \min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{i,i'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}} \right) (\bar{t}, \bar{x}). \end{aligned}$$

This completes the proof.

3.3. Conclusion

Now we prove Theorem 1.5, which is our main result in this paper. Take an arbitrary subsequence (\bar{n}) of $(n)_{n \in \mathbf{N}}$ and define W_f as in Proposition 3.2. Using Proposition 3.2 (3), We have $\inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[f(xw(T-t))] \leq W_f(t, x)$ for $(t, x) \in [0, T] \times [0, \infty)^M$. On the other hand, we have the inverse inequality by Theorem A.2, Proposition A.4, and Proposition 3.2 (3). Then we have

$$W_f(t, x) = \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[f(xw(T-t))].$$

Since the subsequence (\bar{n}) is arbitrary, $U(t, x) = \lim_{n \rightarrow \infty} (H^{(n)})^{n-[nt/T]T/n} f(x)$ exists for any $t \in [0, T]$, $x \in [0, \infty)^M$, and equals $\inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[f(xw(T-t))]$. Then we see that U is a viscosity solution of (13) because $U(t, x) = W_f(t, x) = E^{\bar{P}}[f(xw(T-t))]$ is both a supersolution and subsolution. Also the uniqueness holds by Corollary A.2. Hence we have the assertion from Theorem 2.2. This completes the proof.

Appendix A. Some remarks on a Bellman equation and viscosity solution

We recall the definition and some property of viscosity solution in this appendix. The reader also refer to [10] for detail.

DEFINITION A.1. We say that a continuous function $U : [0, T] \times [0, \infty)^M \rightarrow \mathbf{R}$ is a viscosity supersolution (*resp. subsolution*) of Hamilton-Jacobi-Bellman equation (13), if

$$\left(\frac{\partial \hat{U}}{\partial t} + \inf_{\gamma \in \Gamma} \sum_{i, i'=1}^M \frac{1}{2} \gamma_{i, i'} x_i x_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}\right)(\bar{t}, \bar{x}) \leq 0 \text{ (resp. } \geq 0)$$

holds for any $(\bar{t}, \bar{x}) \in [0, T] \times [0, \infty)^M$ and $\hat{U} \in C^\infty([0, T] \times [0, \infty)^M)$ such that $\hat{U}(\bar{t}, \bar{x}) = U(\bar{t}, \bar{x})$ and $\hat{U} - U$ takes its local maximum (*resp. local minimum*) value 0 at (\bar{t}, \bar{x}) . Also we say that a function $U : [0, T] \times [0, \infty)^M \rightarrow \mathbf{R}$ is a viscosity solution if it is both a viscosity supersolution and subsolution.

We will need the following comparison theorem for a viscosity supersolution and a subsolution due to [17].

THEOREM A.2. *Let \bar{U} and \underline{U} be a viscosity supersolution and a subsolution of Hamilton-Jacobi-Bellman equation (13). If the following inequalities:*

$$\begin{aligned} \sup_{[0,T] \times [0,\infty)^M} \underline{U}(t,x)/(|x|^2 + 1)^m &< \infty, \\ \inf_{[0,T] \times [0,\infty)^M} \bar{U}(t,x)/(|x|^2 + 1)^m &> -\infty, \quad m > 0, \\ \underline{U}(T,x) &\leq \bar{U}(T,x), \quad x \in [0,\infty)^M, \end{aligned}$$

hold, then we have $\underline{U}(t,x) \leq \bar{U}(t,x)$, $(t,x) \in [0,T] \times [0,\infty)^M$. In particular, the viscosity solution U of Hamilton-Jacobi-Bellman equation (13) satisfying $U \in \hat{C}([0,\infty)^M : \mathbf{R})$ is unique.

Before we state a proposition on Hamilton-Jacobi-Bellman equation (13), we prove a lemma.

LEMMA A.3. *There exists $C_m > 0$ for each $m \in \mathbf{N}$ such that*

$$\begin{aligned} E^{\bar{P}}[\max_{u \in [0,t]} |w(u) - \mathbf{1}|^{2m}] &\leq C_m t^m, \\ E^{\bar{P}}[\max_{u \in [0,T]} |w(u)|^{2m}] &\leq C_m, \quad t \in [0,T], \quad \bar{P} \in \bar{\mathcal{P}}. \end{aligned}$$

PROOF. We see by Burkholder's inequality that

$$\begin{aligned} (37) \quad E^{\bar{P}}[\max_{u \in [0,t]} |w(u) - \mathbf{1}|^{2m}] &\leq c E^{\bar{P}}[\sum_{i=1}^M \langle w_i \rangle_u^m] \\ &\leq c E^{\bar{P}}[(\int_0^t \max_{\gamma \in \Gamma} \sum_{i=1}^M \gamma_{ii} |w_i(u)|^2 du)^m] \leq c \bar{\gamma}^m E^{\bar{P}}[(\int_0^t |w(u)|^2 du)^m] \\ &\leq ct^m E^{\bar{P}}[\int_0^t |w(u)|^{2m} du] \leq c + ct^m E^{\bar{P}}[\int_0^t \max_{s \in [0,u]} |w(u) - \mathbf{1}|^{2m} du] \end{aligned}$$

where all c stand for positive numbers (not necessarily equal) which do not depend on $\bar{P} \in \bar{\mathcal{P}}$ and t . Then we have the assertion by Gronwall's inequality. \square

Let $\bar{U}(t, x) = \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[\tilde{f}(xw(T - t))]$. Using Lemma A.3, We can easily see that $\bar{U} \in \hat{C}([0, \infty)^M : \mathbf{R})$. Now we show the following.

PROPOSITION A.4. $\bar{U}(t, x)$ is a viscosity supersolution of (13).

PROOF. First we denote by Λ a set of control which is composed of pairs $\{(\Omega, \mathcal{F}, P; \{\mathcal{F}_t\}_{t \in [0, T]}, X)\}$ such that the following satisfied:

$(\Omega, \mathcal{F}, P; \{\mathcal{F}_t\}_{t \in [0, T]})$ is a filtered probability space,

$X = (X_1, X_2, \dots, X_M)$ is a continuous positive martingale with respect to $\{\mathcal{F}_t\}_{t \in [0, T]}$ under P ,

$P(X_i(0) = 1, i = 1, 2, \dots, M) = 1$.

$\langle X_i, X_{i'} \rangle, i, i' \in \{1, 2, \dots, M\}$ are absolutely continuous with respect to Lebesgue measure, and

$$(38) \quad \left(\frac{1}{X_i(u)X_{i'}(u)} \frac{d\langle X_i, X_{i'} \rangle}{dt}(u) \right)_{i, i' = 1, 2, \dots, M} \in \Gamma, \quad u \in [0, T], \quad P \text{ a.s.}$$

We define $Q_t g(x), t \in [0, T], x \in [0, \infty)^M$, and $g \in \bar{C}([0, \infty)^M : \mathbf{R})$ by

$$(39) \quad Q_t g(x) = \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[g(xw(t))].$$

Then we can easily see that $Q_t g(x) = \inf_{\lambda \in \Lambda} E^P[g(xX_t)]$ since $PX^{-1} \in \bar{\mathcal{P}}$.

Step 1. We show that $Q_{t+t'}g(x) \geq Q_t Q_{t'}g(x), t, t' \in [0, T], x \in [0, \infty)^M$. We define a filtration $\{\hat{\mathcal{B}}_u\}_{u \in [0, T]}$ on $(C[0, \infty)^M, \mathcal{B}[C[0, \infty)^M])$ and a M -dimensional $\{\hat{\mathcal{B}}_u\}$ -adapted process $\hat{X}_u = (\hat{X}_{1,u}, \hat{X}_{2,u}, \dots, \hat{X}_{M,u}), u \in [0, T]$, by

$$\hat{\mathcal{B}}_u = \begin{cases} \mathcal{B}_{t+u}, & 0 \leq u \leq T - t, \\ \mathcal{B}_T, & T - t < u \leq T. \end{cases}$$

$$\hat{X}_{i,u}(w) = \begin{cases} w_i(t+u)/w_i(t), & 0 \leq u \leq T - t, \\ w_i(T)/w_i(t), & T - t < u \leq T. \end{cases}$$

Since $(C[0, \infty)^M, \mathcal{B}[C[0, \infty)^M], \bar{P})$ is standard probability space for each $\bar{P} \in \bar{\mathcal{P}}$, there exists a regular conditional measure $\bar{P}_t(w, B) : C[0, \infty)^M \times \mathcal{B}[C[0, \infty)^M] \rightarrow [0, 1]$, $t \in [0, T]$. Then we can easily see that

$$((C[0, \infty)^M, \mathcal{B}[C[0, \infty)^M], \bar{P}_t(w, \cdot) ; \{\hat{\mathcal{B}}_u\}_{u \in [0, T]}, \hat{X}) \in \Lambda,$$

and

$$\begin{aligned} E^{\bar{P}}[g(xw(t+t'))] &= E^{\bar{P}}[E^{\bar{P}}[g(xw(t+t'))|\mathcal{B}_t]] \\ &= E^{\bar{P}}\left[\int g(xw(t')\hat{X}_{t'})\hat{P}_t(w, dw')\right] \geq E^{\hat{P}}[Q_{t'}g(xw(t))] \geq Q_t Q_{t'}g(x), \end{aligned}$$

for $\bar{P} \in \bar{\mathcal{P}}$. Hence we have the assertion.

Step 2. Take $(\bar{t}, \bar{x}) \in [0, T] \times [0, \infty)^M$ and fix it. Let $\hat{U} \in C^\infty([0, T] \times [0, \infty)^M)$ be a function such that $\hat{U}(\bar{t}, \bar{x}) = \bar{U}(\bar{t}, \bar{x})$ and $\hat{U} \leq \bar{U}$ on some neighbourhood V of (\bar{t}, \bar{x}) hold. We may assume that $\hat{U}(t, x) \leq \bar{U}(t, x)$ and,

$$(40) \quad |\hat{U}(t, x)| + \left|\frac{\partial \hat{U}}{\partial t}(t, x)\right| + \sum_{i, i'} |x_i x_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(t, x)| \leq K(1 + |x|^{2m}),$$

for any $(t, s) \in [0, T] \times [0, \infty)$, because $\bar{U} \in \hat{C}([0, \infty)^M : \mathbf{R})$.

From here to the end of this proof, $c > 0$ will stand for positive numbers (not necessarily equal) which do not depend on $\bar{P} \in \bar{\mathcal{P}}$, $t \in [0, T]$, $\gamma \in \Gamma$, and $R > 0$. First we claim that there exists $M(R) > 0$ for each $R > 0$ such that $M(R)$ does not depend on $\bar{P} \in \bar{\mathcal{P}}$, $t \in [0, T]$, $\gamma \in \Gamma$, and

$$\begin{aligned} &|E^{\bar{P}}\left[\int_0^{t-\bar{t}} \frac{\partial \hat{U}}{\partial t}(\bar{t} + u, \bar{x}w(u))du\right. \\ &\quad \left. + \int_0^{t-\bar{t}} \sum_{i, i'=1}^M \frac{1}{2} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t} + u, \bar{x}w(u))d\langle w_i, w_{i'} \rangle_u\right] \\ &\quad - \left\{\frac{\partial \hat{U}}{\partial t}(\bar{t}, \bar{x})(t - \bar{t}) + E^{\bar{P}}\left[\sum_{i, i'=1}^M \frac{1}{2} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x})\langle w_i, w_{i'} \rangle_{t-\bar{t}}\right]\right\} \\ &\leq \frac{c(t - \bar{t})}{R} + cM(R)(t - \bar{t})^{3/2}, \end{aligned}$$

for $t > \bar{t}$. To show this claim, we estimate

$$\begin{aligned}
 & I_1 + I_2 \\
 &= E^{\bar{P}} \left[\left| \int_0^{t-\bar{t}} \sum_{i,i'=1}^M \frac{1}{2} \bar{x}_i \bar{x}_{i'} F_{ii'}(u, w) d\langle w_i, w_{i'} \rangle_u \right| 1_{\left\{ \max_{u \in [0, t-\bar{t}]} |w(u)| > R \right\}} \right] \\
 (41) \quad &+ E^{\bar{P}} \left[\left| \int_0^{t-\bar{t}} \sum_{i,i'=1}^M \frac{1}{2} \bar{x}_i \bar{x}_{i'} F_{ii'}(u, w) d\langle w_i, w_{i'} \rangle_u \right| 1_{\left\{ \max_{u \in [0, t-\bar{t}]} |w(u)| \leq R \right\}} \right],
 \end{aligned}$$

where $F_{i,i'}(u, w) = \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t} + u, \bar{x}w(u)) - \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x})$, $i, i' = 1, 2, \dots, M$, $R > 0$, $\bar{P} \in \bar{\mathcal{P}}$, and

$$\begin{aligned}
 J_1 + J_2 &= E^{\bar{P}} \left[\int_0^{t-\bar{t}} |G(u, w)| du 1_{\left\{ \max_{u \in [0, t-\bar{t}]} |w(u)| > R \right\}} \right] \\
 (42) \quad &+ E^{\bar{P}} \left[\int_0^{t-\bar{t}} |G(u, w)| du 1_{\left\{ \max_{u \in [0, t-\bar{t}]} |w(u)| \leq R \right\}} \right],
 \end{aligned}$$

where $G(u, w) = \frac{\partial \hat{U}}{\partial t}(\bar{t} + u, \bar{x}w(u)) - \frac{\partial \hat{U}}{\partial t}(\bar{t}, \bar{x})$, $R > 0$, $\bar{P} \in \bar{\mathcal{P}}$. We have the claim by Lemma A.3, Hölder’s inequality, Tchebychev’s inequality, Burkholder’s inequality, and Lipschitz continuity of all derivatives of \hat{U} on a bounded interval. Also we have

$$\begin{aligned}
 & |E^{\bar{P}} \left[\int_0^{t-\bar{t}} \left(\min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{ii'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}) w_i(u) w_{i'}(u) \right) du \right]| \\
 & - (t - \bar{t}) \left(\min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{ii'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}) \right) \leq c|t - \bar{t}|^{3/2}.
 \end{aligned}$$

by Lemma A.3 and Hölder’s inequality.

Step 3. We see by Ito’s formula that

$$\begin{aligned}
 \hat{U}(\bar{t}, \bar{x}) &= \bar{U}(\bar{t}, \bar{x}) = Q(T - \bar{t}) \tilde{f}(\bar{t}, \bar{x}) \\
 &\geq Q(t - \bar{t}) Q(T - \bar{t}) \tilde{f}(x) = Q(t - \bar{t}) \bar{U}(t, \cdot)(x)
 \end{aligned}$$

$$\begin{aligned}
 &= \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[\bar{U}(t, \bar{x}w(t - \bar{t}))] \geq \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}[\hat{U}(t, \bar{x}w(t - \bar{t}))] \\
 &= \hat{U}(\bar{t}, \bar{x}) + \inf_{\bar{P} \in \bar{\mathcal{P}}} E^{\bar{P}}\left[\int_0^{t-\bar{t}} \frac{\partial \hat{U}}{\partial t}(\bar{t} + u, \bar{x}w(u))du \right. \\
 &\quad \left. + \int_0^{t-\bar{t}} \sum_{i,i'=1}^M \frac{1}{2} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t} + u, \bar{x}w(u))d\langle w_i, w_{i'} \rangle_u\right].
 \end{aligned}$$

Then we have

$$\begin{aligned}
 0 &\geq -c(M(R) + 1)(t - \bar{t})^{3/2} - \frac{c(t - \bar{t})}{R} + \frac{\partial \hat{U}}{\partial t}(\bar{t}, \bar{x})(t - \bar{t}) \\
 &\quad + (t - \bar{t})\left(\min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{ii'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x})\right), \quad \bar{t} < t \leq T, \quad R > 0,
 \end{aligned}$$

by the consequence of step 2. Dividing both sides of the above inequality by $t - \bar{t} > 0$ and letting $t \rightarrow \bar{t}$, we have

$$0 \geq -\frac{c}{R} + \frac{\partial \hat{U}}{\partial t}(\bar{t}, \bar{x}) + \min_{\gamma \in \Gamma} \sum_{i,i'=1}^M \frac{1}{2} \gamma_{ii'} \bar{x}_i \bar{x}_{i'} \frac{\partial^2 \hat{U}}{\partial x_i \partial x_{i'}}(\bar{t}, \bar{x}),$$

for all $R > 0$. Hence we have the assertion. This completes the proof. \square

Acknowledgement. The author would like to thank Professor Shigeo Kusuoka for his useful advices.

References

- [1] Artzner, P., Delbaen, F., Eber, J.-M. and D. Heath, Coherent measures of risk, *Math. Finance* **9**(3) (1999), 203–228.
- [2] Artzner, P., Delbaen, F., Eber, J.-M., Heath, D. and H. Ku, Multiperiod risk and coherent multiperiod risk measurements of risk, Preprint, ETH Zürich (2002).
- [3] Acerbi, C. and D. Tasche, On the Coherence of Expected Shortfall, Working Paper (2001).
- [4] Barrieu, P. and N. El Karoui, Inf-Convolution of risk measures and optimal risk transfer, *Finance Stoch.* **9** (2005), 269–298.
- [5] Barrieu, P. and N. El Karoui, Pricing, hedging and optimally designing derivatives via minimization of risk measures, Preprint (2005).

- [6] Cheridito, P., Delbaen, F. and M. Kupper, Coherent and convex risk measures for bounded cadlag processes, *Stoc. Appl.* **112** (2004), 1–22.
- [7] Cherny, A. S., Pricing with coherent risk, *Probability Theory and Its Applications* **52**(3) (2007), 506–540.
- [8] Cherny, A. S., Pricing and hedging European options with discrete-time coherent risk, *Finance Stoch.* **11** (2007), 537–569.
- [9] Cherny, A. S. and D. B. Madan, Pricing and hedging in incomplete markets with coherent risk, Preprint (2006).
- [10] Crandall, M. G., Ishii, H. and P. L. Lions, User’s guide to viscosity solutions of second order partial differential equations, *Bull. A. M. S.* **27** (1993), 1–67.
- [11] Delbaen, F., Coherent risk measures, Lecture notes, Pisa (2001).
- [12] Delbaen, F., The structure of m-stable sets and in particular of the set of risk neutral measures, Preprint, ETH Zürich (2001).
- [13] Detlefsen, K. and G. Scandolo, Conditional and dynamic convex risk measures, *Finance Stoch.* **9** (2005), 539–561.
- [14] Föllmer, H. and A. Schied, Convex measures of risk and trading constraints, *Finance Stoch.* **6** (2002), 429–447.
- [15] Föllmer, H. and A. Schied, *Stochastic Finance*, 2nd Ed., Walter de Gruyter (2004).
- [16] Kusuoka, S. and Y. Morimoto, Homogeneous law invariant coherent multiperiod value measures and their limits, *J. Math. Sci. Univ. Tokyo* **14**(2) (2007), 117–156.
- [17] Nagai, H., *Stochastic Differential Equations*, Kyoritsu (In Japanese) (1999).
- [18] Nakano, Y., Efficient hedging with coherent risk measure, *J. Math. Anal. Appl.* **293**(1) (2004), 345–354.
- [19] Roorda, B., Martingale characterizations of coherent acceptability measures, working paper, Twente university (2002).
- [20] Takahashi, W., Nonlinear variational inequalities and fixed point theorems, *J. Math. Soc. Japan* **28**(1) (1976), 168–181.
- [21] Umezawa, Y., The minimal risk of hedging with a convex risk measure, *Advances in Mathematical Economics* **9** (2006), 109–116.

(Received March 2, 2010)

(Revised February 28, 2011)

Mizuho-DL Financial Technology Co. Ltd.
Ote Center Building 16F
1-3, Otemachi 1-chome, Chiyodaku
Tokyo 100-0004, Japan
E-mail: yuji-umezawa@fintec.co.jp