


UTMS 2008–18

July 2, 2008

**Nonexistence of
global solutions in time for
reaction-diffusion systems with
inhomogeneous terms in cones**

by

Takefumi IGARASHI and Noriaki UMEDA



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

Nonexistence of Global Solutions in Time for Reaction-Diffusion Systems with Inhomogeneous Terms in Cones

Takefumi Igarashi,

College of Science and Technology, Nihon University,
7-24-1, Narashino-dai, Funabashi-shi, Chiba, 274-8501, Japan,
E-mail: igarashit@penta.ge.cst.nihon-u.ac.jp

and

Noriaki Umeda,

Graduate School of Mathematical Sciences, University of Tokyo,
3-8-1, Komaba, Meguro-ku, Tokyo, 153-8914, Japan,
E-mail: umeda@ms.u-tokyo.ac.jp

Abstract

We consider initial-boundary value problems for the reaction-diffusion systems with inhomogeneous terms in cones. In this paper we show the nonexistence of global solutions of the problems in time.

Keyword and Phrases: reaction-diffusion, global nonexistence, inhomogeneous term, cone.

AMS subject classifications: 35K45, 35K57.

1 Introduction

We consider nonnegative solutions of initial-boundary value problems for the reaction-diffusion systems of the form

$$\begin{cases} u_t = \Delta u + K_1(x, t)v^{p_1}, & x \in D, t > 0, \\ v_t = \Delta v + K_2(x, t)u^{p_2}, & x \in D, t > 0, \\ u(x, t) = v(x, t) = 0, & x \in \partial D, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in D, \end{cases} \quad (1)$$

where $p_1, p_2 \geq 1$ with $p_1 p_2 > 1$. The domain D is a cone in \mathbf{R}^N , such as

$$D = \{x \in \mathbf{R}^N; x \neq 0 \text{ and } x/|x| \in \Omega\}, \quad (2)$$

where Ω is some region on S^{N-1} smooth enough.

The initial data $u_0(x)$ and $v_0(x)$ are nonnegative, bounded and continuous in \bar{D} , and $u_0(x) = v_0(x) = 0$ on ∂D . The inhomogeneous terms K_i ($i = 1, 2$) are nonnegative continuous functions in $D \times (0, \infty)$.

In this paper we denote by BC the set of all bounded continuous functions in \bar{D} . The ‘‘nontrivial solution’’ denotes the solution u satisfying $(u, v) \not\equiv 0$ in $D \times (0, T)$ with some $T > 0$, it thus means that $(u_0, v_0) \not\equiv 0$ with the condition $(u_0, v_0) \in BC$.

For the Laplace-Beltrami operator with homogeneous Dirichlet boundary condition on $\Omega \in S^{N-1}$, define ω_n as Dirichlet eigenvalues and $\psi_n(\theta)$ as the Dirichlet eigenfunctions corresponding to ω_n which is normalized so that

$$\int_{\Omega} \psi_n(\theta) d\theta = 1.$$

It is following that

$$\int_{\Omega} \psi_m(\theta) \psi_n(\theta) d\theta = 0$$

for $m \neq n$. We introduce the Green’s function $G(x, y, t) = G(r, \theta, \rho, \phi, t)$ for the linear heat equation in the cone D , where

$$r = |x|, \quad \rho = |y|, \quad \theta = x/|x| \text{ and } \phi = y/|y| \in \Omega \quad (3)$$

The Green’s function is expressed to

$$G(r, \theta, \rho, \phi, t) = \frac{1}{2t} (r\rho)^{-(N-2)/2} \exp\left(-\frac{\rho^2 + r^2}{4t}\right) \sum_{n=1}^{\infty} I_{\nu_n}\left(\frac{r\rho}{2t}\right) \psi_n(\theta) \psi_n(\phi), \quad (4)$$

where $\nu_n = [(N-2)^2/4 + \omega_n]^{1/2}$, and I_{ν} is the modified Bessel function or

$$I_{\nu}(z) = \left(\frac{z}{2}\right)^{\nu} \sum_{k=0}^{\infty} \frac{(z/2)^{2k}}{k! \Gamma(\nu + k + 1)} \quad (5)$$

with the Gamma function $\Gamma(z) = \int_0^{\infty} s^{z-1} e^{-s} ds$ (see Watson [27, p.p.395]).

For our first theorem we shall give the conditions of the inhomogeneous terms K_i ($i = 1, 2$) as following:

$$\left. \begin{array}{l} \text{there exist } C_U, \hat{\sigma}_i \text{ and } \hat{q}_i \geq 0 \text{ such that} \\ K_i(x, t) \leq C_U \langle x \rangle^{\hat{\sigma}_i} (t+1)^{\hat{q}_i} \quad \text{for any } x \in D, t \geq 0, \end{array} \right\} \quad (6)$$

where $\langle x \rangle = (|x|^2 + 1)^{1/2}$.

Let L_a^∞ be a Banach space of L^∞ -functions in D with the norm

$$\|\xi\|_{\infty, a} \equiv \text{esssup}_{x \in D} (\langle x \rangle^a |\xi(x)|).$$

For $T > 0$, set

$$E_T = \{(u, v) : [0, T] \rightarrow L_{\delta_1}^\infty \times L_{\delta_2}^\infty; \|(u, v)\|_{E_T} < \infty\} \quad (7)$$

with the norm

$$\|(u, v)\|_{E_T} := \sup_{t \in [0, T]} \{\|u(t)\|_{\infty, \delta_1} + \|v(t)\|_{\infty, \delta_2}\},$$

where

$$\delta_i = \frac{\hat{\sigma}_j p_i + \hat{\sigma}_i}{p_i p_j - 1} \quad ((i, j) = (1, 2), (2, 1)). \quad (8)$$

It is easily seen that E_T is a Banach space.

We begin with stating the existence of the local solution for (1).

Theorem 1. *Assume that $u_0, v_0 \in BC$, $u_0 \equiv v_0 \equiv 0$ on ∂D , and $\langle x \rangle^{\delta_1} u_0(x)$, $\langle x \rangle^{\delta_2} v_0(x)$ are bounded in \bar{D} . Suppose that $K_i(x, t)$ ($i = 1, 2$) satisfy (6). Then there exists a nonnegative solution $(u, v) \in E_T$ which solves (1) in $D \times (0, T)$ for some $T > 0$.*

For given initial values (u_0, v_0) , let $T^* = T^*(u_0, v_0)$ be a maximal existence time of the solution of (1). If $T^* = \infty$, the solutions are global in time. On the other hand, if $T^* < \infty$, then the solutions are not global in time. If the solution blows up in finite time such that

$$\limsup_{t \rightarrow T^*} \|u(\cdot, t)\|_\infty + \limsup_{t \rightarrow T^*} \|v(\cdot, t)\|_\infty = \infty, \quad (9)$$

then the solution is not global, where $\|\cdot\|_\infty$ denotes the L^∞ -norm with respect to space variable.

For our second theorem we shall define a region \tilde{D} such that

$$\left. \begin{array}{l} \text{there exist } k > 0 \text{ and } \{x_m\}_{m=1}^\infty \text{ satisfying } 0 < |x_m| < |x_{m+1}|, \\ B(x_m, k|x_m|) \subset \tilde{D} \subset D \text{ for any } m, \text{ and } \lim_{m \rightarrow \infty} |x_m| = \infty, \end{array} \right\} \quad (10)$$

where $B(x, r)$ denotes the ball with radius r centered at x . We let the inhomogeneous terms K_i ($i = 1, 2$) satisfy

$$\left. \begin{array}{l} \text{there exist } C_L > 0, \check{\sigma}_i, \check{q}_i \geq 0 \text{ and } \tilde{D} \text{ satisfying (10) such that} \\ K_i(x, t) \geq C_L |x|^{\check{\sigma}_i} t^{\check{q}_i} \quad \text{for any } x \in \tilde{D}, t \geq 0. \end{array} \right\} \quad (11)$$

For the theorem we should define γ_+ denoting the positive root of the equation $\gamma(\gamma + N - 2) = \omega_1$,

$$\alpha_i = \frac{(2 + \check{\sigma}_i + 2\check{q}_i) + (2 + \check{\sigma}_j + 2\check{q}_j)p_i}{p_i p_j - 1} \quad ((i, j) = (1, 2), (2, 1)), \quad (12)$$

and

$$H_a = \{\xi \in C(\tilde{D}); \xi(x) \geq M \langle x \rangle^{-a} \psi_1(x/|x|) \text{ for } x \in \tilde{D} \text{ with some } M > 0\}.$$

The main result of this paper is summarized in the following theorem.

Theorem 2. *Assume that $u_0, v_0 \in BC$, $u_0 \equiv v_0 \equiv 0$ on ∂D , and $K_i(x, t)$ ($i = 1, 2$) satisfy (11). Suppose that one of the following two conditions holds;*

$$(i) \max\{\alpha_1, \alpha_2\} \geq N + \gamma_+.$$

$$(ii) u_0 \in H_{a_1} \text{ with } a_1 < \alpha_1 \text{ or } v_0 \in H_{a_2} \text{ with } a_2 < \alpha_2.$$

Then, there exists no nontrivial nonnegative global solution of (1).

The method using the sequence of balls in (11) was used in [4, 22] and other papers.

Remark. (i) *It is easily seen that $\gamma_+ = \nu_1 - (N - 2)/2$.*

(ii) *If both (6) and (11) hold, then it is necessarily that $C_U \geq C_L$, $\hat{\sigma}_i \geq \check{\sigma}_i$ and $\hat{q}_i \geq \check{q}_i$.*

We briefly recall a history of the study on global nonexistence of solutions to the system (1). First, the global nonexistence of solutions in the case $D = \mathbf{R}^N$ ($\Omega = S^{N-1}$), $u = v$, $p_i = p$ and $K_i(x, t) = 1$ ($i = 1, 2$), that is

$$\begin{cases} u_t = \Delta u + u^p, & x \in \mathbf{R}^N, t > 0, \\ u(x, 0) = u_0(x) \geq 0, & x \in \mathbf{R}^N, \end{cases} \quad (13)$$

was studied by Fujita [3]. Fujita proved that when $p < 1 + 2/N$ the solution of (13) is not global in time for any nonnegative bounded and continuous initial data $u_0 \not\equiv 0$. Fujita's results were also extended by some researcher. Hayakawa [8], Kobayashi-Sirao-Tanaka [11] and Weissler [28] proved that

when $p = 1 + 2/N$, the solution of (13) blows up in finite time for any $u_0 \not\equiv 0$. For the case $p > 1 + 2/N$, Lee-Ni [12] studied that if $\|u_0\|_\infty$ is large enough or $\liminf_{|x| \rightarrow \infty} |x|^a u_0(x) > 0$ with $a < 2/(p-1)$, the solution of (13) is not global in time. When D is a cone, that is

$$\begin{cases} u_t = \Delta u + u^p, & x \in D, t > 0, \\ u(x, t) = 0, & x \in \partial D, t > 0, \\ u(x, 0) = u_0(x) \geq 0, & x \in D, \end{cases} \quad (14)$$

Levine-Meier [14], [15] proved that if $p \leq 1 + 2/(N + \gamma_+)$, there is no global solution of (14).

Fujita's results were extended to the case $D = \mathbf{R}^N$, $u = v$, $p_i = p$ and $K_i(x, t) = K(x, t)$ for $i = 1, 2$, that is

$$\begin{cases} u_t = \Delta u + K(x, t)u^p, & x \in \mathbf{R}^N, t > 0, \\ u(x, 0) = u_0(x) \geq 0, & x \in \mathbf{R}^N. \end{cases} \quad (15)$$

In the case $K(x, t) = |x|^\sigma$ with $\sigma \geq 0$, Bandle-Levine [1] had that when $p < 1 + (2 + \sigma)/N$ the solution of (15) is not global in time for any $u_0 \not\equiv 0$. Hamada [6] had the same result for $p = 1 + (2 + \sigma)/N$ (see also [18]). Suzuki [23] extended to the case $\sigma \in \mathbf{R}$ for the quasilinear parabolic equations. Thereafter, Qi [20] extended the result to the case $K(x, t) = t^q |x|^\sigma u^p$ with $q \geq 0$, $\sigma \geq 0$. He proved that when $p \leq 1 + (2 + \sigma + 2q)/N$ there exists no global solution of (15). When D is a cone, that is

$$\begin{cases} u_t = \Delta u + K(x, t)u^p, & x \in D, t > 0, \\ u(x, t) = 0, & x \in \partial D, t > 0, \\ u(x, 0) = u_0(x) \geq 0, & x \in D, \end{cases} \quad (16)$$

in the case $K(x, t) = |x|^\sigma$ with $\sigma \geq 0$, Levine-Meier [14], [15] and Hamada [6] had that if $p \leq 1 + (2 + \sigma)/(N + \gamma_+)$, there is no global solution of (16). For the case $p > 1 + (2 + \sigma)/(N + \gamma_+)$, Hamada [7] studied that if $u_0(x) \geq M \langle x \rangle^{-a} \psi_1(x/|x|)$ with $a \leq (2 + \sigma)/(p-1)$, $0 \leq \sigma \leq (p-1)(N-2)$ and some $M > 0$, the solution of (16) is not global. In the case $K(x, t) \sim t^q$ with $q > -1$ as $t \rightarrow \infty$, Levine-Meier [15] had that if $p \leq 1 + (2 + 2q)/(N + \gamma_+)$, there exists no global solution of (16).

In the case $D = \mathbf{R}^N$, our results are reduced to Escobedo-Herrero [2] and Mochizuki [16] with $K_i(x, t) = 1$ ($i = 1, 2$), to Uda [24] with $K_i(x, t) = t^{q_i}$ ($i = 1, 2$), and to Mochizuki-Huang [17] with $K_i(x, t) = |x|^{\sigma_i}$ with $\sigma_i \in [0, n(p_i - 1))$ ($i = 1, 2$). Additionally, Guedda-Kirane [5] and Kirane-Qafsaoui [10] studied in this field. They studied the case $K_i(x, t) \sim t^{q_i} |x|^{\sigma_i}$ as $t \rightarrow \infty$ and $|x| \rightarrow \infty$. However, they needed the condition $\max\{2q_i, \sigma_i\} < n(p_i - 1)$

($i = 1, 2$). Moreover, when $K_i(x, t)$ ($i = 1, 2$) satisfy (6) and (11) with $D = \mathbf{R}^N$, the system (1) was studied by Igarashi-Umeda [9]. When D is a cone, in the case $K_i(x, t) = 1$, the condition (i) of Theorem 2 is reduced to Levine [13].

The rest of the paper is organized as follows. In section 2, we state the proof of the local existence (Theorem 1). The proof of global nonexistence (Theorem 2) is given in section 3. For the change of variable as (3), we decide $\eta(x, y, t) = \eta(r, \theta, \rho, \phi, t)$, $\eta(x, t) = \eta(r, \theta, t)$ or $\eta_0(x) = \eta_0(r, \theta)$ for any function.

2 Local existence in time

In this section we use $\sigma_i = \hat{\sigma}_i$ and $q_i = \hat{q}_i$ for $i = 1, 2$. In order to show the local solvability of the Cauchy problem (1), we consider the associated integral system

$$u(x, t) = S(t)u_0(x) + \int_0^t S(t-s)K_1(x, s)v(x, s)^{p_1} ds, \quad (17)$$

$$v(x, t) = S(t)v_0(x) + \int_0^t S(t-s)K_2(x, s)u(x, s)^{p_2} ds, \quad (18)$$

where

$$S(t)\xi(x) = \int_D G(x, y, t)\xi(y) dy \quad (19)$$

with G defined by (4). Define

$$\Psi(u, v) = (S(t)u_0(x) + \Phi_1(v), S(t)v_0(x) + \Phi_2(u)), \quad (20)$$

where

$$(\Phi_1(v), \Phi_2(u)) = \int_0^t S(t-s)(K_1(x, s)v(x, s)^{p_1}, K_2(x, s)u(x, s)^{p_2}) ds.$$

Lemma 2.1. *Let $\delta \geq 0$ and $\alpha := \max\{0, -\delta(N - 2 - \delta)/2\}$. If we take $0 < T \leq (\log 2)/\alpha$, then for $0 \leq t < T$*

$$\|S(t)\xi\|_{\infty, \delta} \leq 2\|\xi\|_{\infty, \delta}.$$

Moreover for $0 \leq t < T$

$$\|S(t)\langle \cdot \rangle^{-\delta}\|_{\infty, \delta} \leq 2.$$

Proof. Let $w(x, t) := S(t)\xi(x) - \|\xi\|_{\infty, \delta} \langle x \rangle^{-\delta} \exp(\alpha t)$, then we have

$$\begin{aligned} \Delta w - w_t &= [\alpha|x|^4 + \{2\alpha + \delta(N - 2 - \delta)\}|x|^2 + N\delta + \alpha] \|\xi\|_{\infty, \delta} \langle x \rangle^{-\delta-4} \exp(\alpha t) \geq 0. \end{aligned}$$

Combining this with Protter-Weinberger [19, Theorem 10, p.p.183-184], we get $w(x, t) \leq 0$; that is,

$$\langle x \rangle^\delta S(t)\xi(x) \leq \|\xi\|_{\infty, \delta} \exp(\alpha t).$$

Then we obtain $\|S(t)\xi\|_{\infty, \delta} \leq \|\xi\|_{\infty, \delta} \exp(\alpha t)$. Moreover, if we take $0 < T \leq (\log 2)/\alpha$, then for $0 < t < T$

$$\|S(t)\xi\|_{\infty, \delta} \leq 2\|\xi\|_{\infty, \delta}.$$

□

Lemma 2.2. (i) Assume that $\langle x \rangle^{\delta_1} u_0(x)$ and $\langle x \rangle^{\delta_2} v_0(x)$ are bounded in \bar{D} . $(S(\cdot)u_0, S(\cdot)v_0) \in E_T$ for $0 < T \leq (\log 2)/\alpha$, and we have

$$\|(S(\cdot)u_0, S(\cdot)v_0)\|_{E_T} \leq 2\{\|u_0\|_{\infty, \delta_1} + \|v_0\|_{\infty, \delta_2}\},$$

where E_T is defined in (7).

(ii) Let $(u, v) \in E_T$. Suppose that $K_i(x, t)$ ($i = 1, 2$) satisfy (6). Then $(\Phi_1(v), \Phi_2(u)) \in E_T$ for some $T > 0$, and we have

$$\|(\Phi_1(v), \Phi_2(u))\|_{E_T} \leq 2C_U(\tilde{T}_1(T) + \tilde{T}_2(T))\{\|(0, v)\|_{E_T}^{p_1} + \|(u, 0)\|_{E_T}^{p_2}\},$$

where the constant C_U is appeared in (6), and $\tilde{T}_i(t) = \{(t+1)^{q_i+1} - 1\}/(q_i+1)$ ($i = 1, 2$).

Proof. (i) It is obvious from Lemma 2.1 with $\delta = \delta_i$ ($i = 1, 2$).

(ii) Note that

$$\begin{aligned} &\int_0^t S(t-s)K_1(x, s)v(x, s)^{p_1} ds \\ &\leq \int_0^t S(t-s)C_U(s+1)^{q_1} \langle x \rangle^{\sigma_1 - \delta_2 p_1} ds \sup_{s \in [0, t]} \|v(\cdot, s)\|_{\infty, \delta_2}^{p_1}. \end{aligned}$$

A simple calculation gives $-\sigma_1 + \delta_2 p_1 = \delta_1$. Then it follows from Lemma 2.1 that

$$\|S(t-s)\langle \cdot \rangle^{\sigma_1 - \delta_2 p_1}\|_{\infty, \delta_1} \leq 2.$$

Thus we have

$$\|\Phi_1(v)\|_{\infty, \delta_1} \leq 2C_U \tilde{T}_1(t) \sup_{s \in [0, t]} \|v(s)\|_{\infty, \delta_2}^{p_1}.$$

Similarly, we have

$$\|\Phi_2(u)\|_{\infty, \delta_2} \leq 2C_U \tilde{T}_2(t) \sup_{s \in [0, t]} \|u(s)\|_{\infty, \delta_1}^{p_2}.$$

We conclude from these inequations. \square

Proof of Theorem 1.

Let $B_R = \{(u, v) \in E_T; \|(u, v)\|_{E_T} \leq R\}$ and $P_T = \{(u, v) \in E_T; u \geq 0, v \geq 0\}$, and define \tilde{T}_i same as in Lemma 2.2 (ii). For $(u_1, v_1), (u_2, v_2) \in B_R \cap P_T$ with $R \geq 1$ sufficient large, we have

$$\|\Psi(u_1, v_1) - \Psi(u_2, v_2)\|_{E_T} = \|(\Phi_1(v_1) - \Phi_1(v_2), \Phi_2(u_1) - \Phi_2(u_2))\|_{E_T}. \quad (21)$$

We consider

$$\begin{aligned} & |\Phi_1(v_1) - \Phi_1(v_2)| \langle x \rangle^{\delta_1} \\ & \leq \int_0^t S(t-s) C_U (s+1)^{q_1} \langle x \rangle^{\sigma_1} |v_1(x, s)^{p_1} - v_2(x, s)^{p_1}| ds \langle x \rangle^{\delta_1}. \end{aligned}$$

Then, since $(u_2, v_2) \in B_R$ we obtain

$$|\Phi_1(v_1) - \Phi_1(v_2)| \langle x \rangle^{\delta_1} \leq 2^{p_1} C_U \tilde{T}_1(T) \sup_{s \in [0, t]} \|R^{p_1-1} p_1(v_1(\cdot, s) - v_2(\cdot, s))\|_{\infty, \delta_2}. \quad (22)$$

By the same argument we have

$$|\Phi_2(u_1) - \Phi_2(u_2)| \langle x \rangle^{\delta_2} \leq 2^{p_2} C_U \tilde{T}_2(T) \sup_{s \in [0, t]} \|R^{p_2-1} p_2(u_1(\cdot, s) - u_2(\cdot, s))\|_{\infty, \delta_1}. \quad (23)$$

Substitute (22) and (23) into (21). Since $\max\{p_1, p_2\} \leq p_1 p_2$, we obtain

$$\begin{aligned} & \|\Psi(u_1, v_1) - \Psi(u_2, v_2)\|_{E_T} \\ & \leq 2^{p_1 p_2} C_U (\tilde{T}_1(T) + \tilde{T}_2(T)) R^{p_1 p_2 - 1} p_1 p_2 \|(u_1 - u_2, v_1 - v_2)\|_{E_T}. \end{aligned}$$

Taking $T > 0$ small enough, we have

$$\|\Psi(u_1, v_1) - \Psi(u_2, v_2)\|_{E_T} \leq \rho \|(u_1, v_1) - (u_2, v_2)\|_{E_T}$$

for some $\rho < 1$. Then Ψ is a strict contraction of $B_R \cap P_T$ into itself, whence there exists a unique fixed point $(u, v) \in B_R \cap P_T$ which solves (1). \square

3 Nonexistence of global solution

In this section we treat the nonexistence of global solutions in time of (1). Here, we take the same strategy as in [17], [18], [25] and [26]. Let $\sigma_i = \check{\sigma}_i$ and $q_i = \check{q}_i$ for $i = 1, 2$ through this section.

First, we should consider only the case $k \in (0, 1/2)$ by comparison. Let $\lambda_m > 0$ denote the principal eigenvalue of $-\Delta$ with Dirichlet problem in $B(x_m, k|x_m|)$, and let $\zeta_m(x) > 0$ denote the corresponding positive eigenfunction, normalized by $\int_{B(x_m, k|x_m|)} \zeta_m(x) dx = 1$. Define

$$F_m(t) = \int_{B(x_m, k|x_m|)} u(x, t) \zeta_m(x) dx, \quad G_m(t) = \int_{B(x_m, k|x_m|)} v(x, t) \zeta_m(x) dx. \quad (24)$$

We will show that for an appropriate choice of k , $(F_m(t), G_m(t))$ is not global in time, thereby contradicting the assumption that (u, v) is a global solution. By the same arguments as in [9, §3], [13] and [21], we have the following proposition:

Proposition 3.1. *Let $(F_m(t), G_m(t))$ by (24) for some $t_0 \in (0, t]$ and $m \in \mathbf{N}$. If*

$$F_m(c_1|x_m|^2) > A|x_m|^{-\alpha_1} \quad \text{or} \quad G_m(c_1|x_m|^2) > B|x_m|^{-\alpha_2}$$

with some $A, B > 0$ and some $c_1 > 0$, then $(F_m(t), G_m(t))$ is not global in time.

Lemma 3.1. *Let u_0 and v_0 are BC and $(u_0, v_0) \neq 0$, and let (u, v) be a solution of (1). Then for any $\tau > 0$ and $x \in D$ there exist constants $\mu \geq 1$ and $C = C(N, \tau, u_0, v_0, K_1, K_2, p_1, p_2, \mu) > 0$ such that*

$$u(x, \tau) \geq C|x|^{\gamma_+} e^{-\mu|x|^2} \psi_1(x/|x|) \quad \text{and} \quad v(x, \tau) \geq C|x|^{\gamma_+} e^{-\mu|x|^2} \psi_1(x/|x|).$$

Proof. We may let $u_0(x) \neq 0$ without loss of generality. Since $u(x, t) \geq S(t)u_0(x)$, $I_\nu(z) \geq Cz^\nu$ and $\gamma_+ = \nu_1 - (N - 2)/2$, we obtain

$$\begin{aligned} u(x, t) &\geq \frac{C}{(2t)^{1+\nu_1}} r^{\gamma_+} e^{-r^2/4t} \psi_1(\theta) \\ &\quad \times \int_0^\infty \int_\Omega \rho^{\gamma_++N-1} e^{-\rho^2/4t} \psi_1(\phi) u_0(\rho, \phi) d\phi d\rho. \end{aligned}$$

Then we have, for every $\tau_1 > 0$,

$$u(x, \tau_1) \geq C_1 r^{\gamma_+} e^{-\mu_1 r^2} \psi_1(\theta) \quad (25)$$

with $\mu_1 = \max\{1, 1/4\tau_1\}$ and

$$C_1 = C_1(\tau_1, N, u_0) = \frac{C}{(2\tau_1)^{1+\nu_1}} \int_0^\infty \int_\Omega \rho^{\gamma_++N-1} e^{-\rho^2/4\tau_1} \psi_1(\phi) u_0(\rho, \phi) d\phi d\rho.$$

From (18) and the fact that $I_\nu \geq Cz^\nu$ with some $C > 0$, we have

$$\begin{aligned} v(x, t) &\geq Cr^{\gamma_+} \psi_1(\theta) \int_0^t \int_0^\infty \int_\Omega \frac{1}{(2(t-s))^{\nu_1+1}} e^{-r^2/4(t-s)} \\ &\quad \times \rho^{\gamma_++N-1} e^{-\rho^2/4(t-s)} \psi_1(\phi) K_2(\rho, \phi, s) u^{p_2}(\rho, \phi, s) d\phi d\rho ds. \end{aligned}$$

Then by (25) we obtain for $\tau_2 > 2\tau_1$

$$\begin{aligned} v(x, \tau_2) &\geq C_2 r^{\gamma_+} \psi_1(\theta) \int_{\tau_2/2}^{\tau_2} \frac{1}{(2(\tau_2-s))^{\nu_1+1}} e^{-r^2/4(\tau_2-s)} ds \\ &\geq C_2 r^{\gamma_+} \psi_1(\theta) \frac{1}{\tau_2^{\nu_1+1}} e^{-r^2/2\tau_2} \int_{\tau_2/2}^{\tau_2} ds = C_2 r^{\gamma_+} \psi_1(\theta) \frac{1}{2\tau_2^{\nu_1}} e^{-r^2/2\tau_2} \end{aligned}$$

with

$$\begin{aligned} C_2 &= C_2(\tau_2, N, u_0, K_2) \\ &= \inf_{s \in (\tau_2/2, \tau_2)} C \int_0^\infty \int_\Omega \rho^{\gamma_++N-1} e^{-\rho^2/4(\tau_2-s)} \psi_1(\phi) K_2(\rho, \phi, s) u^{p_2}(\rho, \phi, s) d\phi d\rho. \end{aligned}$$

Then we have

$$v(x, \tau_2) \geq C_3 r^{\gamma_+} e^{-\mu_2 r^2} \psi_1(\theta)$$

with $\mu_2 = \max\{1, 1/2\tau_2\}$ and $C_3 = C_2/2\tau_2^{\nu_1}$. Put $C = \min\{C_1, C_2, C_3\}$ and $\mu = \max\{\mu_1, \mu_2\}$ and $\tau = \tau_2$. Then we have

$$u(x, \tau) \geq Cr^{\gamma_+} e^{-\mu r^2} \psi_1(\theta) \quad \text{and} \quad v(x, \tau) \geq Cr^{\gamma_+} e^{-\mu r^2} \psi_1(\theta).$$

□

Lemma 3.2. For $\sigma \geq 0$, $\mu \geq 1$, $x \in D$ and $t \geq \tau$ with some $\tau > 0$, we have

$$S(t) \chi_B(x) |x|^\sigma e^{-\mu|x|^2} \geq Ct^{(\sigma-\gamma_+)/2} (1+4\mu t)^{-(N+\sigma+\gamma_+)/2} |x|^{\gamma_+} e^{-|x|^2/4t} \psi_1(x/|x|)$$

with some $C > 0$ and $B = B(b, a) \subset D$ with $a > 0$ and $b \in D$, where χ_B is a characteristic function of B such that $\chi_B(x) = 1$ for $x \in B$ and $= 0$ for $x \in D \setminus B$. The domain $B(b, a)$ denotes the open ball of radius a centered at b .

Proof. We can put positive constants a_1, a_2 and domain $\Omega' \subset \Omega$ satisfying $0 < a_1 < a_2 < \infty$, $|\Omega'| \neq 0$ and $D_B = \{x; |x| \in (a_1, a_2), x/|x| \in \Omega'\} \subset B$. By (19) and $\int_{\Omega'} \psi_1(\phi) d\phi = C$ with some $C \in (0, 1]$, we have

$$\begin{aligned} S(t)\chi_B(x)|x|^\sigma e^{-\nu|x|^2} &\geq \int_{a_1}^{a_2} \int_{\Omega'} G(r, \theta, \rho, \phi, t) \rho^\sigma e^{-\mu\rho^2} \rho^{N-1} d\phi d\rho \\ &\geq \frac{C}{(2t)^{1+\nu_1}} r^{\gamma_+} e^{-r^2/4t} \psi_1(\theta) \int_{a_1}^{a_2} \int_{\Omega'} \rho^{\gamma_++\sigma+N-1} e^{-(1+4\mu t)\rho^2/4t} \psi_1(\phi) d\phi d\rho \\ &\geq \frac{Cr^{\gamma_+} e^{-r^2/4t} \psi_1(\theta)}{(2t)^{1+\nu_1} \tilde{\mu}(t)^{\gamma_++\sigma+N}} \int_{\tilde{\mu}(t)a_1}^{\tilde{\mu}(t)a_2} s^{\gamma_++\sigma+N-1} e^{-s^2} ds \end{aligned}$$

where $\tilde{\mu}(t) = \sqrt{(1+4\mu t)/4t}$. Since $1 \leq \sqrt{\mu} \leq \tilde{\mu}(t) \leq \tilde{\mu}(\tau)$ for $t \geq \tau$, we have

$$S(t)\chi_B(x)|x|^\sigma e^{-\nu|x|^2} \geq Ct^{(\sigma-\gamma_+)/2} (1+4\mu t)^{-(N+\sigma+\gamma_+)/2} r^{\gamma_+} e^{-r^2/4t} \psi_1(\theta).$$

□

By Lemma 3.1, we can assume

$$u_0(x) \geq C|x|^{\gamma_+} e^{-\mu|x|^2} \psi_1(x/|x|)$$

for some $C > 0$ and $\mu > 0$. Then we have, for $t \geq \tau$

$$u(x, t) \geq C(1+4\mu t)^{-N/2-\gamma_+} |x|^{\gamma_+} e^{-|x|^2/4t} \psi_1(x/|x|). \quad (26)$$

Lemma 3.3. *Let v be the second element of the solution of (1). Then we have*

$$v(x, t) \geq Ct^{((p_2-1)\gamma_++\sigma_2+2q_2+2)/2} (t+1)^{-\gamma_++p_2-Np_2/2} |x|^{\gamma_+} e^{-|x|^2/2t} \psi(x/|x|)^{p_2+1}$$

for $t \geq \tau$ with some $\tau > 0$ and $C = C(\tau, u_0, v_0, K_1, K_2, p_1, p_2) > 0$.

Proof. It follows from (11), (18) and (26), we obtain

$$\begin{aligned} v(x, t) &\geq C \int_0^t S(t-s) \chi_{\tilde{B}_{k,1}}(x) |x|^{\sigma_2+p_2\gamma_+} s^{q_2} \\ &\quad \times (4s+1/\mu)^{-Np_2/2-p_2\gamma_+} e^{-p_2|x|^2/4s} \psi_1^{p_2}(x/|x|) ds. \end{aligned}$$

By Lemma 3.2, we then have

$$\begin{aligned} v(x, t) &\geq C(t/2)^{(p_2-1)\gamma_+/2+\sigma_2/2} (t/4)^{q_2} (2t+1/\mu)^{-\gamma_++p_2-Np_2/2} \\ &\quad \times |x|^{\gamma_+} e^{-|x|^2/2t} \psi_1^{p_2+1}(x/|x|) \int_{t/4}^{t/2} ds. \end{aligned}$$

Thus, the inequality of the lemma holds. □

Lemma 3.4. *Let u be first element of the solution of (1) and $\alpha_1 \geq N + \gamma_+$. Then for $t \geq a$*

$u(x, t)$

$$\geq \begin{cases} Ct^{-N/2-\gamma_+}|x|^{\gamma_+}e^{-|x|^2/2t}\psi_1(x/|x|)^{p_1p_2+p_1+1}\log(t/2a), & \text{if } \alpha_1 = N + \gamma_+, \\ Ct^{-N/2-\gamma_+}|x|^{\gamma_+}e^{-|x|^2/2t}\psi_1(x/|x|)^{p_1p_2+p_1+1}(t^{\tilde{p}} - (2a)^{\tilde{p}}), & \text{if } \alpha_1 > N + \gamma_+ \end{cases}$$

with $C = C(a, u_0, v_0, K_1, K_2, p_1, p_2, N) > 0$, where $a > 0$ is a small constant and $\tilde{p} = (p_1p_2 - 1)(\alpha_1 - N - \gamma_+)/2$.

Proof. By Lemma 3.3, we have

$$\begin{aligned} u(x, t) &\geq C \int_a^t S(t-s)\chi_{\bar{B}_{k,1}}(x)|x|^{\sigma_1}S^{q_1}S^{(1+\sigma_2/2+q_2)p_1+p_1(p_2-1)\gamma_+/2} \\ &\quad \times (s+1)^{-Np_1p_2/2-\gamma_+p_1p_2}|x|^{p_1\gamma_+}e^{-p_1|x|^2/2s}\psi_1(x/|x|)^{p_1(p_2+1)}ds. \end{aligned}$$

It follows from Lemma 3.2 that

$$\begin{aligned} u(x, t) &\geq C(t/2)^{(\sigma_1+p_1\gamma_+-\gamma_+)/2}t^{-(N+\sigma_1+p_1\gamma_++\gamma_+)/2}|x|^{\gamma_+}e^{-|x|^2/2t} \\ &\quad \times \psi_1(x/|x|)^{p_1p_2+p_1+1} \int_a^{t/2} s^{\{(-N-\gamma_+)(p_1p_2-1)+(2+\sigma_2+2q_2)p_1+\sigma_1+2q_1\}/2}ds \end{aligned}$$

for small $a > 0$. Since

$$\begin{aligned} &\{(-N-\gamma_+)(p_1p_2-1)+(2+\sigma_2+2q_2)p_1+\sigma_1+2q_1\}/2 \\ &= \{(p_1p_2-1)(\alpha_1-N-\gamma_+)\}/2-1, \end{aligned}$$

this proves the inequality of the lemma. \square

Proof of Theorem 2.

First we consider the case (i). We may assume $\alpha_1 \geq \alpha_2$. Put $Y_m = \sqrt{c_1}|x_m|$. From the definition, we have $\alpha_1 \geq N + \gamma_+$. By Lemma 3.4, since $x \in B(x_m, k|x_m|)$, we have

$$\begin{aligned} F_m(Y_m^2) &\geq CY_m^{-N-\gamma_+}h_m \\ &\quad \times \int_{B(x_m, k|x_m|)} \frac{|x|^{\gamma_+}}{|x_m|^{\gamma_+}} \exp\left(-\frac{|x|^2}{2Y_m^2}\right) \zeta_m(x)\psi_1(x/|x|)^{p_1p_2+p_1+1}dx \\ &\geq C|x_m|^{-N-\gamma_+}(c_1)^{-N/2-\gamma_+}h_m(1+k)^{\gamma_+} \exp(-(1+k)^2c_1/2), \end{aligned}$$

where $h_m = \log(Y_m^2/2a)$ for $\alpha_1 = N + \gamma_+$ and $h_m = Y_m^{2\tilde{p}} - (2a)^{\tilde{p}}$ for $\alpha_1 > N + \gamma_+$ with $C = C(a, u_0, v_0, K_1, K_2, p_1, p_2, N) > 0$ and \tilde{p} defined in Lemma 3.4. Since $\alpha_1 \geq N + \gamma_+$, it follows that

$$\begin{aligned} |x_m|^{\alpha_1}F_m(Y_m^2) &\geq C|x_m|^{\alpha_1-N-\gamma_+}c_1^{-N/2-\gamma_+} \\ &\quad \times h_m(1+k)^{\gamma_+} \exp(-(1+k)^2c_1/2) > A \end{aligned}$$

for m large enough. Thus, $(F_m(t), G_m(t))$ is not global in time by Proposition 3.1.

Next, we consider the case (ii). Since $u(x, t) \geq S(t)u_0(x)$, $u \in H_{a_1}$, $I_\nu(z) \geq Cz^\nu$ and $\int_\Omega \psi_1(\phi)^2 d\phi$ is constant, it follows that

$$\begin{aligned} u(x, t) &\geq \int_0^\infty \int_\Omega G(r, \theta, \rho, \phi, t) u_0(\rho, \phi) \rho^{N-1} d\phi d\rho \\ &\geq C \int_0^\infty \frac{r^{\gamma_+} \rho^{N+\gamma_+-1}}{t^{\gamma_++N/2}} \exp\left(-\frac{r^2 + \rho^2}{4t}\right) (1 + \rho^2)^{-a_1/2} \psi_1(\theta) d\rho. \end{aligned}$$

Then, since $\gamma_+ = \nu_1 - (N - 2)/2$, we obtain

$$\begin{aligned} u(x, Y_m^2) &\geq C \left(\frac{r}{Y_m}\right)^{\gamma_+} \exp\left(-\frac{r^2}{4Y_m^2}\right) \int_0^\infty \frac{1}{Y_m} \left(\frac{\rho}{Y_m}\right)^{\frac{N}{2}+\nu_1} \\ &\quad \times \exp\left(-\frac{\rho^2}{4Y_m^2}\right) (1 + \rho^2)^{-a_1/2} \psi_1(\theta) d\rho. \end{aligned}$$

Since $Y_m = \sqrt{c_1|x_m|^2}$, we have for $x \in B(x_m, k|x_m|)$

$$\begin{aligned} u(x, Y_m^2) &\geq C(1+k)^{-\frac{N}{2}}(1-k)^{1+\nu_1} c_1^{-\gamma_+/2} \exp\left\{-\frac{(1+k)^2}{c_1}\right\} \\ &\quad \times \psi_1(\theta) \int_0^\infty \frac{1}{Y_m} \left(\frac{\rho}{Y_m}\right)^{\frac{N}{2}+\nu_1} \exp\left(-\frac{\rho^2}{4Y_m^2}\right) (1 + \rho^2)^{-a_1/2} d\rho. \end{aligned}$$

Putting $\chi = \rho/Y_m$, we have

$$u(x, Y_m^2) \geq C\psi_1(\theta) \int_0^\infty \chi^{\frac{N}{2}+\nu_1} \exp\left(-\frac{\chi^2}{4}\right) (1 + \chi^2 Y_m^2)^{-a_1/2} d\chi.$$

Note that $1 + \chi^2 Y_m^2 \leq Y_m^2(1 + \chi^2)$ if m is large enough. Then, we obtain

$$\begin{aligned} u(x, Y_m^2) &\geq C\psi_1(\theta) |Y_m|^{-a_1} \int_0^\infty \chi^{\frac{N}{2}+\nu_1} \exp\left(-\frac{\chi^2}{4}\right) (1 + \chi^2)^{-a_1/2} d\chi \\ &\geq C\psi_1(x/|x|) |x_m|^{-a_1} \end{aligned}$$

for sufficiently large m . Since $\int_{B(x_m, k|x_m|)} \psi_1(x/|x|) \zeta_m(x) dx$ is constant, we have

$$\begin{aligned} F_m(Y_m^2) &\geq \int_{B(x_m, k|x_m|)} u(x, Y_m^2) \zeta_m(x) dx \\ &\geq C|x_m|^{-a_1} \int_{B(x_m, k|x_m|)} \psi_1(x/|x|) \zeta_m(x) dx \geq C|x_m|^{-a_1}. \end{aligned}$$

Since $u_0 \in H_{a_1}$ with $a_1 < \alpha_1$, we have

$$|x_m|^{\alpha_1} F_m(Y_m^2) \geq C|x_m|^{\alpha_1 - a_1} > A$$

for sufficiently large m . If $v_0 \in H_{a_2}$ with $a_2 < \alpha_2$, we similarly have

$$|x_m|^{\alpha_2} G_m(Y_m^2) \geq C|x_m|^{\alpha_2 - a_2} > B$$

for m large enough. Thus, $(F_m(t), G_m(t))$ is not global in time by Proposition 3.1. \square

Acknowledgement. Much of the work of the second author was done while he visited the University of Tokyo during 2005-2008 as a postdoctoral fellow. Its hospitality is gratefully acknowledged as well as support from formation of COE “New Mathematical Development Center to Support Scientific Technology” (University of Tokyo), supported by JSPS.

References

- [1] C. Bandle and H. A. Levine, *On the existence and nonexistence of global solution of reaction-diffusion equation in sectorial domains*, Trans. Amar. Math. Sec. **316** (1989), 595–622.
- [2] M. Escobedo and M. A. Herrero, *Boundness and blow up for a semilinear reaction-diffusion system*, J. Diff. Eqns. **89** (1991), 176–202.
- [3] H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Univ. Tokyo Sect. A Math. **16** (1966), 109–124.
- [4] Y. Giga and N. Umeda, *Blow-up directions at space infinity for solutions of semilinear heat equations*, Bol. Soc. Parana. Mat. **23** (2005), 9–28.
- [5] M. Guedda and M. Kirane, *Criticality for some evolution equations*, Differential Equations **37** (2001), 540–550.
- [6] T. Hamada, *Nonexistence of global solutions of parabolic equations in conical domains*, Tsukuba J. Math. **19** (1995), 15-25.
- [7] T. Hamada, *On the existence and nonexistence of global solutions of semilinear parabolic equations with slowly decaying initial data*, Tsukuba J. Math. **21** (1997), 505-514.
- [8] K. Hayakawa, *On nonexistence of global solution of some semilinear parabolic equations*, Proc. Japan. Acad. **49** (1973), 503–505.

- [9] T. Igarashi and N. Umeda, *Existence and nonexistence of global solutions in time for a reaction-diffusion system with inhomogeneous terms*, Funkcialaj Ekvacioj **51** (2008), 17–37.
- [10] M. Kirane and M. Qafsaoui, *Global nonexistence for the Cauchy problem of some nonlinear reaction-diffusion systems*, J. Math. Anal. Appl. **268** (2002), 217-243.
- [11] K. Kobayashi, T. Sirao and H. Tanaka, *On glowing up problem for semilinear heat equations*, J. Math. Soc. Japan **29** (1977), 407-424.
- [12] T.-Y. Lee and W.-M. Ni, *Global existence, large time behavior and life span on solutions of semilinear Cauchy problem*, Trans. Amer. Math. Soc. **333** (1992), 365-378.
- [13] H. A. Levine, *A Fujita type global existence-global nonexistence theorem for a weakly coupled system of reaction-diffusion equations*, J. Appl. Math. Phys. (ZAMP) **42** (1991), 408-430.
- [14] H. A. Levine and P. Meier, *The value of critical exponent for reaction-diffusion equation in cones*, Arch. Ratl. Mech. Anal. **109** (1990), 73-80.
- [15] H. A. Levine and P. Meier, *A blowup result for the critical exponent in cones*, Israel J. Math. **67** (1989), 129-136.
- [16] K. Mochizuki, *Blow-up, life-span and large time behavior of solutions of a weakly coupled system of reaction-diffusion equations*, Adv. Math. Appl. Sci. **48**, World Scientific 1998, 175-198.
- [17] K. Mochizuki and Q. Huang, *Existence and behavior of solutions for a weakly coupled system of reaction-diffusion equations*, Methods Appl. Anal. **5** (2) (1998), 109-124.
- [18] R. G. Pinsky, *Existence and nonexistence of global solutions for $u_t = \Delta u + a(x)u^p$ in \mathbf{R}^n* , J. Differential Equations **133** (1997), 152-177.
- [19] M. H. Protter and H. F. Weinberger, “Maximum principles in Differential Equations”, Prentice-Hall, Englewood Cliffs, New Jersey, 1967.
- [20] Y.-W. Qi, *The critical exponents of parabolic equations and blow-up in \mathbf{R}^n* , Proc. Roy. Soc. Edinburgh Sect. **128A** (1998), 123-136.
- [21] Y.-W. Qi and H. A. Levine, *The critical exponent of degenerate parabolic systems*, Z. Angew. Math. Phys. **44** (1993), 249-265.

- [22] M. Shimojyo, *On blow-up phenomenon at space infinity and its locality for semilinear heat equations (in Japanese)*, Master's Thesis, The University of Tokyo (2005).
- [23] R. Suzuki, *Existence and nonexistence of global solutions of quasilinear parabolic equations*, J. Math. Soc. Japan, **54** (2002), 747–792.
- [24] Y. Uda, *The critical exponent for a weakly coupled system of the generalized Fujita type reaction-diffusion equations*, Z. Angew Math. Phys. **46** (1995), 366–383.
- [25] N. Umeda, *Blow-up and large time behavior of solutions of a weakly coupled system of reaction-diffusion equations*, Tsukuba J. Math. **27** (2003) 31–46.
- [26] N. Umeda, *Existence and nonexistence of global solutions of a weakly coupled system of reaction-diffusion equations*, Comm. Appl. Anal. **10** (2006) 57–78.
- [27] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd Ed., Cambridge University Press, London/New York 1944.
- [28] F. B. Weissler, *Existence and nonexistence of global solutions for semilinear heat equation*, Israel J. Math. **38** (1981) 29–40.

UTMS

- 2008–7 Noriaki Umeda: *Blow-up at space infinity for nonlinear equations.*
- 2008–8 Masahiko shimojo and Noriaki Umeda: *Blow-up at space infinity for solutions of cooperative reaction-diffusion systems.*
- 2008–9 Shouhei Ma: *Fourier-Mukai partners of a K3 surface and the cusps of its Kähler Moduli.*
- 2008–10 Takashi Tsuboi: *On the uniform perfectness of diffeomorphism groups.*
- 2008–11 Takashi Tsuboi: *On the simplicity of the group of contactomorphisms.*
- 2008–12 Hajime Fujita, Mikio Furuta and Takahiko Yoshida: *Acyclic polarizations and localization of Riemann-Roch numbers I.*
- 2008–13 Rolci Cipolatti and Masahiro Yamamoto: *Inverse hyperbolic problem by a finite time of observations with arbitrary initial values.*
- 2008–14 Yoshifumi Matsuda: *Groups of real analytic diffeomorphisms of the circle with a finite image under the rotation number function.*
- 2008–15 Shoichi Kaji: *The (\mathfrak{g}, K) -module structures of the principal series representations of $SL(4, R)$.*
- 2008–16 G. Bayarmagnai: *The (\mathfrak{g}, K) -module structures of principal series of $SU(2,2)$.*
- 2008–17 Takashi Tsuboi: *On the group of real analytic diffeomorphisms.*
- 2008–18 Takefumi Igarashi and Noriaki Umeda: *Nonexistence of global solutions in time for reaction-diffusion systems with inhomogeneous terms in cones.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153-8914, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012