


UTMS 2011–26

November 16, 2011

**Analysis of the fictitious domain method
with penalty for elliptic problems**

by

Guanyu ZHOU and Norikazu SAITO



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

ANALYSIS OF THE FICTITIOUS DOMAIN METHOD WITH PENALTY FOR ELLIPTIC PROBLEMS

GUANYU ZHOU AND NORIKAZU SAITO

ABSTRACT. The fictitious domain method with H^1 -penalty for elliptic problems is considered. We propose a new way to derive the sharp error estimates between the solutions of original elliptic problems and their H^1 -penalty problems. We find our method of analysis is applicable to parabolic problems while retaining the sharpness of the error estimates. We also prove some regularity theorems for H^1 -penalty problems. The $P1$ finite element approximation to H^1 -penalty problems is investigated. We study error estimates between the solutions of H^1 -penalty problems and discrete problems in H^1 norm, as well as in L^2 norm, which is not currently found in the literature. Thanks to regularity theorems, we can simplify the analysis of error estimates. Due to the integration on a curved domain, the discrete problem is not suitable for computation directly. Hence an approximation of the discrete problem is necessary. We provide an approximation scheme for the discrete problem and derive its error estimates. The validity of theoretical results is confirmed by numerical examples.

1. INTRODUCTION

The principle of the fictitious domain method is to solve the problem in a larger domain (the *fictitious domain*) containing the domain of interest with a very simple shape. Then, the fictitious domain is discretized by a uniform mesh, independent of the original boundary. The advantage of this approach is that we can avoid the time-consuming construction of a boundary-fitted mesh. One of these approaches is the penalty fictitious domain method which is based on a reformulation of the original problem in the fictitious domain by using a penalty parameter(see [2] for an introduction of other kinds of fictitious domain methods). In this article, we consider only the fictitious domain method with penalty. Obviously, the fictitious domain method is of use for time-dependent moving-boundary problems. Although there exist some ways to derive the sharp error estimates for elliptic problems(cf. [10, 15, 16]), it seems none of them has been applied to parabolic problem such that the sharpness of the error estimates are maintained. Our motivation lies in the study of the penalty fictitious domain method which can be applied to these time-dependent moving-boundary problems. This is of obvious

importance, and it seems that little is known in this direction. The fictitious domain method with penalty for parabolic problem firstly appeared in [8] to prove the existence of the solution for parabolic problem in time-dependent domain. Then, in [9], the convergence and finite difference approximation is given, but without error estimates. The H^1 -penalty parabolic problem equals to a special interface problem, and in [1] the error estimate for elliptic and parabolic interface problem is studied. However, it is not so suitable to the H^1 -penalty problem, and still, is only for time-independent domain when considering parabolic interface problem. As a primary step towards this final end, herein we examine some new methods of error analysis for elliptic problems that can be easily applied to parabolic problems while maintaining the sharpness of the error estimates. This is the purpose of this paper.

In order to illustrate our results, we consider the Dirichlet boundary value problem for the Poisson equation. The weak form (Q) reads as

$$(1.1) \quad \begin{cases} \text{Find } u \in H_0^1(\Omega) \text{ such that} \\ (\nabla u, \nabla v)_\Omega = (f, v)_\Omega, \quad \forall v \in H_0^1(\Omega), \end{cases}$$

where $\Omega \subset \mathbb{R}^2$ denotes a smooth bounded domain, $(\cdot, \cdot)_\Omega$ is the inner product of $L^2(\Omega)$ and $f \in L^2(\Omega)$. We can find a rectangular domain $D \supset \Omega$, $\Omega_1 = D \setminus \bar{\Omega}$, and turn to solve the H^1 -penalty problem (Q_ϵ) with penalty coefficient $0 < \epsilon \ll 1$,

$$(1.2) \quad \begin{cases} \text{Find } u_\epsilon \in H_0^1(D) \text{ such that} \\ (\nabla u_\epsilon, \nabla v)_\Omega + \frac{1}{\epsilon}(\nabla u_\epsilon, \nabla v)_{\Omega_1} = (\tilde{f}, v)_D, \quad \forall v \in H_0^1(D), \end{cases}$$

where \tilde{f} is the zero extension of f into D .

Another example of applying the fictitious domain method with penalty to (1.1) is,

$$(\nabla u_\epsilon, \nabla v)_\Omega + \frac{1}{\epsilon}(u_\epsilon, v)_{\Omega_1} = (\tilde{f}, v), \quad \forall v \in H_0^1(D),$$

which we call the L^2 -penalty problem. This is of interest; however, the L^2 -penalty problem is beyond the scope of this paper, in which we shall concentrate our attention to the H^1 -penalty problem.

The error $\|u_\epsilon - u\|_{1,\Omega}$ ($\|\cdot\|_{1,\Omega} = \|\cdot\|_{0,\Omega} + \|\cdot\|_{1,\Omega}$) has been analyzed by many authors, where $\|\cdot\|_{1,\Omega}$ is the $H^1(\Omega)$ norm. In [9], it is bounded by $C\sqrt{\epsilon}$ (C is some constant, so as in the following), and in [10, 15, 16] the sharp estimate $C\epsilon$ is achieved. In this paper, we give a new way to derive the sharp estimate. Moreover, we present some regularity analysis of u_ϵ , which is useful for studying the H^1 and L^2 error between the solutions of H^1 problem (Q_ϵ) and its discrete problem, which is denoted as $(Q_{\epsilon,h})$.

A Cartesian mesh can be introduced to the rectangular domain D to get a uniformed triangulation \mathcal{T}_h , h is the maximum diameter of the triangles of \mathcal{T}_h .

$V_h(D)$ is the subspace of all piecewise linear continuous functions subordinate to \mathcal{T}_h . Then $(Q_{\epsilon,h})$ reads as:

$$(1.3) \quad \begin{cases} \text{Find } u_{\epsilon,h} \in V_h(D) \text{ such that} \\ (\nabla u_{\epsilon,h}, \nabla v_h)_\Omega + \frac{1}{\epsilon} (\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega_1} = (\tilde{f}, v_h)_D, \quad \forall v_h \in V_h(D). \end{cases}$$

In the literature, there are several works devoted to the study of the H^1 error between $u_{\epsilon,h}$ and u_ϵ in Ω . For example, in [16], it is proved that the H^1 error is bounded by $C(\epsilon + \epsilon\sqrt{h} + h\sqrt{\epsilon} + \sqrt{h})$. In our work, we prove a similar result with the analysis with a much simpler method of the analysis. The analysis of [16] is to consider the estimate of $u_{\epsilon,h} - u^0 - \epsilon u^1$ to derive the final estimate of $u_\epsilon - u_{\epsilon,h}$, where u^0 is the zero extension of u , and u^1 is the solution of problem:

$$-\Delta u^1 = 0 \text{ in } \Omega_1, \quad \frac{\partial u^1}{\partial n^-} = \frac{\partial u^0}{\partial n}, \quad \text{and } u^1 = 0 \text{ on } \partial D.$$

Here, n is the unit outward normal to Γ viewed as a boundary of Ω , and n^- is opposite to n . We found the analysis in [16] to be complicated and not directly. Our method for estimating of $u_\epsilon - u_{\epsilon,h}$ is simpler, which is to find some interpolation of u_ϵ , denoted as v_h , and then estimate $u_\epsilon - v_h$ by using a regularity theorem of (Q_ϵ) .

Moreover, we show the L^2 error is bounded by $C(\epsilon + h + \sqrt{\epsilon h})$. A similar result of H^1 error for the elliptic problem in a specific domain is given in [10].

In discrete problem $(Q_{\epsilon,h})$, we notice that we have to calculate the inner-product in a curved domain, for example, $(\nabla u_{\epsilon,h}, \nabla v_h)_\Omega$. So the discrete problem cannot be directly computed. We find that few prior works have provided a sufficient discussion on this issue; however, it is necessary to give an approximation scheme for solving $(Q_{\epsilon,h})$ and the associated error estimates when applying the finite element method to computation.

Herein, we present an approximation scheme, that is, instead of solving $(Q_{\epsilon,h})$ we solve some problem $(\hat{Q}_{\epsilon,h})$ approximating to $(Q_{\epsilon,h})$. $(\hat{Q}_{\epsilon,h})$ reads as:

$$(1.4) \quad \begin{cases} \text{Find } \hat{u}_{\epsilon,h} \in V_h(D) \text{ such that} \\ (\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega}} + \frac{1}{\epsilon} (\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega}_1} = (f_h, v_h)_D, \quad \forall v_h \in V_h(D), \end{cases}$$

where $\hat{\Omega}$ is a polygon approximating to Ω , with $\hat{\Omega} \subset \Omega$, f_h is some interpolation of \tilde{f} . With assumptions that $\|f_h - f\|_{0,D} \leq Ch$, $\hat{\Omega}$ is convex and $(\hat{\Omega} \setminus \Omega) \cup (\Omega \setminus \hat{\Omega}) \subset \Gamma_\delta$, $\delta = O(h^2)$, where $\Gamma_\delta = \{x \in \mathbb{R}^2 \mid \text{dist}(x, \Gamma = \partial\Omega) \leq \delta\}$, we find that the error $\hat{u}_{\epsilon,h} - u_{\epsilon,h}$ is bounded by Ch in H^1 norm.

In above, we have restricted our attention to Dirichlet boundary problem. For Neumann and mixed boundary problems we also consider the approximation of H^1 -penalty problems. Further, for Neumann boundary problem, the

discrete problem is investigated. Although some results we obtain are similar to those of [16], as we mentioned in the beginning, our work is focus on solving parabolic problems with time-dependent domain, and all methods of analysis are applicable to this class of problems.

The rest of this paper is organized as follows. The H^1 -penalty problems for original problems with Dirichlet, Neumann, and mixed boundaries are given in Section 2, as well as the analysis of error estimates between the solutions of original problems and H^1 -penalty problems, in a different way from that in [10, 15].

In Section 3, we present some regularity theorems for H^1 -penalty problems. The H^1 -penalty problem is in a sense equivalent to a kind of interface problem. The regularity theorem for the interface problem has been studied in [12]. However, we make several improvements in priori estimates and identify some higher-order regularity for our problems. The theorems will be used in Section 4 to make the error estimate more simple than that of [16].

The Section 4 is devoted to discrete problems. Finite element approximations are investigated. Using the same separation method of the triangulated domain as in [16], regularity theorems in Section 3 and some lemmas from [3, 13], we obtain the error estimates in H^1 norm with the same order as in [10, 16]. Moreover, we give the higher-order L^2 norm error estimates.

We consider a scheme approximating to the discrete problem in Section 5. We introduce a new discrete problem ($\hat{Q}_{\epsilon,h}$) to approximate to the discrete problem ($Q_{\epsilon,h}$). Of necessity, due to insufficient prior reported works on this issue in the literature, we derive some error estimates of the scheme to make the numerical analysis of the fictitious domain method with H^1 -penalty more complete.

Finally, we give some numerical experiments to verify our theoretical results in Section 6.

2. H^1 -PENALTY PROBLEMS OF FICTITIOUS DOMAIN METHOD FOR ELLIPTIC PROBLEMS

Following the notation given in the previous section, we state the H^1 -penalty problem for the original elliptic problem with homogeneous Dirichlet, Neumann and mixed boundary conditions. In addition, we write $\Gamma = \partial\Omega$.

2.1. Dirichlet boundary value problem. First, we consider the Dirichlet boundary value problem (1.1) and its H^1 -penalty problem (1.2).

Theorem 2.1. *There exist unique solutions u and u_ϵ for (1.1) and (1.2), respectively, and we have the following estimates:*

$$(2.1) \quad \|u_\epsilon - u\|_{1,\Omega} \leq C\epsilon,$$

$$(2.2) \quad \|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon.$$

Those error estimates themselves are not new: they have been stated in [10] and [15]. The main process to prove those estimates in [10] and [15] are different (see Remark 3 below). We shall give a somewhat new proof which will be used for parabolic problems. Before stating that, we recall the well-known extension and trace theorems that we frequently use.

Lemma 2.2. (Theorem 8.1 of Chapter 1 in [7]) *Let $\omega \subset \mathbb{R}^2$ be a bounded domain with the smooth boundary $\partial\omega$. Then, for any integer $k > 0$, there exists an operator $E_k(\omega) : H^k(\omega) \rightarrow H^k(\mathbb{R}^2)$ with the properties*

$$E_k(\omega)u = u \text{ a.e. on } \omega \quad (u \in H^k(\omega)),$$

$$\|E_k(\omega)u\|_{H^k(\mathbb{R}^2)} \leq C_k \|u\|_{H^k(\omega)} \quad (u \in H^k(\omega)),$$

with a domain constant $C_k > 0$.

The following lemma is a readily obtainable consequence of Theorem 8.3 of Chapter 1 in [7].

Lemma 2.3. *Let $\omega \subset \mathbb{R}^2$ be a bounded domain. Assume that the boundary $\partial\omega$ consists of two disjoint and smooth components $\partial\omega_1$ and $\partial\omega_2$; $\partial\omega = \partial\omega_1 \cup \partial\omega_2$. Then, the mapping $u \mapsto u|_{\partial\omega_1}$ of $C^\infty(\bar{\omega}) \rightarrow C^\infty(\partial\omega_1)$ is extended by continuity to a continuous linear mapping, which is called the trace operator and denoted by $\gamma(\omega, \partial\omega_1)$, of $H^1(\omega) \rightarrow H^{\frac{1}{2}}(\partial\omega_1)$. This mapping is surjective and there exists a continuous linear right inverse, which is called the lifting operator, $g \mapsto \gamma(\omega, \partial\omega_1)^{-1}g$ of $H^{\frac{1}{2}}(\partial\omega_1) \rightarrow H^1(\omega)$ such that $\gamma(\omega, \partial\omega_1)\gamma(\omega, \partial\omega_1)^{-1}g = g$.*

Remark 1. In view of Lemma 2.2, there exists an operator $E_{1,0}(\Omega) : H^1(\Omega) \rightarrow H_0^1(D)$ with properties:

$$E_{1,0}(\Omega)u = u \text{ a.e. in } \Omega \quad (u \in H^1(\Omega)),$$

$$\|E_{1,0}(\Omega)u\|_{H^1(D)} \leq C \|u\|_{H^1(\Omega)} \quad (u \in H^1(\Omega)).$$

In fact, taking $\phi \in C_0^\infty(D)$ with $0 \leq \phi \leq 1$ and $\phi = 1$ in Ω' , where $\Omega \subset \Omega' \subset D$, then the desired operator is defined as $E_{1,0}(\Omega)u = \phi \cdot E_1(\Omega)u$ for $u \in H^1(\Omega)$.

Now, we can state the following proof.

Proof of Theorem 2.1. Firstly, by the Lax-Milgram Theorem, the unique existence of u and u_ϵ is obvious. And we can obtain the estimate for the solution of the H^1 -penalty problem (1.2),

$$\|u_\epsilon\|_{1,D} \leq C \|f\|_0 \inf(1, \frac{1}{\epsilon}).$$

Without loss of generality, we assume $0 < \epsilon \leq 1$ in what follows. Substituting $v = E_{1,0}(\Omega_1)u_\epsilon \in H_0^1(D)$ into (1.2), we obtain

$$\epsilon(\nabla u_\epsilon, v)_\Omega + \|\nabla u_\epsilon\|_{0,\Omega_1}^2 = \epsilon(\tilde{f}, v)_D,$$

which leads to an estimate of $u_\epsilon|_{\Omega_1}$, in particular,

$$C_1 \|u_\epsilon\|_{1,\Omega_1}^2 \leq \|\nabla u_\epsilon\|_{0,\Omega_1}^2 \leq C\epsilon \|f\|_0 \|v\|_1 + \epsilon |u_\epsilon|_{1,\Omega} |v|_{1,\Omega}.$$

Therein, the first inequality is deduced by *Friedrichs'* inequality, and the second term of the right-hand side is bounded by $C_2\epsilon \|f\|_0 \|u_\epsilon\|_{1,\Omega_1}$. Thus we have

$$\|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon.$$

Next, we consider the trace operators

$$\gamma(\Omega, \Gamma) : H^1(\Omega) \rightarrow H^{\frac{1}{2}}(\Gamma); \quad \gamma(\Omega_1, \Gamma) : H^1(\Omega_1) \rightarrow H^{\frac{1}{2}}(\Gamma).$$

Since $u_\epsilon \in H^1(D)$, we have $\gamma(\Omega, \Gamma)u_\epsilon = \gamma(\Omega_1, \Gamma)u_\epsilon \in H^{\frac{1}{2}}(\Gamma)$. Hence,

$$\gamma(\Omega, \Gamma)(u - u_\epsilon) = \gamma(\Omega, \Gamma)u - \gamma(\Omega, \Gamma)u_\epsilon = 0 - \gamma(\Omega_1, \Gamma)u_\epsilon.$$

Setting $w = u - u_\epsilon|_\Omega$, we have

$$\|\gamma(\Omega, \Gamma)w\|_{\frac{1}{2},\Gamma} = \|\gamma(\Omega_1, \Gamma)u_\epsilon\|_{\frac{1}{2},\Gamma} \leq C \|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon.$$

We define an operator $A : H^1(\Omega) \rightarrow H^{-1}(\Omega)$ by $\langle Au, v \rangle = (\nabla u, \nabla v)_\Omega$, $\forall v \in H_0^1(\Omega)$. We observe that for any $v \in H_0^1(\Omega)$,

$$\begin{aligned} \langle Aw, v \rangle &= (\nabla u, \nabla v)_\Omega - (\nabla u_\epsilon, \nabla v)_\Omega = (\nabla u, \nabla v)_\Omega - (\nabla u_\epsilon, \nabla \tilde{v})_\Omega \\ &= (f, v)_\Omega - (\tilde{f}, \tilde{v})_D + \frac{1}{\epsilon} (\nabla u_\epsilon, \nabla \tilde{v})_{\Omega_1} \\ &= (f, v)_\Omega - (f, v)_\Omega + \frac{1}{\epsilon} (\nabla u_\epsilon, 0)_{\Omega_1} = 0, \end{aligned}$$

where \tilde{v} means the zero extension of v into D . This implies $Aw = 0$ in $H^{-1}(\Omega)$.

Since $w \rightarrow \{Aw, \gamma(\Omega, \Gamma)w\}$ is an isomorphic map of $H^1(\Omega) \rightarrow H^{-1}(\Omega) \times H^{\frac{1}{2}}(\Gamma)$, we obtain that

$$\|u - u_\epsilon\|_{1,\Omega} = \|w\|_{1,\Omega} \leq C(\|Aw\|_{H^{-1}(\Omega)} + \|\gamma(\Omega, \Gamma)w\|_{\frac{1}{2},\Gamma}) \leq C(0 + C\epsilon) \leq C\epsilon,$$

which completes the proof. \square

Remark 2. The conclusion of Theorem 2.1 remains valid even for $f \in H^{-1}(\Omega)$.

Remark 3. The saddle-point method in [10] requires a symmetric variational form, and the operator method in [15] is not so easy to deal with the operator of time-derivative when one considers parabolic problems. However, our method of analysis is applicable to parabolic problems; this is a recent achievement, and will be presented in our future work.

2.2. Neumann boundary value problem. The original problem (Q) with homogeneous Neumann boundary condition reads as:

$$(2.3) \quad \begin{cases} \text{Find } u \in H^1(\Omega) \text{ such that} \\ (u, v)_{1, \Omega} = (f, v)_{\Omega}, \quad \forall v \in H^1(\Omega), \end{cases}$$

where $(u, v)_{1, \Omega} := (\nabla u, \nabla v)_{\Omega} + (u, v)_{\Omega}$.

The H^1 -penalty problem (Q_{ϵ}) reads as:

$$(2.4) \quad \begin{cases} \text{Find } u_{\epsilon} \in H_0^1(D) \text{ such that} \\ (u_{\epsilon}, v)_{1, \Omega} + \epsilon(\nabla u_{\epsilon}, \nabla v)_{\Omega_1} = (\tilde{f}, v)_D, \quad \forall v \in H_0^1(D). \end{cases}$$

Theorem 2.4. *There exist unique solutions u and u_{ϵ} for (2.3) and (2.4), respectively, and we have the following estimates:*

$$(2.5) \quad \|u_{\epsilon} - u\|_{1, \Omega} \leq C\epsilon,$$

$$(2.6) \quad \|u_{\epsilon}\|_{1, \Omega_1} \leq C.$$

Proof. Firstly, substituting $v = u_{\epsilon}$ into (2.4), we obtain

$$\|u_{\epsilon}\|_{1, \Omega}^2 + \epsilon\|\nabla u_{\epsilon}\|_{\Omega_1}^2 = (f, u_{\epsilon})_{\Omega}.$$

This gives

$$\|u_{\epsilon}\|_{1, \Omega} \leq C\|f\|_{\Omega}.$$

Since $v \in H_0^1(D)$ implies $v|_{\Omega} \in H^1(\Omega)$, subtracting (2.3) from (2.4), we have

$$(2.7) \quad (u_{\epsilon} - u, v)_{1, \Omega} + \epsilon(\nabla u_{\epsilon}, \nabla v)_{\Omega_1} = 0, \quad \forall v \in H_0^1(D).$$

Setting $v = u_{\epsilon} - E_{1,0}(\Omega)u_{\epsilon}$ in (2.7), and noticing that $v|_{\Omega} = 0$, we have

$$(u_{\epsilon} - u, 0)_{1, \Omega} + \epsilon(\nabla u_{\epsilon}, \nabla(u_{\epsilon} - E_{1,0}(\Omega)u_{\epsilon})).$$

And we see that

$$\begin{aligned} \|\nabla u_{\epsilon}\|_{\Omega_1}^2 &= (\nabla u_{\epsilon}, \nabla E_{1,0}(\Omega)u_{\epsilon})_{\Omega_1} \\ &\leq \|\nabla u_{\epsilon}\|_{\Omega_1} \|\nabla E_{1,0}(\Omega)u_{\epsilon}\|_{\Omega_1} \leq \|\nabla u_{\epsilon}\|_{\Omega_1} \|E_{1,0}(\Omega)u_{\epsilon}\|_{1, D} \\ &\leq C\|\nabla u_{\epsilon}\|_{\Omega_1} \|u_{\epsilon}\|_{1, \Omega} \leq C\|\nabla u_{\epsilon}\|_{\Omega_1} \|f\|_{\Omega}. \end{aligned}$$

Together with *Friedrichs'* inequality, we have

$$\|u_{\epsilon}\|_{1, \Omega_1} \leq C\|\nabla u_{\epsilon}\|_{\Omega_1} \leq C\|f\|_{\Omega}.$$

Thus, we proved (2.6).

In order to derive (2.5), we substitute $v = E_{1,0}(\Omega)(u_{\epsilon}|_{\Omega} - u) \in H_0^1(D)$ in (2.7). Then, together with (2.6), we have

$$\|u_{\epsilon} - u\|_{1, \Omega}^2 = -\epsilon(\nabla u_{\epsilon}, \nabla v_1)_{\Omega_1} \leq C\epsilon\|v_1\|_{1, \Omega_1} \leq C_1\epsilon\|u_{\epsilon}|_{\Omega} - u\|_{1, \Omega},$$

which implies (2.5). We complete the proof. \square

Remark 4. Recall Remark 3 again. It should be noticed that our analysis method is much simpler than that in [10, 15].

2.3. Mixed boundary value problem. Let $\Gamma = \Gamma_1 \cup \Gamma_2$, $\Gamma_1 \cap \Gamma_2 = \emptyset$. The domain $D \setminus \overline{\Omega}$ is assumed to be split into two parts Ω_1 and Ω_2 , which respectively share the boundaries Γ_1 and Γ_2 with Ω . In addition both share non-empty measure boundary with D (see Figure 3). Set

$$V = \{u \in H^1(\Omega) \mid \gamma(\Omega, \Gamma_1)u = 0\}.$$

The original problem (Q) with homogeneous mixed boundary is stated as:

$$(2.8) \quad \begin{cases} \text{Find } u \in V \text{ such that} \\ (\nabla u, \nabla v)_\Omega = (f, v)_\Omega, \quad \forall v \in V. \end{cases}$$

The H^1 -penalty problem (Q_ϵ) reads as:

$$(2.9) \quad \begin{cases} \text{Find } u_\epsilon \in H_0^1(D) \text{ such that} \\ (\nabla u_\epsilon, \nabla v)_\Omega + \frac{1}{\epsilon} (\nabla u_\epsilon, \nabla v)_{\Omega_1} \\ \quad + \epsilon (\nabla u_\epsilon, \nabla v)_{\Omega_2} = (\tilde{f}, v), \quad \forall v \in H_0^1(D). \end{cases}$$

Theorem 2.5. *There exist unique solutions u and u_ϵ for (2.8) and (2.9), respectively, and we have the following estimates:*

$$(2.10) \quad \|u_\epsilon - u\|_{1,\Omega} \leq C\epsilon,$$

$$(2.11) \quad \|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon,$$

$$(2.12) \quad \|u_\epsilon\|_{1,\Omega_2} \leq C.$$

Proof. The following results have already been achieved (Theorem I-8 in [9]):

$$(2.13) \quad u_\epsilon|_\Omega \rightarrow u \text{ in } H^1(\Omega);$$

$$(2.14) \quad \frac{u_\epsilon|_{\Omega_1}}{\sqrt{\epsilon}} \rightarrow 0 \text{ in } H^1(\Omega_1);$$

$$(2.15) \quad \sqrt{\epsilon}u_\epsilon|_{\Omega_2} \rightarrow 0 \text{ in } H^1(\Omega_2),$$

as $\epsilon \rightarrow 0$. Noting $\Omega^* = \Omega \cup \Omega_1 \cup \Gamma_1$, we set $v = u_\epsilon - E_{1,0}(\Omega^*)u_\epsilon \in H_0^1(D)$. From (2.13) and (2.14), we have

$$(2.16) \quad \|E_{1,0}(\Omega^*)u_\epsilon\|_{1,D} \leq C\|u_\epsilon\|_{1,\Omega^*} \leq C.$$

We find that $v|_{\Omega^*} = 0$ and substitute this v into (2.9) to obtain

$$\epsilon(\nabla u_\epsilon, \nabla v)_{\Omega_2} = 0.$$

This gives

$$|u_\epsilon|_{1,\Omega_2}^2 = (\nabla u_\epsilon, \nabla E_{1,0}(\Omega^*)u_\epsilon)_{\Omega_2} \leq |u_\epsilon|_{1,\Omega} \|E_{1,0}(\Omega^*)u_\epsilon\|_{1,D}.$$

Combining with $\|u_\epsilon\|_{1,\Omega_2} \leq C|u_\epsilon|_{1,\Omega_2}$ and (2.16), it shows (2.12).

Substituting $v = E_{1,0}(\Omega_1)u_\epsilon$ into (2.9), we have

$$(\nabla u_\epsilon, \nabla E_{1,0}(\Omega_1)u_\epsilon)_\Omega + \frac{1}{\epsilon} \|\nabla u_\epsilon\|_{\Omega_1}^2 + \epsilon (\nabla u_\epsilon, \nabla E_{1,0}(\Omega_1)u_\epsilon)_{\Omega_2} = (\tilde{f}, E_{1,0}(\Omega_1)u_\epsilon),$$

from which we obtain that

$$\begin{aligned} \|u_\epsilon\|_{1,\Omega_1}^2 &\leq C|u_\epsilon|_{1,\Omega_1}^2 \\ &= C \left(\epsilon (\nabla u_\epsilon, \nabla E_{1,0}(\Omega_1)u_\epsilon)_\Omega + \epsilon^2 (\nabla u_\epsilon, \nabla E_{1,0}(\Omega_1)u_\epsilon)_{\Omega_2} + \epsilon (\tilde{f}, E_{1,0}(\Omega_1)u_\epsilon) \right) \\ &\leq C\epsilon (\|f\|_\Omega + |u_\epsilon|_{1,\Omega} + \epsilon |u_\epsilon|_{1,\Omega_2}) \|E_{1,0}(\Omega_1)u_\epsilon\|_{1,D} \leq C\epsilon \|u_\epsilon\|_{1,\Omega_1}, \end{aligned}$$

which implies (2.11).

Setting $w = u_\epsilon|_\Omega - u$, we have $\gamma(\Omega, \Gamma_1)w = \gamma(\Omega, \Gamma_1)u_\epsilon - 0 = \gamma(\Omega_1, \Gamma_1)u_\epsilon$, which shows

$$\|\gamma(\Omega, \Gamma_1)w\|_{\frac{1}{2},\Gamma_1} \leq C\|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon.$$

Applying Lemma 2.3, there exists $w_1 = \gamma(\Omega, \Gamma_1)^{-1}\gamma(\Omega, \Gamma_1)w \in H^1(\Omega)$, satisfying

$$\|w_1\|_{1,\Omega} \leq C\|\gamma(\Omega, \Gamma_1)w\|_{\frac{1}{2},\Gamma_1} \leq C\epsilon.$$

Next, we define a continuous operator $\bar{A} : V \rightarrow V'$ (V' is the dual space of V) by $\langle \bar{A}\phi, v \rangle = (\nabla\phi, \nabla v)_\Omega$, $\phi \in V$, $\forall v \in V$. Applying *Friedrichs'* inequality, we have

$$(\nabla v, \nabla v)_\Omega = |v|_{1,\Omega}^2 \geq C\|v\|_{1,\Omega}^2 = \|v\|_{V'}^2.$$

Hence, the bilinear form $a(\phi, v)$ on $V \times V$ defined by $a(\phi, v) = \langle \bar{A}\phi, v \rangle$ is V -elliptic. It follows from the Lax-Milgram theorem that

$$\|w - w_1\|_V \leq C\|\bar{A}(w - w_1)\|_{V'} \leq C(\|\bar{A}w\|_{V'} + \|\bar{A}w_1\|_{V'}).$$

As $\|u_\epsilon - u\|_{1,\Omega} = \|w\|_V \leq \|w - w_1\|_V + \|w_1\|_V \leq \|w - w_1\|_V + C\epsilon$, to prove (2.10) we need to show that $\|w - w_1\|_V \leq C\epsilon$, where we already know $\|\bar{A}w_1\|_{V'} \leq C\|w_1\|_{1,\Omega} \leq C\epsilon$ by the continuity of \bar{A} . To estimate $\|\bar{A}w\|_{V'}$, we observe that $\forall v \in V$, $\bar{v} = E_{1,0}(\Omega)v \in H_0^1(D)$ with $E_{1,0}(\Omega)v|_{\Omega_1} = 0$,

$$\begin{aligned} \langle \bar{A}w, v \rangle &= (\nabla u_\epsilon, \nabla v)_\Omega - (\nabla u, \nabla v)_\Omega = (\nabla u_\epsilon, \nabla \bar{v})_\Omega - (\nabla u, \nabla v)_\Omega \\ &= (\tilde{f}, \bar{v})_D - \frac{1}{\epsilon} (\nabla u_\epsilon, \nabla \bar{v})_{\Omega_1} - \epsilon (\nabla u_\epsilon, \nabla \bar{v})_{\Omega_2} - (f, v)_\Omega \\ &= (f, v)_\Omega - \frac{1}{\epsilon} (\nabla u_\epsilon, 0)_{\Omega_1} - \epsilon (\nabla u_\epsilon, \nabla \bar{v})_{\Omega_2} - (f, v)_\Omega \\ &\leq \epsilon |u_\epsilon|_{1,\Omega_2} |\bar{v}|_{1,\Omega_2} \\ &\leq C\epsilon \|v\|_{1,\Omega}, \end{aligned}$$

that is $\|\bar{A}w\|_{V'} \leq C\epsilon$. Thus, we have $\|w - w_1\|_V \leq C(\|\bar{A}w\|_{V'} + \|\bar{A}w_1\|_{V'}) \leq C\epsilon$, and the proof is completed. \square

Remark 5. Basically, the proof for the mixed boundary case is a combination of those for the Dirichlet and Neumann boundary cases.

Remark 6. All of the above in this section, which involve homogeneous boundary conditions are also suitable for the non-homogeneous boundary value problems.

3. THE REGULARITY OF THE SOLUTIONS OF H^1 -PENALTY PROBLEMS

This section is devoted to the regularity theorems for (Q_ϵ) with homogeneous Dirichlet, Neumann, and mixed boundary conditions respectively. As it is shown in [9], these (Q_ϵ) are equal to certain interface elliptic problems, denoted as (P_ϵ) . There are some regularity theorems for the interface elliptic problems in the literature(see [1, 12] for example), however, these are not specific to our problems. Our particular objective is to deduce explicit dependence of various norms of u_ϵ on the penalty parameter ϵ . We show some estimates which are only suitable for our problems, as well as some higher-order regularity which will be used in the study of the H^1 -penalty parabolic problem.

3.1. Dirichlet boundary value problem. As a first step, let us assume D is sufficiently smooth. Then, we have the following theorem.

Theorem 3.1. *Let $u_\epsilon \in H_0^1(D)$ be the solution of the H^1 -penalty problem (1.2) for $f \in L^2(\Omega)$. Then, we have*

$$u_\epsilon|_\Omega \in H^2(\Omega), \quad u_\epsilon|_{\Omega_1} \in H^2(\Omega_1),$$

$$(3.1) \quad \|u_\epsilon\|_{2,\Omega} \leq C\|f\|_{0,\Omega},$$

$$(3.2) \quad \|u_\epsilon\|_{2,\Omega_1} \leq C\epsilon\|f\|_{0,\Omega}.$$

First, we recall the following basic regularity result:

Lemma 3.2. *Let $\omega \subset \mathbb{R}^2$ be a bounded domain. Assume that the boundary $\partial\omega$ is divided into two disjoint smooth components $\partial\omega_1$ and $\partial\omega_2$; $\partial\omega = \partial\omega_1 \cup \partial\omega_2$. Let $v \in H^1(\omega)$ be the unique weak solution of*

$$\Delta v = f \text{ in } \omega, \quad v = g_1 \text{ on } \partial\omega_1, \quad \frac{\partial v}{\partial n} = g_2 \text{ on } \partial\omega_2,$$

where $f \in L^2(\omega)$, $g_1 \in H^{\frac{1}{2}}(\partial\omega_1)$ and $g_2 \in L^2(\partial\omega_2)$.

Then, if $g_1 \in H^{\frac{3}{2}}(\partial\omega_1)$ and $g_2 \in H^{\frac{1}{2}}(\partial\omega_2)$, we have $v \in H^2(\omega)$ and

$$\|v\|_{2,\omega} \leq C(\|f\|_\omega + \|g_1\|_{H^{\frac{3}{2}}(\partial\omega_1)} + \|g_2\|_{H^{\frac{1}{2}}(\partial\omega_2)}).$$

Before the proof, we see that, by applying Green's formula, (1.2) is equivalent to (P_ϵ) , which reads as:

$$(3.3) \quad \begin{cases} \text{Find } u_\epsilon \in H_0^1(D) \text{ such that} \\ -\Delta u_\epsilon = f \text{ in } \Omega, \\ \Delta u_\epsilon = 0 \text{ in } \Omega_1, \\ \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega} = \frac{1}{\epsilon} \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega_1}, \end{cases}$$

where, for example,

$$\left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega} = (\gamma(\Omega, \Gamma) \nabla u_\epsilon) \cdot n.$$

From (P_ϵ) , we see that, if we have $u_\epsilon|_\Omega \in H^2(\Omega)$ with $\|u_\epsilon\|_{2, \Omega} \leq C$, then we have

$$\left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega_1} = \epsilon \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega} \in H^{\frac{1}{2}}(\Gamma).$$

Moreover, according to the trace theorem, we have

$$\left\| \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\frac{1}{2}, \Omega_1} \right\| \leq C \|u_\epsilon\|_{2, \Omega} \leq C.$$

Then, applying Lemma 3.2, we obtain

$$\begin{aligned} \|u_\epsilon\|_{2, \Omega_1} &\leq C \left(\|0\|_\Omega + \|0\|_{H^{\frac{3}{2}}(\partial D)} + \left\| \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega_1} \right\|_{\frac{1}{2}, \Gamma} \right) \\ &= C \left\| \epsilon \left. \frac{\partial u_\epsilon}{\partial n} \right|_{\Gamma, \Omega} \right\|_{\frac{1}{2}, \Gamma} \leq C\epsilon. \end{aligned}$$

Now, we can state the following proof.

Proof of Theorem 3.1. From the discussion above, we only need to show that $u_\epsilon|_\Omega \in H^2(\Omega)$ and $\|u_\epsilon\|_{2, \Omega} \leq C$. This is a well-known result; however, we want to present a brief proof here, because we will show that, by a slight change of this process, we can obtain a higher-order regularity, with smoother assumptions on f .

There exist $\{U_j\}_{j=1}^N$, $U_j \in \mathbb{R}^2$, $\Phi_j \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$, with $\Psi_j = \Phi_j^{-1}$, $j = 1, 2, \dots, N$.

$$\bar{\Omega} \subset \cup_{j=1}^N \Phi_j(U_j) \subset D. \quad U_{j_0} := \Psi_j(\Phi_j(U_j) \cap \Omega) = \mathbb{R}_+^2 \cap U_j,$$

$$U_{j_1} := \Psi_j(\Phi_j(U_j) \cap \Omega_1) = \mathbb{R}_-^2 \cap U_j, \quad j = 1, 2, \dots, N.$$

And also, there exists $\theta_j \in C_0^\infty(\bar{\Omega})$ with $\text{supp } \theta_j \subset \Phi_j(U_j)$, $j = 1, 2, \dots, N$, with

$$\sum_{j=1}^N \theta_j = 1, \text{ on } \bar{\Omega}.$$

Hence, $(\theta_j u_\epsilon) \circ \Phi_j \in H_0^1(U_j)$, $j = 1, 2, \dots, N$. Now we write $U, U_1, U_0, \Phi, \Psi, \theta$ instead of $U_j, U_{j_1}, U_{j_0}, \Phi_j, \Psi, \theta_j$. Setting $u_2 := (\theta_j u_\epsilon) \circ \Phi_j$, we investigate u_2 in the H^2 space for two cases.

(1) The case $U_1 = \emptyset$. We find $u_1 := \theta u_\epsilon$ satisfies

$$(\nabla u_1, \nabla v)_\Omega = (\theta f - \nabla u_\epsilon \nabla \theta - \nabla(u_\epsilon \nabla \theta), v), \quad \forall v \in H_0^1(\Omega).$$

Since $(\theta f - \nabla u_\epsilon \nabla \theta - \nabla(u_\epsilon \nabla \theta))|_\Omega \in L^2(\Omega)$, obviously, we have

$$u_1|_\Omega \in H^2(\Omega), \quad \|u_1\|_2 \leq C.$$

So, $u_2 \in H^2(U)$ and $\|u_2\|_{2,U} \leq C$.

(2) The case $U_0 \neq \emptyset$, and $U_1 \neq \emptyset$. Writing $D_i = \frac{\partial}{\partial x_i}$, ($i = 1, 2$), we have $u_2 \in H_0^1(U)$ satisfies

$$\begin{aligned} & \sum_{i,j=1}^2 \int_{U_0} a_{ij} D_i u_2 D_j v dx + \frac{1}{\epsilon} \sum_{i,j=1}^2 \int_{U_1} a_{ij} D_i u_2 D_j v dx \\ & - \int_{U_0} (u_\epsilon \nabla \theta) \circ \Phi \nabla v \nabla \Psi |D\Phi| dx - \frac{1}{\epsilon} \int_{U_1} (u_\epsilon \nabla \theta) \circ \Phi \nabla v \nabla \Psi |D\Phi| dx \\ & = (f_1 \circ \Phi |D\Phi|, v), \quad \forall v \in H_0^1(U), \end{aligned}$$

where

$$a_{ij} = \left(\sum_{k=1}^2 D_k \psi_i D_k \psi_j \right) \circ \Phi |D\Phi|, \quad i, j = 1, 2, \quad \Psi = (\psi_1, \psi_2).$$

Let \tilde{u}_2 be the zero extension of u_2 onto \mathbb{R}^2 . Substituting in

$$v = \frac{\tau_h - 1}{h} \frac{\tau_{-h} - 1}{h} \tilde{u}_2$$

into the above equation, where τ_h is the translation operator with $\tau_h \phi(x) = \phi(x_1 + h, x_2)$, $\phi(x) \in L^2(\mathbb{R}^2)$. Using some lemmas of $\frac{\tau_h - 1}{h}$ in Chapter 2 of [3], we can obtain

$$\begin{aligned} & \sum_{i=1}^2 \left\| D_i \left(\frac{\tau_h - 1}{h} \tilde{u}_2 \right) \right\|_{U_0}^2 + \frac{1}{\epsilon} \sum_{i=1}^2 \left\| D_i \left(\frac{\tau_h - 1}{h} \tilde{u}_2 \right) \right\|_{U_1}^2 \\ & \leq C \sum_{i=1}^2 \left(\left\| D_i \left(\frac{\tau_h - 1}{h} \tilde{u}_2 \right) \right\|_{U_0} + \left\| D_i \left(\frac{\tau_h - 1}{h} \tilde{u}_2 \right) \right\|_{U_1} \right) \end{aligned}$$

Thus, we get

$$\sum_{i=1}^2 \left\| D_i \left(\frac{\tau_h - 1}{h} \tilde{u}_2 \right) \right\|_{U_0} \leq C.$$

For $i = 1, 2$, we consider any sequence $h_j \rightarrow 0$, as $j \rightarrow \infty$. We see that $D_i \frac{\tau_{h_j} - 1}{h_j} \tilde{u}_2$ converges weakly to some function $\phi \in L^2(\mathbb{R}^2)$, and also

$$D_i \frac{\tau_{h_j} - 1}{h_j} \tilde{u}_2 \rightarrow D_i D_1 \tilde{u}_2, \text{ as } j \rightarrow \infty,$$

in the sense of distribution. Consequently, $D_i D_1 \tilde{u}_2 = \phi \in L^2(\mathbb{R}_+^2)$. This shows that all second derivatives, except $D_2^2 u_2$, are in $L^2(\mathbb{R}^2)$. We find that the equation of u_2 is also equivalent to

$$\begin{aligned} & (\nabla u_1, \nabla v)_\Omega + \frac{1}{\epsilon} (\nabla u_\epsilon, \nabla v)_{\Omega_1} \\ &= (\theta \tilde{f}, v) - (\nabla u_\epsilon, v \nabla \theta)_\Omega - \frac{1}{\epsilon} (\nabla u_\epsilon, v \nabla \theta)_{\Omega_1} - (\nabla(u_\epsilon \nabla \theta))_\Omega - \frac{1}{\epsilon} (\nabla(u_\epsilon \nabla \theta))_{\Omega_1} \\ &+ \int_\Gamma (u_\epsilon \nabla \theta v)|_{\Gamma, \Omega} \cdot \mathbf{n} d\sigma - \frac{1}{\epsilon} \int_\Gamma (u_\epsilon \nabla \theta v)|_{\Gamma, \Omega_1} \cdot \mathbf{n} d\sigma \\ &:= (f_3, v)_D + (g, v)_\Gamma \quad \forall v \in H_0^1(D). \end{aligned}$$

As $\|f_3\|_{0,D} \leq C$ and $\|g\|_{\frac{1}{2},\Gamma} \leq C$ (since $\|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon$), there exists $w \in H^2(\Omega)$ satisfying that $\frac{\partial w}{\partial n} = g$. Thus, $(u_1 - w) \circ \Phi = u_2 - w \circ \Phi$, and from the equation of u_1 above, w satisfies

$$\sum_{i,j=1}^2 D_j (a_{ij} (D_i (u_2 - w \circ \Phi))) = (f_3 + \Delta w) \circ \Phi \text{ in } U_0,$$

which comes from Green's formula.

Thus we have $D_2^2 u_2 \in L^2(U_0)$, so that $\|u_2\|_{2,U_0} \leq C$. And because for every $j = 1, 2, \dots, N$, $\|u_2\|_{2,U_{j_0}} \leq C$. We have $\|u_\epsilon\|_{2,\Omega} \leq C$. \square

Remark 7. With the assumption that $f \in H^1(\Omega)$. In the above proof, we find that by taking $v = \frac{\tau_h - 1}{h} \frac{\tau_h - 1}{h} \frac{\tau_{-h} - 1}{h} \tilde{u}_2$ or $\frac{\tau_h - 1}{h} \frac{\tau_{-h} - 1}{h} \frac{\tau_{-h} - 1}{h} \tilde{u}_2$ on case(2) instead of $\frac{\tau_h - 1}{h} \frac{\tau_{-h} - 1}{h} \tilde{u}_2$, we can obtain $D_1^3 u_2, D_1^2 D_2 u_2, D_2^2 D_1 u_2 \in L^2(U_0)$. Noticing that in this time $f_3 \circ \Phi \in H^1(U_0)$, $g \in H^{\frac{3}{2}}(\Gamma)$, we have $\|u\|_{3,\Omega} \leq C$. And applying the (P_ϵ) , we have $\|u\|_{3,\Omega_1} \leq C\epsilon$. Hence, we can obtain higher-order regularity of (Q_ϵ) .

By an analogue of the proof of Theorem 3.1 and the Remark 7, we have

Theorem 3.3. *Under the assumption that $f \in H^k(\Omega)$, for all non-negative integers k , we have*

$$u_\epsilon|_\Omega \in H^{k+2}(\Omega), \quad u_\epsilon|_{\Omega_1} \in H^{k+2}(\Omega_1).$$

$$(3.4) \quad \|u_\epsilon|_\Omega\|_{k+2,\Omega} \leq C\|f\|_{k,\Omega},$$

$$(3.5) \quad \|u_\epsilon|_{\Omega_1}\|_{k+2,\Omega_1} \leq C\epsilon\|f\|_{k,\Omega}.$$

Remark 8. In the above two theorems, we both assume that D is sufficiently smooth; however, in our case, D is a rectangle (a convex polygon). From the discussion in [3, 4] on elliptic problems in non-smooth domains, we can keep Theorem 3.1 remains true for any convex polygon D .

3.2. Neumann boundary value problem. As a first step, let us assume D is sufficiently smooth. Then, we have the following theorem.

Theorem 3.4. *(Q) is the original problem with homogeneous Neumann boundary. $f \in L^2(\Omega)$, then the corresponding H^1 -penalty problem (Q_ϵ) has a unique solution $u_\epsilon \in H_0^1(D)$. Moreover,*

$$u_\epsilon|_\Omega \in H^2(\Omega), \quad u_\epsilon|_{\Omega_1} \in H^2(\Omega_1),$$

$$(3.6) \quad \|u_\epsilon|_\Omega\|_{2,\Omega} \leq C\|f\|_{0,\Omega},$$

$$(3.7) \quad \|u_\epsilon|_{\Omega_1}\|_{2,\Omega_1} \leq C\|f\|_{0,\Omega}.$$

Before the proof, we show that, by applying the Green's formula, (Q_ϵ) is equivalent to (P_ϵ), which reads as:

$$(3.8) \quad \begin{cases} \text{Find } u_\epsilon \in H_0^1(D) \text{ such that} \\ -\Delta u_\epsilon + u_\epsilon = f \text{ in } \Omega, \\ \Delta u_\epsilon = 0 \text{ in } \Omega_1, \\ \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma,\Omega} = \epsilon \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma,\Omega_1}. \end{cases}$$

From (P_ϵ) we know that, if we have $u_\epsilon|_\Omega \in H^2(\Omega)$ with $\|u_\epsilon\|_{2,\Omega} \leq C$, then we have

$$u_\epsilon|_{\Gamma,\Omega_1} = u_\epsilon|_{\Gamma,\Omega} \in H^{\frac{3}{2}}(\Omega).$$

Since the right-hand-side function is 0, and with the homogeneous Dirichlet boundary of D , it concludes that $u_\epsilon|_{\Omega_1} \in H^2(\Omega_1)$ and

$$\|u_\epsilon\|_{2,\Omega_1} \leq C\|u_\epsilon\|_{\frac{3}{2},\Gamma} \leq C.$$

This means we have left to prove only $u_\epsilon|_\Omega \in H^2(\Omega)$.

Proof. The process of the proof is very similar to that of Theorem 3.1. In fact, we only need to replace the $\frac{1}{\epsilon}$ in the proof of Theorem 3.1 by ϵ . \square

Remark 9. The same comments as those in the Dirichlet case apply here, specifically, we can obtain higher-order regularity for $f \in H^k(\Omega)$, and, if D is a convex polygon, Theorem 3.4 remains true.

3.3. Mixed boundary value problem. The (Q_ϵ) for original problem with homogeneous mixed boundary is equivalent to the problem (P_ϵ) :

$$(3.9) \quad \left\{ \begin{array}{l} \text{Find } u_\epsilon \in H_0^1(D) \text{ such that} \\ -\Delta u_\epsilon = f \text{ in } \Omega, \\ \Delta u_\epsilon = 0 \text{ in } \Omega_1, \\ \Delta u_\epsilon = 0 \text{ in } \Omega_2, \\ \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_1, \Omega} = \frac{1}{\epsilon} \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_1, \Omega_1}, \\ \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_2, \Omega} = \epsilon \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_2, \Omega_2}, \\ \frac{1}{\epsilon} \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_3, \Omega_1} = \epsilon \frac{\partial u_\epsilon}{\partial n} \Big|_{\Gamma_3, \Omega_2}, \end{array} \right.$$

where Γ_3 is the common boundary of Ω_1 and Ω_2 . By an analogue of the previous proof, we can obtain that $u_\epsilon|_\Omega \in H^2(\Omega)$.

Since the domain Ω_1 and Ω_2 have corners at the intersection points of their boundaries(see Fig. 3), $u_\epsilon|_{\Omega_1}$ and $u_\epsilon|_{\Omega_2}$ would not be in H^2 space but $H^{1+\alpha}$, $\alpha \in (0, 1)$ (see [12, 4]).

4. FINITE ELEMENT APPROXIMATION AND DISCRETE PROBLEMS

Recall that the Cartesian mesh is introduced to the rectangular domain D to get a uniform triangulation \mathcal{T}_h , and h is the maximum diameter of the triangles of \mathcal{T}_h . Each $K \in \mathcal{T}_h$ is assumed to be a closed set. $V_h(D) \subset H_0^1(D)$ is the subspace of all piecewise linear continuous functions subordinate to \mathcal{T}_h .

4.1. Dirichlet boundary value problem. We consider the discrete problem (1.3).

Lemma 4.1. *There exists a unique solution $u_{\epsilon, h} \in V_h(D)$ for (1.3). u_ϵ is the solution of (1.2), and we have*

$$(4.1) \quad \begin{aligned} & \|u_\epsilon - u_{\epsilon, h}\|_{1, \Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - u_{\epsilon, h}\|_{1, \Omega_1} \\ & \leq C \inf_{v_h \in V_h(D)} \left(\|u_\epsilon - v_h\|_{1, \Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - v_h\|_{1, \Omega_1} \right). \end{aligned}$$

Proof. Subtracting (1.2) from (1.3), we have

$$(\nabla(u_\epsilon - u_{\epsilon, h}), \nabla v_h)_\Omega + \frac{1}{\epsilon} (\nabla(u_\epsilon - u_{\epsilon, h}), \nabla v_h)_{\Omega_1} = 0, \quad \forall v_h \in V_h(D).$$

Then we find

$$\begin{aligned}
& |u_\epsilon - u_{\epsilon,h}|_{1,\Omega}^2 + \frac{1}{\epsilon} |u_\epsilon - u_{\epsilon,h}|_{1,\Omega_1}^2 \\
& \leq \inf_{v_h \in V_h(D)} \left\{ (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla(u_\epsilon - v_h))_\Omega + \frac{1}{\epsilon} (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla(u_\epsilon - v_h))_{\Omega_1} \right\} \\
& \leq \inf_{v_h \in V_h(D)} \left\{ |u_\epsilon - u_{\epsilon,h}|_{1,\Omega} |u_\epsilon - v_h|_{1,\Omega} + \frac{1}{\epsilon} |u_\epsilon - u_{\epsilon,h}|_{1,\Omega_1} |u_\epsilon - v_h|_{1,\Omega_1} \right\} \\
& \leq \left\{ \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1} \right\} \times \\
& \quad \inf_{v_h \in V_h(D)} \left\{ \|u_\epsilon - v_h\|_{1,\Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - v_h\|_{1,\Omega_1} \right\}.
\end{aligned}$$

Applying the *Poincaré* inequality to the left-hand-side, we have

$$\begin{aligned}
& |u_\epsilon - u_{\epsilon,h}|_{1,\Omega}^2 + \frac{1}{\epsilon} |u_\epsilon - u_{\epsilon,h}|_{1,\Omega_1}^2 \\
& = |u_\epsilon - u_{\epsilon,h}|_{1,D}^2 + \left(\frac{1}{\epsilon} - 1 \right) |u_\epsilon - u_{\epsilon,h}|_{1,\Omega_1}^2 \\
& \geq C \left(\|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega}^2 + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1}^2 \right)
\end{aligned}$$

Thus, we have proved the result. \square

To estimate

$$\inf_{v_h \in V_h(D)} \left(\|u_\epsilon - v_h\|_{1,\Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - v_h\|_{1,\Omega_1} \right),$$

we need some lemmas, which can be found in [16], and several other similar results in [13, 17]. For a curve γ in $C^2(\mathbb{R}^2)$ and $\delta > 0$, we define a δ -neighborhood $\gamma_\delta = \{x \in \mathbb{R}^2 \mid \text{dist}(x, \gamma) \leq \delta\}$.

Lemma 4.2. *Suppose $\gamma_\delta \subset\subset R$, R is a domain in \mathbb{R}^2 , and $v \in H^1(R)$. Then we have*

$$(4.2) \quad \|v\|_{0,\gamma_\delta} \leq C\sqrt{\delta} \|v\|_{1,R}.$$

If we assume $v \in H^2(R)$, then we have

$$(4.3) \quad \|v\|_{1,\gamma_\delta} \leq C\sqrt{\delta} \|v\|_{2,R}.$$

Lemma 4.3. *Suppose $w \in H^2(D)$, and we define $I_K w$ as the linear interpolation of w on the vertices of a triangle $K \in \mathcal{T}_h$. Then, we have*

$$(4.4) \quad |I_K w|_{1,K} \simeq \sum_{i,j=1, i \neq j}^3 |w(\nu_i) - w(\nu_j)|,$$

where ν_i , $i = 1, 2, 3$, are vertices of K . ($A \simeq B$ means that there exist constants depending on the regularity of the triangulation C_1, C_2 such that $C_2 B \leq A \leq C_1 B$.)

Theorem 4.4. *The following error estimate for (1.2) and (1.3) holds:*

$$(4.5) \quad \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega} + \frac{1}{\sqrt{\epsilon}} \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1} \leq C(\sqrt{h} + \sqrt{\epsilon})(\|u_\epsilon\|_{2,\Omega} + \|u_\epsilon\|_{2,\Omega_1}).$$

Before the proof, we define some notations:

$$\begin{aligned} \Lambda_h &= \text{the set of all vertices of } \mathcal{T}_h, \\ \Lambda(K) &= \{\nu_1^K, \nu_2^K, \nu_3^K\} = \text{the set of all vertices of } K \in \mathcal{T}_h, \\ T_\Omega &= \{K \in \mathcal{T}_h \mid K \subset \Omega, K \cap \Gamma = \emptyset\}, \\ T_{\Omega_1} &= \{K \in \mathcal{T}_h \mid K \subset \Omega_1, K \cap \Gamma = \emptyset\}, \\ T_\Gamma &= \{K \in \mathcal{T}_h \mid K \cap \Gamma \neq \emptyset\}, \\ T_0 &= \{K \in T_\Omega \mid K \cap T_\Gamma \neq \emptyset\}, \\ T_1 &= \{K \in T_{\Omega_1} \mid K \cap T_\Gamma \neq \emptyset\}, \\ \bar{\omega}_0 &= \cup_{K \in T_0} K, \\ \bar{\omega}_\Omega &= \cup_{K \in T_\Omega \setminus T_0} K, \\ \bar{\omega}_\Gamma &= \cup_{K \in T_\Gamma} K. \end{aligned}$$

We may assume that $T_\Omega \setminus T_0 \neq \emptyset$ and $T_{\Omega_1} \setminus T_1 \neq \emptyset$ without loss of generality.

Proof of Theorem 4.4. We define v_h by setting,

$$v_h(\nu) = \begin{cases} u_\epsilon(\nu) & \text{for } \nu \in \Lambda(K), K \in T_\Omega \setminus T_0, \\ 0 & \text{for all others vertices } \nu, \end{cases}$$

and substitute this v_h into the right-hand-side of (4.1). We find that $\|u_\epsilon - v_h\|_{1,\Omega_1} = \|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon$. To estimate $\|u_\epsilon - v_h\|_{1,\Omega}$, we use the scheme proposed in [16] by using Lemma 4.2 and Lemma 4.3. However, there are several differences between our analysis and that of [16], because we apply our regularity theorem presented in the previous section, which simplifies the analysis.

We find that $\|u_\epsilon - v_h\|_{1,\Omega}^2 = \|u_\epsilon - v_h\|_{1,\omega_\Omega}^2 + \|u_\epsilon - v_h\|_{1,\omega_0}^2 + \|u_\epsilon - v_h\|_{1,\omega_\Gamma}^2$.

For the first term $\|u_\epsilon - v_h\|_{1,\omega_\Omega} = \|u_\epsilon - I_K u_\epsilon\|_{1,\omega_\Omega} \leq Ch \|u_\epsilon\|_{2,\Omega}$.

For the third term $\|u_\epsilon - v_h\|_{1,\omega_\Gamma} = \|u_\epsilon - 0\|_{1,T_{\omega_\Gamma}} \leq C\sqrt{h} \|u_\epsilon\|_{2,\Omega}$, following from Lemma 4.2.

Then, for the second term $\|u_\epsilon - v_h\|_{1,\omega_0} \leq \|u_\epsilon\|_{1,\omega_0} + \|v_h\|_{1,\omega_0}$, and we have $\|u_\epsilon\|_{1,\omega_0} \leq C\sqrt{h} \|u_\epsilon\|_{2,\Omega}$, again, following from Lemma 4.2. What remains is to estimate $\|v_h\|_{1,\omega_0}$.

For every $K \in T_0$, we have

$$\begin{aligned} & \|v_h\|_{1,K} \leq \|v_h - I_K u_\epsilon\|_{1,K} + \|I_K \bar{u}_\epsilon\|_{1,K} \\ & \leq |v_h - I_K u_\epsilon|_{1,K} + \|v_h - I_K u_\epsilon\|_{0,K} + \|I_K u_\epsilon\|_{1,K} \\ & \leq |v_h - I_K u_\epsilon|_{1,K} + \|v_h\|_{0,K} + \|I_K u_\epsilon\|_{0,K} + \|I_K u_\epsilon\|_{1,K}. \end{aligned}$$

We want to show that $\|v_h\|_{0,K} \leq C\|I_K u_\epsilon\|_{1,K}$. There are two possibilities: (1) if for all $\nu_i = \nu_i^K \in \Lambda(K)$, $i = 1, 2, 3$, $u_\epsilon(\nu_i)$ have the same sign, then obviously, $|v_h(x)|^2 \leq |I_K u_\epsilon(x)|^2$, ($x \in K$) and $\|v_h\|_{0,K} \leq C\|I_K u_\epsilon\|_{0,K}$; (2) if we consider the case where for ν_i , $i = 1, 2, 3$, $u_i = u_\epsilon(\nu_i)$ do not all have the same signs, then, without loss of generality, we assume that $|u_1| = \|u\|_{\infty,K}$ and $u_1 u_2 \geq 0$ and $u_3 \leq 0$. We have $\nabla(I_K u) \cdot \vec{\nu}_3 \vec{\nu}_1 = u_1 - u_3$ on K . Hence, $|u_1| \leq |u_1 - u_3| \leq |\nabla(I_K u)| \cdot |\vec{\nu}_3 \vec{\nu}_1| \leq |\nabla(I_K u)| h_K$ on K , since u_1 and u_3 have the different signs. Therefore,

$$\begin{aligned} \|v_h\|_{0,K} & \leq |K|^{\frac{1}{2}} \|v_h\|_{\infty,K} \\ & \leq |K|^{\frac{1}{2}} \|u_\epsilon\|_{\infty,K} \\ & \leq |K|^{\frac{1}{2}} |I_K u| h_K = h_K |\nabla(I_K u)|_{0,K}. \end{aligned}$$

Thus, we have $\|v_h\|_{0,K} \leq (1 + h_K)\|I_K u\|_{1,K}$, which gives

$$(4.6) \quad \begin{aligned} \|v_h\|_{1,K} & \leq |v_h - I_K u_\epsilon|_{1,K} + C\|I_K u_\epsilon\|_{1,K} \\ & \leq |v_h - I_K u_\epsilon|_{1,K} + C\|I_K u_\epsilon - u_\epsilon\|_{1,K} + C\|u_\epsilon\|_{1,K} \end{aligned}$$

By the standard interpolation error estimates, we have

$$(4.7) \quad \sum_{K \in T_0} \|I_K u_\epsilon - u_\epsilon\|_{1,K}^2 \leq Ch^2 \|u_\epsilon\|_{2,\Omega}^2.$$

We notice that there exists $C' > 0$ such that $\bar{\omega}_\Gamma \subset \gamma_{C'h}$. By Lemma 4.3, setting $\hat{v}_h = v_h - I_K u_\epsilon$, we obtain

$$(4.8) \quad |v_h - I_K u_\epsilon|_{1,K} \leq C \sum_{\substack{\nu_i, \nu_j \in \Lambda(K) \\ i \neq j}} |\hat{v}_h(\nu_i) - \hat{v}_h(\nu_j)|.$$

Next, we set

$$\Lambda_\Omega(K) = \{\nu \in \Lambda(K) \mid \exists K' \in T_\Omega \setminus T_0, \text{ s.t. } \nu \text{ in } \Lambda(K')\},$$

$$\Lambda_\Gamma(K) = \{\nu \in \Lambda(K) \mid \exists K' \in T_\Gamma, \text{ s.t. } \nu \text{ in } \Lambda(K'')\}.$$

By definition, we see that $\Lambda(K) = \Lambda_\Omega(K) \cup \Lambda_\Gamma(K)$, $\Lambda_\Omega(K) \neq \emptyset$, $\Lambda_\Gamma(K) \neq \emptyset$.

There are two possibilities: for $\{\nu_i = \nu_i^K\}_{i=1}^3 = \Lambda(K)$,

- (i) $\Lambda_\Omega(K) = \{\nu_1, \nu_2\}$, $\Lambda_\Gamma(K) = \{\nu_3\}$;
- (ii) $\Lambda_\Omega(K) = \{\nu_1\}$, $\Lambda_\Gamma(K) = \{\nu_2, \nu_3\}$.

(See Figure 1 and Figure 2.)




FIGURE 1. (i)




FIGURE 2. (ii)

For (i), we have

$$\begin{aligned}\hat{v}_h(\nu_i) &= v_h(\nu_i) - (I_K u_\epsilon)(\nu_i) = 0, \quad i = 1, 2, \\ \hat{v}_h(\nu_3) &= v_h(\nu_3) - (I_K u_\epsilon)(\nu_3) = -u_\epsilon(\nu_3).\end{aligned}$$

Hence, $|\hat{v}_h|_{1,K} \leq C|u_\epsilon(\nu_3)|$. Taking arbitrary point $\nu_\Gamma \in \Gamma \cap K''$, we have $|\hat{v}_h|_{1,K} \leq C(|u_\epsilon(\nu_3) - u_\epsilon(\nu_\Gamma)| + |u_\epsilon(\nu_\Gamma)|)$.

At this stage, we apply the *Sobolev* and *Morrey's* inequalities. Let ω be a Lipschitz domain in \mathbb{R}^2 . They are given as

$$\begin{aligned}\|u\|_{L^\infty(\omega)} &\leq C\|u\|_{H^2(\omega)}, \quad (\forall u \in H^2(\omega)) \\ \|u\|_{W^{1,q}(\omega)} &\leq C\|u\|_{H^2(\omega)}, \quad (1 \leq q < \infty, \forall u \in H^2(\omega)) \\ |u(x) - u(y)| &\leq C\|u\|_{W^{1,q}(\omega)}|x - y|^\alpha, \\ (x, y \in \omega, \quad 2 < q, \quad \alpha &= 1 - \frac{2}{q}, \quad u \in W^{1,q}(\omega))\end{aligned}$$

We choose $q = 4$, and define $K''_\Omega = K'' \cap \Omega$, $K''_{\Omega_1} = K'' \cap \Omega_1$, and we have

$$|u_\epsilon(\nu_3) - u_\epsilon(\nu_\Gamma)| \leq C\|u_\epsilon\|_{W^{1,4}(K''_\Omega)}|\nu_3 - \nu_\Gamma|^{\frac{1}{2}} \leq Ch^{\frac{1}{2}}\|u_\epsilon\|_{2,K''_\Omega},$$

$$|u_\epsilon(\nu_\Gamma)| \leq C\|u_\epsilon\|_{2,K''_{\Omega_1}}.$$

Hence, we obtain $|\hat{v}_h|_{1,K} \leq C\|u_\epsilon\|_{2,K''_{\Omega_1}} + Ch^{\frac{1}{2}}\|u_\epsilon\|_{2,K''_\Omega}$.

Then, applying the same trick to (ii), we can show that

$$\begin{aligned}\hat{v}_h(\nu_1) &= v_h(\nu_1) - (I_K u_\epsilon)(\nu_1) = 0, \\ \hat{v}_h(\nu_i) &= v_h(\nu_i) - (I_K u_\epsilon)(\nu_i) = -u_\epsilon(\nu_i), \quad i = 2, 3, \\ |\hat{v}_h|_{1,K} &\leq C(|u_\epsilon(\nu_2) - u_\epsilon(\nu_3)| + |u_\epsilon(\nu_2)| + |u_\epsilon(\nu_3)|) \\ &\leq C \sum_{i=2,3} |u_\epsilon(\nu_i) - u_\epsilon(\nu_\Gamma)| + C|u_\epsilon(\nu_\Gamma)| \\ &\leq C\|u_\epsilon\|_{2,K''_{\Omega_1}} + Ch^{\frac{1}{2}}\|u_\epsilon\|_{2,K''_{\Omega}}.\end{aligned}$$

Combining (i) and (ii), we have

$$(4.9) \quad \begin{aligned}\sum_{K \in T_0} |\hat{v}_h|_{1,K}^2 &\leq C \sum_{K \in T_0} \left(\|u_\epsilon\|_{2,K''_{\Omega_1}}^2 + Ch\|u_\epsilon\|_{2,K''_{\Omega}}^2 \right) \\ &\leq C\|u_\epsilon\|_{2,\Omega_1}^2 + Ch\|u_\epsilon\|_{2,\Omega}^2.\end{aligned}$$

From (4.6), (4.7), (4.8), (4.9), we can easily derive that

$$\begin{aligned}\|v_h\|_{1,\omega_0}^2 &= \sum_{K \in T_0} \|v_h\|_{1,K}^2 \\ &\leq (\|u_\epsilon\|_{2,\Omega_1}^2 + h\|u_\epsilon\|_{2,\Omega}^2 h^2 \|u_\epsilon\|_{2,\Omega}^2 + h\|u_\epsilon\|_{2,\Omega}^2) \\ &\leq \epsilon^2 \|f\|_{0,\Omega}^2 + C(h^2 + h)\|f\|_{0,\Omega}^2,\end{aligned}$$

where the last inequality is from Theorem 3.1. Hence, we get

$$\|v_h\|_{1,\omega_0} \leq C(\epsilon + h + h^{\frac{1}{2}})\|f\|_{0,\Omega}.$$

Recalling that $\|u_\epsilon - v_h\|_{1,\Omega_1} = \|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon$ and other estimates from the beginning of the proof, we have

$$\|u_\epsilon - v_h\|_{1,\Omega} + \frac{1}{\sqrt{\epsilon}}\|u_\epsilon - v_h\|_{1,\Omega_1} \leq C(\sqrt{\epsilon} + \sqrt{h}).$$

Hence, the theorem follows from Lemma 4.1. \square

Remark 10. Since we have $\|u_\epsilon\|_{2,\Omega_1} \leq C\epsilon$, other choices for v_h than that above can be taken, such as

$$v_h = \begin{cases} u_\epsilon(\nu) & \text{for } \nu \in \Lambda(K), K \in T_\Omega \setminus T_0, \\ \bar{u}_\epsilon(\nu) & \text{for } \nu \in \Lambda(K), K \in T_{\Omega_1} \cup T_\Gamma, \end{cases}$$

where \bar{u}_ϵ is the extension of $u_\epsilon|_{\Omega_1}$ onto D with $\|\bar{u}_\epsilon\|_{2,D} \leq C\|u_\epsilon\|_{2,\Omega_1}$, and the estimate result still holds.

To estimate $\|u_\epsilon - u_{\epsilon,h}\|_{0,D}$, we need the adjoint boundary value problem, which reads as:

$$(4.10) \quad \begin{cases} \text{For any given } f \in L^2(\Omega), \text{ find } u_{\epsilon f} \in H_0^1(D) \text{ such that} \\ (\nabla v, \nabla u_{\epsilon f})_\Omega + \frac{1}{\epsilon}(\nabla v, \nabla u_{\epsilon f})_{\Omega_1} = (\tilde{f}, v), \quad \forall v \in H_0^1(D). \end{cases}$$

(\tilde{f} is the zero extension of f .) We see that there exists a unique solution $u_{\epsilon f} \in H_0^1(D)$, with

$$\|u_{\epsilon f}\|_{2,\Omega} \leq C\|f\|_{0,\Omega}, \quad \|u_{\epsilon f}\|_{2,\Omega_1} \leq C\epsilon\|f\|_{0,\Omega},$$

which follows from Theorem 3.1.

Theorem 4.5. *We have the error estimate in L^2 norm for (1.2) and (1.3),*

$$(4.11) \quad \|u_\epsilon - u_{\epsilon,h}\|_{0,D} \leq C(\epsilon + h + \sqrt{h\epsilon}).$$

Proof. Let $f = (u_\epsilon - u_{\epsilon,h})|_\Omega$ and substitute $v = u_\epsilon - u_{\epsilon,h}$ into the equation of the adjoint problem, we get

$$\begin{aligned} & \|u_\epsilon - u_{\epsilon,h}\|_{0,\Omega}^2 = (f, u_\epsilon - u_{\epsilon,h})_{0,\Omega} \\ & = (\tilde{f}, u_\epsilon - u_{\epsilon,h})_{0,D} \\ & = (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla u_{\epsilon f})_\Omega + \frac{1}{\epsilon} (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla u_{\epsilon f})_{\Omega_1} \\ & \leq (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla(u_{\epsilon f} - v_h))_\Omega + \frac{1}{\epsilon} (\nabla(u_\epsilon - u_{\epsilon,h}), \nabla(u_{\epsilon f} - v_h))_{\Omega_1}, \quad \forall v_h \in V_h(D) \\ & \leq \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega} \|u_{\epsilon f} - v_h\|_{1,\Omega} + \frac{1}{\epsilon} \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1} \|\nabla u_{\epsilon f} - v_h\|_{1,\Omega_1} \\ & \leq C(\sqrt{\epsilon} + \sqrt{h})(\sqrt{\epsilon} + \sqrt{h})\|f\|_{0,\Omega} + C\frac{1}{\epsilon}\sqrt{\epsilon}(\sqrt{\epsilon} + \sqrt{h})\sqrt{\epsilon}(\sqrt{\epsilon} + \sqrt{h})\|f\|_{0,\Omega} \end{aligned}$$

The last inequality follows from Theorem 3.1 and Theorem 4.4. Noticing that $f = (u_\epsilon - u_{\epsilon,h})|_\Omega$, we get

$$\|u_\epsilon - u_{\epsilon,h}\|_{0,\Omega} \leq C(\sqrt{\epsilon} + \sqrt{h})^2$$

With $\|u_\epsilon\|_{1,\Omega_1} \leq C\epsilon$ and $\|u_{\epsilon,h}\|_{1,\Omega_1} \leq C\epsilon$, we have proved the result. \square

4.2. Neumann boundary value problem. The discrete problem $(Q_{\epsilon,h})$ reads as:

$$(4.12) \quad \begin{cases} \text{Find } u_{\epsilon,h} \in V_h(D) \text{ such that} \\ (u_{\epsilon,h}, v_h)_{1,\Omega} + \epsilon(\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega_1} = (\tilde{f}, v_h)_D, \quad \forall v_h \in V_h(D). \end{cases}$$

Lemma 4.6. *There exists a unique solution $u_{\epsilon,h} \in V_h(D)$ for (4.12). u_ϵ is the solution of (2.4), and we have*

$$(4.13) \quad \begin{aligned} & \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega} + \sqrt{\epsilon}\|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1} \\ & \leq C \inf_{v_h \in V_h(D)} (\|u_\epsilon - v_h\|_{1,\Omega} + \sqrt{\epsilon}\|u_\epsilon - v_h\|_{1,\Omega_1}). \end{aligned}$$

Proof. The proof of this lemma is an analogue of that of Lemma 4.1. \square

Then, we have the error estimate theorem:

Theorem 4.7. *We have the error estimate for (2.4) and (4.12)*

$$(4.14) \quad \|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega} + \sqrt{\epsilon}\|u_\epsilon - u_{\epsilon,h}\|_{1,\Omega_1} \leq C(h + \epsilon + \sqrt{h\epsilon})(\|u_\epsilon\|_{2,\Omega} + \|u_\epsilon\|_{2,\Omega_1}).$$

Proof. By taking

$$v_h = \begin{cases} u_\epsilon(\nu) & \text{for } \nu \in \Lambda(K), K \in T_\Omega \cup T_\Gamma, \\ \bar{u}_\epsilon(\nu) & \text{for } \nu \in \Lambda(K), K \in T_{\Omega_1} \setminus T_1, \end{cases}$$

the proof is an analogue of that of Theorem 4.4. \square

With the analogue of the proof of the Dirichlet case, we have the error estimate in L^2 norm for Neumann case.

Theorem 4.8. *For u_ϵ and $u_{\epsilon,h}$ are the solutions of (2.4) and (4.12), respectively, we have*

$$(4.15) \quad \|u_\epsilon - u_{\epsilon,h}\|_{0,D} \leq C(\epsilon + h + \sqrt{h\epsilon})^2.$$

4.3. Mixed boundary value problem. Since the regularity theorem of (Q_ϵ) for mixed boundary case is weak, we will not put a discussion on the error estimates of the discrete problem for this case. Also, we could not find any discussion on this issue in [16, 10] *etc.*

5. AN APPROXIMATION FOR DISCRETE PROBLEMS

In the discrete problem, we find the inner-product $(\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega \text{ or } \Omega_1}$ and $(\tilde{f}, v_h)_D$ (since \tilde{f} is the zero extension of f from Ω onto D) are not applicable to computation, because we assumed that Ω has a curved boundary Γ . The integral of the elements crossing Γ becomes a problem when doing computation. Thus, we need a proper approximation. One way is to replace the integral in the open triangle K , $K \cap \Gamma \neq \emptyset$, of

$$(\nabla u_\epsilon, v_h)_{K \cap \Omega} + \frac{1}{\epsilon} (\nabla u_\epsilon, v_h)_{K \cap \Omega_1}$$

by the integral of

$$(\nabla u_\epsilon, v_h)_{K \cap \hat{\Omega}} + \frac{1}{\epsilon} (\nabla u_\epsilon, v_h)_{K \cap \hat{\Omega}_1}$$

where $\hat{\Omega}$ is a polygon with vertices which are the points of intersection between Γ and the triangles' edges. $\hat{\Omega}$ satisfies $(\Omega \setminus \hat{\Omega}) \cup (\hat{\Omega} \setminus \Omega) \subset \Gamma_\delta$, $\delta = O(h^2)$. $\hat{\Omega}_1 = D \setminus \hat{\Omega}$. The approximation problem of $(Q_{\epsilon,h})$ is denoted as $(\hat{Q}_{\epsilon,h})$.

5.1. Dirichlet boundary value problem. The problem (1.4) is considered. We assume that f_h is some interpolation of \tilde{f} , such that $(f_h, v_h)_D$ is applicable to computation and has $\|f_h - \tilde{f}\|_0 \leq Ch$ holds. For example, suppose $f \in C^1(\Omega)$; then we can choose f_h is the linear interpolation of f on the vertices ν of triangles for every $\nu \in \hat{\Omega}$ and zero on other vertices. Before giving the estimate of $\|\hat{u}_{\epsilon,h} - u_{\epsilon,h}\|_{1,D}$, we quote a lemma from [17]. For any open triangle K , we denote

$$\pi_K = (\Omega_1 \setminus \hat{\Omega}_1) \cap K = (\hat{\Omega} \setminus \Omega) \cap K$$

and

$$\begin{aligned}\hat{\pi}_K &:= (\hat{\Omega}_1 \setminus \Omega_1) \cap K = (\Omega \setminus \hat{\Omega}) \cap K. \\ \pi &:= \cup_K \pi_K. \\ \hat{\pi} &:= \cup_K \hat{\pi}_K.\end{aligned}$$

Lemma 5.1. *As we have $(\Omega \setminus \hat{\Omega}) \cup (\hat{\Omega} \setminus \Omega) \subset \Gamma_\delta$, $\delta = O(h^2)$, the following estimates hold for any $v_h \in V_h$,*

$$(5.1) \quad |v_h|_{1,\pi} \leq h^{\frac{1}{2}} |v_h|_{1,\Omega_1} \quad (\text{or } h^{\frac{1}{2}} |v_h|_{1,\hat{\Omega}}),$$

$$(5.2) \quad |v_h|_{1,\hat{\pi}} \leq h^{\frac{1}{2}} |v_h|_{1,\hat{\Omega}_1} \quad (\text{or } h^{\frac{1}{2}} |v_h|_{1,\Omega}).$$

Then we have the following theorem.

Theorem 5.2. *There exists a unique solution $\hat{u}_{\epsilon,h}$ for (1.4). $u_{\epsilon,h}$ is the solution of (1.3), and we have*

$$(5.3) \quad \|u_{\epsilon,h} - \hat{u}_{\epsilon,h}\|_{1,D} \leq Ch.$$

Proof. Subtracting the equation in (1.4) from that in (1.3), we have

$$\begin{aligned}& (\nabla(u_{\epsilon,h} - \hat{u}_{\epsilon,h}), \nabla v_h)_{\Omega \cap \hat{\Omega}} + \frac{1}{\epsilon} (\nabla(u_{\epsilon,h} - \hat{u}_{\epsilon,h}), \nabla v_h)_{\Omega \cap \hat{\Omega}_1} + (\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega \setminus \hat{\Omega}} \\ & + \frac{1}{\epsilon} (\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega_1 \setminus \hat{\Omega}_1} - (\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega} \setminus \Omega} - \frac{1}{\epsilon} (\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega}_1 \setminus \Omega_1} \\ & = (\tilde{f} - f_h, v_h)_D, \quad \forall v_h \in V_h.\end{aligned}$$

Since $\Omega_1 \setminus \hat{\Omega}_1 = \hat{\Omega} \setminus \Omega$ and $\hat{\Omega}_1 \setminus \Omega_1 = \Omega \setminus \hat{\Omega}$, the above equation can be written as

$$\begin{aligned}& (\nabla(u_{\epsilon,h} - \hat{u}_{\epsilon,h}), \nabla v_h)_{\Omega \cup \hat{\Omega}} + \frac{1}{\epsilon} (\nabla(u_{\epsilon,h} - \hat{u}_{\epsilon,h}), \nabla v_h)_{\Omega \cap \hat{\Omega}_1} \\ & = \left(\frac{1}{\epsilon} - 1\right) (\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega}_1 \setminus \Omega_1} + \left(1 - \frac{1}{\epsilon}\right) (\nabla u_{\epsilon,h}, \nabla v_h)_{\Omega_1 \setminus \hat{\Omega}_1} + (\tilde{f} - f_h, v_h)_D.\end{aligned}$$

We apply Lemma 5.1 to obtain

$$\begin{aligned}|\hat{u}_{\epsilon,h}|_{1,\hat{\Omega}_1 \setminus \Omega_1} &\leq C\sqrt{h} |\hat{u}_{\epsilon,h}|_{1,\hat{\Omega}_1} \leq C\sqrt{h}\epsilon, \\ |v_h|_{1,\hat{\Omega}_1 \setminus \Omega_1} &\leq C\sqrt{h} |v_h|_{1,\hat{\Omega}_1} \quad (\text{or } \sqrt{h} |v_h|_{1,\Omega}), \\ |u_{\epsilon,h}|_{1,\Omega_1 \setminus \hat{\Omega}_1} &\leq C\sqrt{h} |u_{\epsilon,h}|_{1,\Omega_1} \leq C\sqrt{h}\epsilon, \\ |v_h|_{1,\Omega_1 \setminus \hat{\Omega}_1} &\leq C\sqrt{h} |v_h|_{1,\Omega_1} \quad (\text{or } \sqrt{h} |v_h|_{1,\hat{\Omega}}).\end{aligned}$$

Since we have $\|\tilde{f} - f_h\|_{0,D} \leq Ch$, for $v_h = u_{\epsilon,h} - \hat{u}_{\epsilon,h}$, we get

$$\begin{aligned}& |u_{\epsilon,h} - \hat{u}_{\epsilon,h}|_{1,\Omega \cup \hat{\Omega}}^2 + \frac{1}{\epsilon} |u_{\epsilon,h} - \hat{u}_{\epsilon,h}|_{1,\Omega_1 \cap \hat{\Omega}_1}^2 \\ & \leq Ch(|u_{\epsilon,h} - \hat{u}_{\epsilon,h}|_{1,\Omega \cup \hat{\Omega}} + \|u_{\epsilon,h} - \hat{u}_{\epsilon,h}\|_{0,D}).\end{aligned}$$

This, together with the *Poincaré* inequality, implies the desired result. \square

5.2. **Neumann boundary value problem.** $(\hat{Q}_{\epsilon,h})$ reads as:

$$(5.4) \quad \begin{cases} \text{Find } \hat{u}_{\epsilon,h} \in V_h(D) \text{ such that} \\ (\hat{u}_{\epsilon,h}, v_h)_{1,\hat{\Omega}} + \epsilon(\nabla \hat{u}_{\epsilon,h}, \nabla v_h)_{\hat{\Omega}_1} = (f_h, v_h)_D, \quad \forall v_h \in V_h(D). \end{cases}$$

Lemma 5.3. $(\Omega \setminus \hat{\Omega}) \cup (\hat{\Omega} \setminus \Omega) \subset \Gamma_\delta$, $\delta = O(h^2)$ implies that:

$$(5.5) \quad \|v_h\|_{0,\Omega \setminus \hat{\Omega}} \leq Ch \|v_h\|_{1,\Omega} \quad (\text{or } h \|v_h\|_{1,\hat{\Omega}_1}).$$

$$(5.6) \quad \|v_h\|_{0,\hat{\Omega} \setminus \Omega} \leq Ch \|v_h\|_{1,\hat{\Omega}} \quad (\text{or } h \|v_h\|_{1,\Omega_1}).$$

(This lemma is quoted from [13].)

Theorem 5.4. There exists a unique solution $\hat{u}_{\epsilon,h}$ for (5.4). If $\tilde{f} - f_h = 0$ on $\Omega_1 \cap \hat{\Omega}_1$, and $u_{\epsilon,h}$ is the solution of (4.12), then we have,

$$(5.7) \quad \|u_{\epsilon,h} - \hat{u}_{\epsilon,h}\|_{1,\Omega \cup \hat{\Omega}} \leq Ch.$$

The proof is an analogue of that of Theorem 5.2, with using Lemma 5.1 and Lemma 5.3.

6. NUMERICAL EXPERIMENTS

Let $\tilde{u} \in H_0^1(D)$ be the zero extension of the solution $u \in H_0^1(\Omega)$ of the Dirichlet boundary value problem (1.1). Then, from the results of previous sections, we find that

$$\begin{aligned} \|\hat{u}_{\epsilon,h} - \tilde{u}\|_{1,D} &\leq \|\hat{u}_{\epsilon,h} - u_{\epsilon,h}\|_{1,D} + \|u_{\epsilon,h} - u_\epsilon\|_{1,D} + \|u_\epsilon - \tilde{u}\|_{1,D} \\ &= \|\hat{u}_{\epsilon,h} - u_{\epsilon,h}\|_{1,D} + \|u_{\epsilon,h} - u_\epsilon\|_{1,D} + \|u_\epsilon|_\Omega - u\|_{1,\Omega} + \|u_\epsilon\|_{1,\Omega_1} \\ &\leq Ch + C(\sqrt{h} + \sqrt{\epsilon}) + C\epsilon + C\epsilon \\ &\leq C(\sqrt{h} + \sqrt{\epsilon}). \end{aligned}$$

In addition, for the error in L^2 norm, we also have

$$\|\hat{u}_{\epsilon,h} - \tilde{u}\|_{0,D} \leq C(\epsilon + h + \sqrt{h\epsilon}).$$

So in our computation, to calculate L^2 and H^1 errors of $\hat{u}_{\epsilon,h} - \tilde{u}$ on D is sufficient to verify the theoretical results, which is more practical than computing the norm of $\hat{u}_{\epsilon,h}|_\Omega - u$ in Ω , because of the curved boundary of Ω . Now, let $\Omega = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 4\}$. The original problem reads as:

$$\begin{cases} -\Delta u = 4 \text{ in } \Omega, \\ u = 0 \text{ on } \Gamma. \end{cases}$$

The exact solution is $u = 4 - x^2 - y^2$. Let $D = \{(x, y) \mid -3 < x < 3, -3 < y < 3\}$, $\Omega \subset D$, and introduce a Cartesian mesh to D (see Fig. 4).

The error estimates are showed in Fig. 5 and Fig. 6, from which we see that for fixed ϵ , L^2 error behaves as Ch and H^1 error behaves as $C\sqrt{h}$. But at the same time, they also have lower bounds even if we allow h to become




FIGURE 3. On mixed boundary case




FIGURE 4. Ω , D and the mesh

arbitrarily small. The lower bounds are controlled by ϵ . And we can observe that L^2 error has the lower bound that behaves as $C\epsilon$, and H^1 error behaves as $C\sqrt{\epsilon}$. This confirms our theoretical results.




FIGURE 5. $\frac{\|u_{\epsilon,h} - \tilde{u}\|_{0,D}}{\|\tilde{u}\|_{0,D}}$ with different ϵ and h




FIGURE 6. $\frac{\|u_{\epsilon,h} - \tilde{u}\|_{1,D}}{\|\tilde{u}\|_{1,D}}$ with different ϵ and h

REFERENCES

- [1] Z. Chen and J. Zou, *Finite element methods and their convergence for elliptic and parabolic interface problems*, Numer. Math. **79** (1998), no. 2, 175202
- [2] S. Del Pino and O. Pironneau, *A fictitious domain based general PDE solver*, Numerical methods for scientific computing. Variational problems and applications, 2030, Internat. Center Numer. Methods Eng. (CIMNE), Barcelona, 2003.
- [3] P. Grisvard, *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics, **24**. Pitman (Advanced Publishing Program), Boston, MA, 1985.
- [4] P. Grisvard, *Singularities in boundary value problems* Recherches en Mathématiques Appliquées [Research in Applied Mathematics], 22. Masson, Paris; Springer-Verlag, Berlin, 1992.
- [5] V. Girault and R. Glowinski, *Error analysis of a fictitious domain method applied to a Dirichlet problem*, Japan J. Indust. Appl. Math. **12** (1995), no. 3, 487514.
- [6] V. Girault and P. -A. Raviart, *Finite element methods for Navier-Stokes equations. Theory and algorithms*, Springer Series in Computational Mathematics, **5**. Springer-Verlag, Berlin, 1986.
- [7] J. L. Lions and E. Magenes, *Non-homogeneous boundary value problems and applications. Vol. I*, Springer-Verlag, New York-Heidelberg, 1972.
- [8] J. L. Lions, *Une remarque sur les problèmes d'évolution non linéaires dans des domaines non cylindriques*, Rev. Roumaine Math. Pures Appl. **9** 1964 1118.
- [9] A. L. Mignot, *Méthodes d'approximation des solutions de certains problèmes aux limites linéaires. I.* (French) Rend. Sem. Mat. Univ. Padova **40** 1968 1138.
- [10] B. Maury, *Numerical analysis of a finite element/volume penalty method*, SIAM J. Numer. Anal. **47** (2009), no. 2, 11261148.

- [11] J. Nečas, *Les méthodes directes en théorie des équations elliptiques*, (French) Masson et Cie, Éditeurs, Paris; Academia, Éditeurs, Prague 1967 351 pp.
- [12] K. Ohmori and N. Saito, *On the convergence of finite element solutions to the interface problem for the Stokes system*, J. Comput. Appl. Math. **198** (2007), no. 1, 116128.
- [13] M. Tabata, *Uniform solvability of finite element solutions in approximate domains*, Japan J. Indust. Appl. Math. **18** (2001), no. 2, 567585.
- [14] J. Wloka, *Partial differential equations*, Translated from the German by C. B. Thomas and M. J. Thomas. Cambridge University Press, Cambridge, 1987.
- [15] S. Zhang, *A domain embedding method for mixed boundary value problems*, C. R. Math. Acad. Sci. Paris **343** (2006), no. 4, 287290.
- [16] S. Zhang, *Analysis of finite element domain embedding methods for curved domains using uniform grids*, SIAM J. Numer. Anal. **46** (2008), no. 6, 28432866.
- [17] A. Ženíšek, *Nonlinear elliptic and evolution problems and their finite element approximations*, Computational Mathematics and Applications. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London, 1990.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1
KOMABA MEGURO-KU TOKYO 153-8914, JAPAN

E-mail address: zhoug@ms.u-tokyo.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1
KOMABA MEGURO-KU TOKYO 153-8914, JAPAN

UTMS

- 2011–15 Yusaku Tiba: *Kobayashi hyperbolic imbeddings into toric varieties.*
- 2011–16 Yuko Hatano, Junichi Nakagawa, Shengzhang Wang and Masahiro Yamamoto: *Determination of order in fractional diffusion equation.*
- 2011–17 Takashi Tsuboi: *Homeomorphism groups of commutator width one.*
- 2011–18 J. Noguchi: *Another direct proof of Oka's theorem (Oka IX).*
- 2011–19 M. Abe, S. Hamano and J. Noguchi: *On Oka's extra-zero problem.*
- 2011–20 Oleg Yu. Imanuvilov and Masahiro Yamamoto : *On reconstruction of Lamé coefficients from partial Cauchy data.*
- 2011–21 Pen-Yuan Hsu, Yasunori Maekawa: *On nonexistence for stationary solutions to the Navier-Stokes equations with a linear strain.*
- 2011–22 Hitoshi Kitada: *Timeless system, superluminal phenomena, dark matter and Big Bang.*
- 2011–23 Takashi Kato: *An optimal execution problem with market impact.*
- 2011–24 Takashi Kato: *An optimal execution problem in geometric Ornstein Uhlenbeck price process.*
- 2011–25 Takashi Kato: *Theoretical sensitivity analysis for quantitative operational risk management.*
- 2011–26 Guanyu Zhou and Norikazu Saito: *Analysis of the fictitious domain method with penalty for elliptic problems.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153-8914, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012