


UTMS 2016–5

December 20, 2016

**Well-posedness of Hamilton-Jacobi
equations with
Caputo's time-fractional derivative**

by

Yoshikazu GIGA and Tokinaga NAMBA



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

WELL-POSEDNESS OF HAMILTON-JACOBI EQUATIONS WITH CAPUTO'S TIME-FRACTIONAL DERIVATIVE

YOSHIKAZU GIGA AND TOKINAGA NAMBA

ABSTRACT. A Hamilton-Jacobi equation with Caputo's time-fractional derivative of order less than one is considered. The notion of a viscosity solution is introduced to prove unique existence of a solution to the initial value problem under periodic boundary conditions. For this purpose comparison principle as well as Perron's method is established. Stability with respect to the order of derivative as well as the standard one is studied. Regularity of a solution is also discussed. Our results in particular apply to a linear transport equation with time-fractional derivatives with variable coefficients.

1. INTRODUCTION

Let $\alpha \in (0, 1]$ and $0 < T < \infty$ be given constants. We consider the initial-value problem for the Hamilton-Jacobi equation of the form

$$(1.1) \quad \partial_t^\alpha u + H(t, x, u, Du) = 0 \quad \text{in } (0, T] \times \mathbb{T}^d =: Q_T$$

and

$$(1.2) \quad u|_{t=0} = u_0 \quad \text{in } \mathbb{T}^d.$$

Here $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$ is a d -dimensional torus, $u : \overline{Q_T} \rightarrow \mathbb{R}$ is an unknown function and $H : \overline{Q_T} \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ is a given function called a *Hamiltonian*. Moreover, Du denotes the spatial gradient, i.e., $Du = (\partial u / \partial x_1, \dots, \partial u / \partial x_d)$ and $\partial_t^\alpha u$ denotes Caputo's (time-)fractional derivative which is defined by

$$(1.3) \quad (\partial_t^\alpha f)(t) := \begin{cases} \frac{1}{\Gamma(1-\alpha)} \int_0^t \frac{f'(s)}{(t-s)^\alpha} ds & \text{for } \alpha \in (0, 1), \\ f'(t) & \text{for } \alpha = 1, \end{cases}$$

where $\Gamma(\cdot)$ is the usual gamma function. Here, throughout this paper, a function v on \mathbb{T}^d is regarded as a function defined on \mathbb{R}^d with \mathbb{Z}^d -periodically, i.e., $v(x+z) = v(x)$ for all $x \in \mathbb{R}^d$ and $z \in \mathbb{Z}^d$. Although some part of our arguments can be easily extended to other boundary conditions we now restrict ourselves only under periodic boundary conditions.

The goal of this paper is to extend a notion of viscosity solutions to (1.1)-(1.2) and to establish unique existence, stability and some regularity results of viscosity solutions for (1.1)-(1.2). Here we will consider only for $\alpha \in (0, 1)$ since the case of $\alpha = 1$ has been well studied. All results excepts for Section 6 and Section 7 will be established under the following assumptions:

(A1) $H : \overline{Q_T} \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ is continuous,

2010 *Mathematics Subject Classification.* 26A33; 35D40; 35F21.

Key words and phrases. Viscosity solutions; Hamilton-Jacobi equations; Caputo's time-fractional derivatives.

(A2) there is a modulus $\omega : [0, \infty) \rightarrow [0, \infty)$ such that

$$|H(t, x, r, p) - H(t, y, r, p)| \leq \omega(|x - y|(1 + |p|))$$

for all $(t, x, y, r, p) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$,

(A3) $r \mapsto H(t, x, r, p)$ is nondecreasing for all $(t, x, p) \in [0, T] \times \mathbb{T}^d \times \mathbb{R}^d$,

(A4) $u_0 : \mathbb{T}^d \rightarrow \mathbb{R}$ is continuous.

We emphasize that these assumptions are fairly standard for $\alpha = 1$. Of course there might be several generalizations but we do not touch them. Note that we do not assume coercivity, i.e.,

$$(1.4) \quad \liminf_{r \rightarrow \infty} \{H(t, x, r, p) \mid (t, x, r) \in Q_T \times \mathbb{R}, |p| \geq r\} = +\infty.$$

Hence our results apply to a transport equation

$$(1.5) \quad \partial_t^\alpha u + b \cdot Du = 0$$

for $b = b(t, x) : \overline{Q_T} \rightarrow \mathbb{R}^d$.

Since a notion of viscosity solutions was introduced by Crandall and Lions [8], its theory has developed rapidly and by now there is a large number of literature. The reader is referred to [1], [4] and [24] for basic theory and to [7] and [12] for more advanced theory. The theory of viscosity solutions had been applied initially to local partial differential equations (pdes for short) and soon has been extended by Soner [39] to pdes with space-fractional derivatives which are defined non-locally. See also [5], [2] and references therein. In these papers the authors are commonly interested in Lévy operators, which can be represented (formally) as

$$(1.6) \quad g[f](x) = - \int_{\mathbb{R}^d} \left(f(x+z) - f(x) - \frac{Df(x) \cdot z}{1 + |z|^2} \right) d\mu(z),$$

where $d\mu$ is the Lévy measure. An example of Lévy operators is the fractional Laplacian:

$$(1.7) \quad (-\Delta)^\alpha f(x) = C \int_{\mathbb{R}^d} \frac{f(x) - f(y)}{|x - y|^{d+2\alpha}} dy,$$

where C is a constant depending on d and α .

Above works are motivated from applied fields such as physics, engineering and finances. Applicabilities of pdes with time-fractional derivatives has been discussed by many researchers in wide fields as well; [10], [11], [38] and [40] for instance. We here refer to several mathematical works for pdes with Caputo's time-fractional derivative (CTFD for short) in order to motivate our research. Although many definitions of a different kind of fractional derivatives have been suggested, we will not touch them in this paper and instead the reader is referred to [9], [13], [23], [22], [35], [37] and [44]. A typical example of pdes with CTFD is

$$(1.8) \quad \partial_t^\alpha u + L(u) = F,$$

where L consists of a symmetric uniformly elliptic operator and a transport term and $F = F(t, x)$ is a given function. This can be considered as a equation describing diffusion phenomena in complex media like fractals and then is called *anomalous diffusion* or *singular diffusion*. There seem to be several previous works for (1.8) (see [36] and references therein) and Luchko's works have a close relationship with ours. He established a maximum principle for Caputo's fractional derivative in [30] and, based on it, proved a uniqueness of classical solutions for an initial-boundary value problem of (1.8) with the type of $L(u) = -\operatorname{div}(p(x)Du) + q(x)u$, where typically

p is smooth and uniformly positive with continuous q . In [31] he established an existence of classical solutions for same equations as well. His research has been continued in a work by Sakamoto and Yamamoto [36], which is a pioneer work in the theory of weak solutions for (1.8). They defined weak solutions in the sense of distribution for a similar equation as one Luchko considered and established well-posedness in order to consider inverse problems. Researches on this line have been growing rapidly; see, e.g., [29] for multi-term time-fractional derivatives and [32] for (1.1) with nonlinear source terms.

Anomalous diffusion equations are modelled by the continuous-time random walk (CTRW for short) introduced by Montroll and Weiss ([34]). More recently, Kolokoltsov and Veretennikova ([25]) extended the notion of CTRW so that its processes can be controlled and then derived (heuristically) Hamilton-Jacobi-Bellman equations with CTFD, fractional Laplacian and some additional term. We note that this does not include second order spatial-derivative. In [26] they also defined mild solutions that belong to $C^1([0, T]; C_\infty^1(\mathbb{R}^d))$ and proved well-posedness for an initial-value problem of

$$\partial_t^\alpha u = -a(-\Delta)^{\beta/2}u + H(t, x, Du).$$

Here $C_\infty^1(\mathbb{R}^d)$ is a set of C^1 functions that decreasing rapidly at infinity and $\beta \in (1, 2]$ and $a > 0$ are given constants. In the case of [26] it is assumed that mild solutions can be defined thanks to fractional Laplacian. On the other hand we need to deal with solutions in the weak sense also in the space direction, so viscosity solutions are expected to be reasonable in our case. However, for this direction there seems to be only one result by Allen ([3]) for CTFD by a viscosity approach as far as we know. He has discussed regularity problems of viscosity solutions for space-time nonlocal equations of the form

$$\partial_t^\alpha u - \sup_i \inf_j \left(\int_{\mathbb{R}^d} \frac{u(x+y) - u(x)}{|y|^{d+2\sigma}} a^{ij}(t, x, y) dy \right) = f,$$

where a_{ij} is positive, bounded function that is symmetric with respect to the third variable and f is a given function. His definition of viscosity solutions seems to be based on one of pdes with space-fractional derivatives mentioned above. However, the form of Caputo's fractional derivative is different from (1.6) or (1.7). Thus, it was not clear why Allen's definition of viscosity solutions is useful.

We explained so far second-order pdes or first-order pdes including fractional Laplacian with CTFD. For first-order pdes with CTFD but without any higher-order terms than one such as (1.5), a formula of solution for (1.5) is given by Mainardi, Mura and Pagnini ([33]) for instance, but for equations with constant coefficients. However, there seems to be no theoretical studies. Our aim of this research is to construct the synthetic theory of viscosity solutions so that (fully nonlinear) pdes with CTFD mentioned above can be considered. However, extensions to second order equations expects some technical issues, so we only treat first order equations in this paper as the first step. For second order problems the reader is referred to one of forthcoming papers of the second author.

We motivate our definition of viscosity solutions (Definition 2.5) by recalling the case of $\alpha = 1$. Let us suppose that u is a classical subsolution of (1.1), that is,

$$(\partial_t^\alpha u)(t, x) + H(t, x, u(t, x), Du(t, x)) \leq 0$$

for all $(t, x) \in Q_T$ and that

$$(1.9) \quad \max_{[0, T] \times \mathbb{R}^d} (u - \phi) = (u - \phi)(\hat{t}, \hat{x})$$

for a text function ϕ . The classical maximum principle in space implies that $Du = D\phi$ at (\hat{t}, \hat{x}) . With respect to time, the maximum principle for CTFD ([30, Theorem 1]) implies that $\partial_t^\alpha u \geq \partial_t^\alpha \phi$ at (\hat{t}, \hat{x}) . Hence, that u is a classical subsolution yields to

$$(1.10) \quad (\partial_t^\alpha \phi)(\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) \leq 0.$$

In the spirit of the case of $\alpha = 1$, it is natural to define weak subsolutions of (1.1) by (1.10). Let us call it *provisional subsolutions* in this paper. Similarly, if u is a classical supersolution and we replace the maximum by a minimum in (1.9), then the converse inequality of (1.10) is led. Then let us call such u *provisional supersolutions* of (1.1). Let us call u a *provisional solution* of (1.1) if it is a both provisional sub- and supersolution of (1.1).

Provisional solutions looks easy to deal with but it is technically difficult to establish a comparison principle, so we do not know whether it is a proper notion of solution or not (see Section 7 for detail). One reason is that the so-called *doubling variable method* (see, e.g., [12, Section 3.3]) does not work and a main problem is, roughly speaking, that $(\partial_t^\alpha \phi)(\hat{t}, \hat{x})$ in (1.10) is not an appropriate substitute of $(\partial_t^\alpha u)(\hat{t}, \hat{x})$. In a proof of comparison principle in the theory of viscosity solutions, we often aim to derive a contradiction by using the doubling variable method under a suitable supposition. For provisional sub/supersolutions, we cannot derive a contradiction because of unnecessary values caused by $\partial_t^\alpha \phi$.

This fact makes us realize that it is necessary to bring a function that has a closer value to $\partial_t^\alpha u$ at each point. After integration by parts and changing the variable of integration, we get another form of $\partial_t^\alpha u$ of the form

$$(1.11) \quad K_0[u](t, x) = \frac{u(t, x) - u(0, x)}{t^\alpha \Gamma(1 - \alpha)} + \frac{\alpha}{\Gamma(1 - \alpha)} \int_0^t (u(t, x) - u(t - \tau, x)) \frac{d\tau}{\tau^{\alpha+1}}.$$

Here the integral is interpreted as an improper integral

$$(1.12) \quad \int_0^t (u(t, x) - u(t - \tau, x)) \frac{d\tau}{\tau^{\alpha+1}} = \lim_{r \searrow 0} \int_r^t (u(t, x) - u(t - \tau, x)) \frac{d\tau}{\tau^{\alpha+1}}.$$

It is easy to see that $\partial_t^\alpha u = K_0[u]$ if u is smooth. Since the convergence of (improper) integration (1.12) is not trivial, we apply test functions instead of unknown functions near the singular time, i.e., the lower end of interval. This idea is taken in our definition. By the way, the integral term in (1.11) is close to the fractional Laplacian (1.7) (or Lévy operators (1.6) without a derivative term in an integrand). Our definition is able to be regarded as an analogy of one for pdes with space-fractional derivatives referred above. It is worth clarifying a relationship between our way to handle CTFD and Allen's ([3]). To give a definition of viscosity solutions he introduced the function

$$\widetilde{K}_0[u](t, x) = \frac{\alpha}{\Gamma(1 - \alpha)} \int_{-\infty}^t \frac{\tilde{u}(t, x) - \tilde{u}(\tau, x)}{(t - \tau)^\alpha} d\tau,$$

where $\tilde{u} : (-\infty, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ is an extension of u defined by $\tilde{u}(t, \cdot) = u(0, \cdot)$ for $t < 0$. He defined a viscosity sub/supersolution by substituting a test function near

the singularity of the integral. As a result, our way is the same as Allen's since clearly $K_0[u] = \widetilde{K}_0[u]$.

Let us give our strategy of proofs in this paper. We will show that (1.12) exists as a finite number at points such that $u - \phi$ attains a maximum/minimum (Lemma 2.9), where ϕ is a test function. This is an analogy of [5] or [6, Lemma 3.3] and it is a key fact in a proof of our comparison principle. An idea of the proof of comparison principle is the same as usual ones, that is, doubling variable method under contradiction. We get a term that yields a contradiction from a non-integration term in a difference of (1.11) for viscosity sub- and supersolutions.

A proof of existence of viscosity solutions follows Perron's method, that is, a construction of maximal subsolutions. Since (1.1) is nonlocal, we need some efforts to handle nonlocal terms compared with the case of local equations. For this purpose we will employ the idea used in [16, Theorem 3] for example.

We will establish stability results under limit operation from two perspectives. One of them is for a family of solutions of (1.1) with a Hamiltonian depending on a parameter. Our statement and proof are almost the same as for $\alpha = 1$ (see, e.g., [4]). The other stability discussed here is the case when time-derivative's orders are regarded as parameters. The latter can be proved under the same idea as the former by defining analogous functions of half-relaxed limits.

We will show that viscosity solutions are Lipschitz continuous in space and α -Hölder continuous in time under some additional assumptions on H and u_0 . When the regularity problems for viscosity solutions are discussed, the coercivity condition is often assumed. However, transport equations are not coercive. In view of applications we will derive the above regularity results without the coercivity assumption. Our proofs follow basically ones for $\alpha = 1$ ([1]) but a proof of the temporal regularity may be not standard. We will construct a viscosity solution of (1.1)-(1.2) that is α -Hölder continuous in time by Perron's method for a family of viscosity subsolutions of (1.1)-(1.2) that is α -Hölder continuous in time. In this argument we should be carefully for a dependence of the Hölder constant of viscosity subsolutions. We will show that viscosity solutions are Hölder continuous in time at the initial time, where its Hölder constant depends only on T , α , H and u_0 . By restricting viscosity subsolutions to Hölder continuous functions with such a constant, we will obtain viscosity solution with the desired regularity.

Our results are new even for transport equations (1.5) with variable coefficients although a formula of solution for constant coefficients is known only in the one dimensional case (see Section 6). For this reason it is worth summarizing here.

Theorem 1.1. *Let $b : \overline{Q_T} \rightarrow \mathbb{R}^d$ be a continuous function. Assume that there is constants $C_1 > 0$ and $C_2 > 0$ such that*

$$(1.13) \quad |b(t, x) - b(t, y)| \leq C_1 |x - y|$$

and

$$(1.14) \quad |u_0(x) - u_0(y)| \leq C_2 |x - y|$$

for all $(t, x, y) \in (0, T] \times \mathbb{R}^d \times \mathbb{R}^d$. Then there exists at most one viscosity solution $u \in C(\overline{Q_T})$ of (1.5)-(1.2). Moreover there exists a constant $C_3 > 0$ such that

$$(1.15) \quad |u(t, x) - u(s, y)| \leq C_3 (|t - s|^\alpha + |x - y|)$$

for all $(t, x, s, y) \in ([0, T] \times \mathbb{R}^d)^2$.

If one does not require (1.15), conditions (1.13) and (1.14) can be weakened so that Hamiltonian $H(t, x, p) = b(t, x) \cdot p$ satisfies (A1) and (A2).

We finally compare with viscosity solutions with weak solutions in the sense of distribution (weak solution for short) and mention several open problems. A weak solution for linear second order problems (1.8) given by Sakamoto and Yamamoto ([36]) was constructed by Galerkin method. Since approximate equations have no comparison principle, it is difficult to compare two notions under the current circumstances. Of course, the case of $\alpha = 1$ has the same difficulty and, even such simple looking case, there seems to be few literatures ([15], [19], [20] and [21]). In order to overcome such a difficulty, analyses for further regularities of weak solutions in the both senses will be needed.

As another direction of researches for weak solutions of pdes with CTFD, we should mention fractional derivative of the form

$$(1.16) \quad (D_t^\alpha f)(t) := \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t \frac{f(\tau) - f(0)}{(t-\tau)^{\alpha+1}} d\tau.$$

The original definition of Caputo's fractional derivative by himself was actually given as this form and hence this derivative is also called Caputo's fractional derivative. We note that $D_t^\alpha f = \partial_t^\alpha f$ almost everywhere on $[0, T]$ if f is absolutely continuous on $[0, T]$. See [9, Chapter 3] for a brief history of Caputo's fractional derivative and the above relationship between two definitions. There are some works for weak solutions of pdes with (1.16). Zacher ([41]) considered abstract evolutional equations of parabolic type including

$$D_t^\alpha u - \operatorname{div}(ADu) + b \cdot Du + cu = 0$$

and, by introducing a notion of a weak solution, he established a unique existence. Here, $A = A(t, x)$ is a symmetric and positive defined matrix-valued function with L^∞ elements and $b = b(t, x)$ and $c = c(t, x)$ are L^∞ functions. See [42], [43] and [28] for related works. An analysis of weak solutions for pdes with (1.16) involves the problem of the trace $u(0, \cdot)$ of u up to $t = 0$ since $D_t^\alpha u$ includes the value $u(0, \cdot)$. This needs some regularity up to $t = 0$ which forced as to restrict range of α , say, for example $\alpha > 1/2$ or regularity of some of given functions A , b and c compared with the case $\alpha = 1$ ([27]). We note that such a trace problem was not considered in [41]; moreover, assumptions of [41] seem to be too weak to get necessary regularity. In view of such restrictions, our viscosity solutions might look a better notion of weak solutions since we are able to obtain a continuous (viscosity) solution for every $\alpha \in (0, 1)$ with no special assumptions on H . However, we cannot compare two notions since it is not guaranteed that our solution u is absolutely continuous in time, so it is not clear whether or not $D_t^\alpha u = \partial_t^\alpha u$ for our solution. Even for this problem, further analyses from both aspects of viscosity solutions and weak solutions are needed.

This paper is organized as follows: In Section 2 we give a definition of viscosity solutions after and summarize some facts used in the other sections. In Section 3 we prove a comparison principle and in Section 4 we establish an existence result. In Section 5 we prove two types of stability results and in Section 6 we study regularity problem for (1.1). Finally, in Section 7 we give a definition of provisional solutions as another possible notion of weak solutions and mention the technical difficulty for them.

2. DEFINITION AND PROPERTIES OF SOLUTIONS

In this section we assume that Hamiltonian H is merely continuous on $\overline{Q_T} \times \mathbb{R} \times \mathbb{R}^d$.

2.1. Preliminaries. To give a definition of viscosity solutions we first introduce a function space of the type

$$\mathcal{C}^1([a, b] \times O) := \{\phi \in C^1((a, b] \times O) \cap C([a, b] \times O) \mid \partial_t \phi(\cdot, x) \in L^1(a, b) \text{ for every } x \in O\}.$$

Here $a, b \in \mathbb{R}$ are constants such that $a < b$, O is a domain in \mathbb{R}^d , \mathbb{T}^d and $\mathbb{R}^d \times \mathbb{R}^d$ and $L^1(a, b)$ is the space of Lebesgue integrable functions on (a, b) . Note that $u \in \mathcal{C}^1([a, b] \times O)$ may not be C^1 up to $t = a$. This space will be used as a space of test functions as well as of classical solutions of (1.1)-(1.2). Here we define classical solutions of (1.1)-(1.2) as follows:

Definition 2.1 (Classical solutions). A function $u \in \mathcal{C}^1(\overline{Q_T})$ is called a *classical solution* of (1.1)-(1.2) if $u(0, \cdot) = u_0$ on \mathbb{T}^d and

$$(\partial_t^\alpha u)(t, x) + H(t, x, u(t, x), Du(t, x)) = 0$$

for all $(t, x) \in Q_T$.

Note that $\partial_t^\alpha \phi$ is bounded in $(0, T] \times \mathbb{R}^d$ if $\phi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$; see [9, Theorem 2.1]. We are tempted to use $\mathcal{C}^1([a, b] \times O)$ as a space of classical solutions since the integrability condition for $\partial_t \phi(\cdot, x)$ is satisfied if ϕ belongs to it: $\mathcal{C}^1([a, b] \times O) \subset \mathcal{C}^1([a, b] \times O)$. However, the class $\mathcal{C}^1([a, b] \times O)$ is too narrow to define classical solutions since it is necessary to include functions that have a fractional power with respect to time at the initial time such as t^α . That is why we do not assume the differentiability at the initial time.

Example 2.2. As an example let us consider a simple ordinary differential equation of the form

$$\partial_t^\alpha f + f = 0 \quad \text{in } (0, \infty)$$

with prescribed data $f(0) = c \in \mathbb{R}$. According to [9, Theorem 4.3] a solution of this equation is given as $f(t) = cE_\alpha(-t^\alpha)$, where E_α is the Mittag-Leffler function defined by

$$E_\alpha(z) := \sum_{j=0}^{\infty} \frac{z^j}{\Gamma(j\alpha + 1)}.$$

In particular, $E_{1/2}(-\sqrt{t}) = e^t \operatorname{erfc}(\sqrt{t})$, where erfc is the complementary error function defined by

$$\operatorname{erfc}(z) := \frac{2}{\sqrt{\pi}} \int_z^\infty e^{-t^2} dt.$$

The function f is not differentiable at $t = 0$ though it is continuous up to $t = 0$; we leave the verification to the reader. Therefore classical solutions of equations with Caputo's (time-)fractional derivative are not always differentiable at the initial time even if an initial datum is smooth.

For a measurable function $f : [0, T] \rightarrow \mathbb{R}$ we define functions $J_r[f], K_r[f] : (0, T] \rightarrow \mathbb{R}$ by

$$J_r[f](t) := \frac{\alpha}{\Gamma(1 - \alpha)} \int_0^r (f(t) - f(t - \tau)) \frac{d\tau}{\tau^{\alpha+1}}$$

and

$$K_r[f](t) := \frac{f(t) - f(0)}{t^\alpha \Gamma(1 - \alpha)} + \frac{\alpha}{\Gamma(1 - \alpha)} \int_r^t (f(t) - f(t - \tau)) \frac{d\tau}{\tau^{\alpha+1}}$$

with a parameter $r \in (0, t)$. For a measurable function $f : [0, T] \times \mathbb{R}^\ell \rightarrow \mathbb{R}$ with $\ell \geq 1$ we define $J_r[f], K_r[f] : (0, T] \times \mathbb{R}^\ell \rightarrow \mathbb{R}$ by $J_r[f](t, x) := J_r[f(\cdot, x)](t)$ and $K_r[f](t, x) := K_r[f(\cdot, x)](t)$ for $(t, x) \in (0, T] \times \mathbb{R}^\ell$.

Proposition 2.3 (Integration by parts). *Let $f : [a, T] \rightarrow \mathbb{R}$ be a function such that $f \in C^1((a, T]) \cap C([a, T])$ and $f' \in L^1(a, T)$, where $a < T$. Then*

$$\frac{1}{\Gamma(1 - \alpha)} \int_a^t \frac{f'(\tau)}{(t - \tau)^\alpha} d\tau = \frac{f(t) - f(a)}{(t - a)^\alpha \Gamma(1 - \alpha)} + \frac{\alpha}{\Gamma(1 - \alpha)} \int_0^{t-a} (f(t) - f(t - \tau)) \frac{d\tau}{\tau^{\alpha+1}}.$$

Proof. The left-hand side (multiplied by $\Gamma(1 - \alpha)$) can be calculated as

$$\begin{aligned} \int_a^t \frac{f'(\tau)}{(t - \tau)^\alpha} d\tau &= \int_a^t \frac{\frac{d}{d\tau}(f(\tau) - f(t))}{(t - \tau)^\alpha} d\tau \\ &= \left[\frac{f(\tau) - f(t)}{(t - \tau)^\alpha} \right]_a^t - \alpha \int_a^t \frac{f(\tau) - f(t)}{(t - \tau)^{\alpha+1}} d\tau \\ &= \lim_{\tau \rightarrow t} \frac{f(\tau) - f(t)}{(t - \tau)^\alpha} + \frac{f(t) - f(a)}{(t - a)^\alpha} + \alpha \int_a^t \frac{f(t) - f(\tau)}{(t - \tau)^{\alpha+1}} d\tau. \end{aligned}$$

Thanks to the smoothness of f , the first term vanishes. By the change of variable $s := t - \tau$ we obtain the desired result. \square

Let us share some words for an integral

$$I[f](t) := \int_a^b f(t, \tau) \frac{d\tau}{\tau^{\alpha+1}}$$

for constants $a, b \in \mathbb{R}$ with $0 \leq a < b \leq T$ and a measurable function $f : [0, T] \times [0, T] \rightarrow \mathbb{R}$. We say that the integral $I[f]$ makes sense if either $I[f^+]$ or $I[f^-]$ is finite (in the sense of Lebesgue integrals) and that $I[f]$ exists if both $I[f^\pm]$ are finite. Here $f^\pm := \max\{\pm f, 0\}$. It is necessary to pay attention when $a = 0$. Then we regard $I[f]$ as an improper integral by $I[f](t) = \lim_{r \searrow 0} I_r[f]$, where

$$I_r[f](t) = \int_r^b f(t, \tau) \frac{d\tau}{\tau^{\alpha+1}}.$$

Thus $I[f]$ exists if $I_r[f^\pm]$ are finite for each r and $\lim_{r \searrow 0} I_r[f^\pm]$ exist as a finite number. Note that, if $\tau \mapsto |f(t, \tau)|/\tau^{\alpha+1}$ is integrable on $(0, b)$, then $I[f]$ exists and it agrees with the Lebesgue integral; this is a direct consequence of the dominated convergence theorem. We abuse above words not only for J_r but also for K_r including a non-integration term.

For a set $E \subset \mathbb{R}^\ell$ with $\ell \geq 1$, let $USC(E)$ and $LSC(E)$ be sets of real-valued upper and lower semicontinuous functions on E , respectively. Note that semicontinuous functions are measurable.

Proposition 2.4 (Properties of J_r and K_r). *Let $f \in USC([0, T])$ (resp. $LSC([0, T])$) and $g \in C^1((0, T])$. Then*

- (i) for each $t \in (0, T]$, $J_r[g](t)$ exists for all $r \in (0, t)$,
- (ii) for each $t \in (0, T]$, $K_r[f](t)$ makes sense and is bounded from below (resp. above) for all $r \in (0, t)$,

- (iii) $K_0[f](\hat{t})$ makes sense and is bounded from below (resp. above) if $f - g$ attains a maximum (resp. minimum) at $\hat{t} \in (0, T]$ over $(0, T]$, i.e.,

$$\sup_{(0, T]} (f - g) = (f - g)(\hat{t}) \quad (\text{resp.} \quad \inf_{(0, T]} (f - g) = (f - g)(\hat{t})),$$

Moreover for each $j \geq 0$ let $t_j \in (0, T]$, $r_j \in (0, t_j)$ and $\alpha_j \in (0, 1)$ be sequences such that $\lim_{j \rightarrow \infty} (t_j, r_j, \alpha_j) = (\hat{t}, \hat{r}, \alpha) \in (0, T] \times \mathbb{R}^d \times [0, \hat{t}) \times (0, 1)$. Let $J_r^{\alpha_j}$ denote a function J_r associated with $\alpha = \alpha_j$. Then

$$(iv) \lim_{j \rightarrow \infty} J_{r_j}^{\alpha_j}[g](t_j) = J_{\hat{r}}^{\alpha}[g](\hat{t}).$$

Proof. (i) Fix $t \in (0, T]$ and $r \in (0, t)$ arbitrarily. Since g is Lipschitz continuous near t due to the smoothness of g , for some constant $C > 0$

$$\int_0^r |g(t) - g(t - \tau)| \frac{d\tau}{\tau^{\alpha+1}} \leq \int_0^r C\tau \frac{d\tau}{\tau^{\alpha+1}} = \frac{Cr^{1-\alpha}}{1-\alpha}.$$

Our assertion follows immediately from this.

(ii) Fix $t \in (0, T]$ and $r \in (0, t)$ arbitrarily. Assume that $f \in USC([0, T])$. Then f attains a maximum and hence

$$f(t) - \max_{[0, T]} f \leq f(t) - f(t - \tau)$$

for all $\tau \in (r, t)$. The left-hand side multiplied by $\tau^{-\alpha-1}$ is integrable on (r, t) since we integrate away from $\tau = 0$. Therefore the negative part $[f(t) - f(t - \tau)]^- / \tau^{\alpha+1}$ is integrable on (r, t) . This implies that $K_r[f](t)$ makes sense and is bounded from below. The similar argument is applied for $f \in LSC([0, T])$. The above yields our assertion.

(iii) Define $h := g + (f - g)(\hat{t})$ and

$$(2.1) \quad v(\tau) := \begin{cases} h(\hat{t}) - h(\hat{t} - \tau) & \text{for } \tau \in [0, \hat{t}/2], \\ f(\hat{t}) - f(\hat{t} - \tau) & \text{for } \tau \in (\hat{t}/2, \hat{t}]. \end{cases}$$

Since $f - h$ attains a maximum at \hat{t} over $(0, T]$, we see $f \leq h$ on $(0, T]$. In addition, $(f - h)(\hat{t}) = 0$ and thus

$$h(\hat{t}) - h(\hat{t} - \tau) \leq f(\hat{t}) - f(\hat{t} - \tau)$$

on $(0, \hat{t})$. By (i) and a similar argument as the proof of (ii) with $r = \hat{t}/2$ it turns out that the negative part $v^-(\tau) / \tau^{\alpha+1}$ is integrable on $(0, \hat{t})$, so is $[f(\hat{t}) - f(\hat{t} - \tau)]^- / \tau^{\alpha+1}$ since $v(\tau) \leq f(\hat{t}) - f(\hat{t} - \tau)$ on $(0, \hat{t})$. This yields our assertion for $f \in USC([0, T])$. Another can be proved similarly.

(iv) Thanks to the smoothness of g the dominated convergence theorem can be applied and ensures our assertion. More precisely, since $\inf_{j \geq 0} (t_j - r_j) > 0$ and $g \in C^1((0, T])$, there exists a constant $C_1 > 0$ such that $|g(t_j) - g(t_j - \tau)| \leq C_1\tau$ on $(0, r_j)$. In particular, we may assume that C_1 does not depend on j since $\lim_{j \rightarrow \infty} t_j = \hat{t} > 0$. Thus we have

$$(2.2) \quad \sup_{j \geq 0} (|g(t_j) - g(t_j - \tau)| \mathbf{1}_{(0, r_j)}(\tau)) \tau^{-\alpha-1} \leq C_1 \mathbf{1}_{(0, \hat{r})}(\tau) \tau^{-\alpha}$$

for all $\tau \in [0, T]$. Here $\mathbf{1}_I$ is the indicator function on an interval I , i.e., $\mathbf{1}_I = 1$ in I and 0 elsewhere. The right-hand side is integrable on $[0, T]$. It remains to check the convergence of $(g(t_j) - g(t_j - \cdot)) \mathbf{1}_{[0, r_j]}(\cdot)$ but this is obvious. \square

2.2. Definition of solutions. We now give our definition of viscosity solutions for (1.1).

Definition 2.5 (Viscosity solutions). A function $u \in USC(\overline{Q_T})$ (resp. $LSC(\overline{Q_T})$) is called a *viscosity subsolution* of (1.1) if, for any constants $a, b \in [0, T]$ with $a < b$ and an open ball B in \mathbb{R}^d ,

$$(2.3) \quad J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) \leq 0 \quad (\text{resp. } \geq 0)$$

whenever $u - \phi$ attains a maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$.

If $u \in C(\overline{Q_T})$ is both a viscosity sub- and supersolution of (1.1), then we call u a *viscosity solution* of (1.1).

Remark 2.6. (i) For an arbitrary function $u : Q_T \rightarrow \mathbb{R}$ an *upper semicontinuous envelope* $u^* : \overline{Q_T} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ and a *lower semicontinuous envelope* $u_* : \overline{Q_T} \rightarrow \mathbb{R} \cup \{\pm\infty\}$ are defined by

$$u^*(t, x) := \lim_{\delta \searrow 0} \sup \{u(s, y) \mid (s, y) \in Q_T \cap \overline{B_\delta(t, x)}\}$$

and $u_* := -(-u)^*$. Here $B_\delta(t, x)$ is an open ball of radius δ centered at (t, x) and $\overline{B_\delta(t, x)}$ is its closure. As for $\alpha = 1$ a viscosity sub- and subsolution of (1.1) can be defined for arbitrary functions $u : Q_T \rightarrow \mathbb{R}$ by using u^* (for a subsolution) and u_* (for a supersolution) in Definition 2.5, where it is further assumed that $u^* < +\infty$ and $u_* > -\infty$ on $\overline{Q_T}$; cf. [12, Definition 2.1.1]. Note that functions u^* and u_* are upper semicontinuous and lower semicontinuous on $\overline{Q_T}$, respectively (see, e.g., [4, Proposition V.2.1]) so they are measurable.

(ii) Although we restrict ourselves for spatially periodic functions, our definition can be easily extended for $(0, T] \times \Omega$, where Ω is a domain in \mathbb{R}^d . In fact, the comparison principle holds for a general bounded domain with necessary modifications.

If a viscosity subsolution (resp. supersolution) u of (1.1) satisfies $u(0, \cdot) \leq u_0$ (resp. $u(0, \cdot) \geq u_0$) on \mathbb{T}^d , u is called a viscosity subsolution (resp. viscosity supersolution) of (1.1)-(1.2). We often suppress the word ‘‘viscosity’’ unless confusion occurs.

2.3. Properties and equivalences of solutions.

Proposition 2.7 (Replacement of test functions). A function $u \in USC(\overline{Q_T})$ (resp. $LSC(\overline{Q_T})$) is a subsolution (resp. supersolution) of (1.1) if and only if, for any $a, b \in [0, T]$ with $a < b$ and an open ball B in \mathbb{R}^d , (2.3) holds whenever

- (i) $u - \phi$ attains a zero maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$ such that $(u - \phi)(\hat{t}, \hat{x}) = 0$, or
- (ii) $u - \phi$ attains a strict maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$, i.e.,

$$\max_{[a, b] \times \overline{B}} (u - \phi) = (u - \phi)(\hat{t}, \hat{x}) > (u - \phi)(t, x)$$

$$(\text{resp. } \min_{[a, b] \times \overline{B}} (u - \phi) = (u - \phi)(\hat{t}, \hat{x}) < (u - \phi)(t, x))$$

for all $(t, x) \in [a, b] \times \overline{B}$.

- (iii) $u - \phi$ attains a maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([a, b] \times \overline{B})$.

Proof. We only prove for a subsolution since a similar argument is applied for a supersolution. It is enough to prove ‘only if’ parts of both assertions since ‘if’ parts are obvious.

(i) Set $\psi := \phi + (u - \phi)(\hat{t}, \hat{x})$. Then $u - \psi$ attains a maximum at (\hat{t}, \hat{x}) over $[a, b] \times \overline{B}$ and $(u - \psi)(\hat{t}, \hat{x}) = 0$. Since u is a subsolution of (1.1),

$$J_{\hat{t}-a}[\psi](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) \leq 0.$$

It is easy to verify from the definition of $J_r[\phi]$ that $J_{\hat{t}-a}[\psi](\hat{t}, \hat{x}) = J_{\hat{t}-a}[\phi](\hat{t}, \hat{x})$. Clearly, $D\psi(\hat{t}, \hat{x}) = D\phi(\hat{t}, \hat{x})$, so that (2.3) holds.

(ii) For $j \geq 0$ we set $\phi_j(t, x) := \phi(t, x) + j^{-1}|t - \hat{t}|^2 + |x - \hat{x}|^2$ on $(0, T] \times \mathbb{R}^d$. Then $\phi_j \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$ and $u - \phi_j$ attains a maximum at (\hat{t}, \hat{x}) over $[a, b] \times \overline{B}$. Since u is a subsolution of (1.1),

$$J_{\hat{t}-a}[\phi_j](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\phi_j(\hat{t}, \hat{x})) \leq 0.$$

By the definition of ϕ_j we have

$$J_{\hat{t}-a}[\phi_j](\hat{t}, \hat{x}) = J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}) - \frac{\alpha}{j\Gamma(1-\alpha)} \int_0^{\hat{t}} \tau^2 \frac{d\tau}{\tau^{\alpha+1}}$$

The last integral in the right-hand side is clearly finite, so vanishes as $j \rightarrow \infty$. Since $D\phi_j(\hat{t}, \hat{x}) = D\phi(\hat{t}, \hat{x})$, we reach (2.3).

(iii) Choose $\delta > 0$ so that $2\delta < \hat{t} - a$ and $\overline{B_{2\delta}(\hat{x})} \subset B$. Let $\xi_1, \xi_2 : [0, T] \times \mathbb{R}^d \rightarrow [0, 1]$ be C^∞ functions such that $\xi_1 + \xi_2 = 1$ in $[0, T] \times \mathbb{R}^d$, $\xi_1 = 1$ on $[\hat{t} - \delta, \hat{t}] \times B_\delta(\hat{x})$ and $\xi_2 = 1$ on $([0, T] \times \mathbb{R}^d) \setminus ([\hat{t} - 2\delta, \hat{t}] \times B_{2\delta}(\hat{x}))$. Set $\psi := \xi_1\phi + \xi_2M + (u - \phi)(\hat{t}, \hat{x})$ on $[0, T] \times \mathbb{R}^d$, where $M := \max_{\overline{Q_T}} u + 1$. Then $\psi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$ and $u - \psi$ attains a zero maximum (\hat{t}, \hat{x}) over $[\hat{t} - \delta, \hat{t}] \times \overline{B} \subset [a, b] \times \overline{B}$. Since u is a subsolution of (1.1),

$$J_\delta[\psi](\hat{t}, \hat{x}) + K_\delta[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) \leq 0.$$

It is easy that $J_\delta[\psi](\hat{t}, \hat{x}) = J_\delta[\phi](\hat{t}, \hat{x})$ and $D\psi(\hat{t}, \hat{x}) = D\phi(\hat{t}, \hat{x})$. Moreover, since $u(\hat{t}, \hat{x}) - u(\hat{t} - \tau) \geq \phi(\hat{t}, \hat{x}) - \phi(\hat{t} - \tau, \hat{x})$ on $[0, \hat{t} - a]$,

$$K_\delta[u](\hat{t}, \hat{x}) \geq K_{\hat{t}-a}[u](\hat{t}, \hat{x}) + \frac{\alpha}{\Gamma(1-\alpha)} \int_\delta^{\hat{t}-a} (\phi(\hat{t}, \hat{x}) - \phi(\hat{t} - \tau, \hat{x})) \frac{d\tau}{\tau^{\alpha+1}}.$$

Thus we have $J_\delta[\psi](\hat{t}, \hat{x}) + K_\delta[u](\hat{t}, \hat{x}) \geq J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x})$, which is nothing but (2.3). \square

Remark 2.8. By the similar way in the proof of (i) it turns out that, if u is a subsolution (resp. supersolution) of (1.1), then $u - C$ (resp. $u + C$) is a subsolution (resp. supersolution) of (1.1) for any positive constant $C > 0$. This is valid even for sub/supersolutions of (1.1)-(1.2). Here a proof needs (A3).

Lemma 2.9 (Equivalence). *A function $u \in USC(\overline{Q_T})$ (resp. $LSC(\overline{Q_T})$) is a subsolution (resp. supersolution) of (1.1) if and only if $K_0[u](\hat{t}, \hat{x})$ exists and*

$$(2.4) \quad K_0[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) \leq 0 \quad (\text{resp. } \geq 0)$$

whenever $u - \phi$ attains a maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ over $[0, T] \times \mathbb{R}^d$ for $\phi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$.

Proof. We only prove for subsolutions since the similar argument is applied for supersolutions.

We first prove the ‘if’ part. To do so, let $a, b \in [0, T]$ with $a < b$ and an open ball B in \mathbb{R}^d fix arbitrarily. Assume that $u - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in \mathcal{C}^1((0, T] \times \mathbb{R}^d)$. Define $\psi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$ similarly as in the proof of Proposition 2.7 (iii). Then $u - \psi$ attains a zero maximum (\hat{t}, \hat{x}) over $[0, T] \times \mathbb{R}^d$. Thus $K_0[u](\hat{t}, \hat{x})$ exists and

$$(2.5) \quad K_0[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) \leq 0.$$

The relationship between u and ϕ implies that

$$u(\hat{t}, \hat{x}) - u(\hat{t} - \tau, \hat{x}) \geq \phi(\hat{t}, \hat{x}) - \phi(\hat{t} - \tau, \hat{x})$$

for all $[0, \hat{t} - a]$, which further yields $J_{\hat{t}-a}[u](\hat{t}, \hat{x}) \geq J_{\hat{t}-a}[\phi](\hat{t}, \hat{x})$. Since $D\psi(\hat{t}, \hat{x}) = D\phi(\hat{t}, \hat{x})$ and $K_0[u](\hat{t}, \hat{x}) = J_{\hat{t}-a}[u](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x})$, the assertion follows immediately.

To prove the ‘only if’ part we assume that $u - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ over $[0, T] \times \mathbb{R}^d$ for $\phi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$. Set $\psi := \phi + (u - \phi)(\hat{t}, \hat{x})$ on $(0, T] \times \mathbb{R}^d$. Let $r > 0$ be a parameter such that $\hat{t} - r > 0$. Then $u - \psi$ attains a zero maximum at (\hat{t}, \hat{x}) over $[\hat{t} - r, \hat{t}] \times \overline{B(\hat{x})}$ for all r , where $B(\hat{x})$ is an open ball centered at \hat{x} in \mathbb{R}^d . Since u is a subsolution of (1.1),

$$(2.6) \quad J_r[\psi](\hat{t}, \hat{x}) + K_r[u](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) \leq 0$$

for all r . From Proposition 2.4 and its proof we know that $J_r[\psi](\hat{t}, \hat{x})$ and $K_r[u^-](\hat{t}, \hat{x})$ exist for each r and moreover $\lim_{r \rightarrow 0} J_r[\psi](\hat{t}, \hat{x}) = 0$. Thus it is enough to show that $K_r[u^+]$ exists for each small r and $\lim_{r \rightarrow 0} K_r[u^\pm] = K_0[u^\pm]$ exist as a finite number. Indeed, if this is proved, it means that $K_0[u](\hat{t}, \hat{x})$ exists and (2.4) follows by passing to the limit $r \rightarrow 0$ in (2.6).

Define a function $v_r : [0, T] \rightarrow \mathbb{R}$ by

$$v_r(\tau) = \begin{cases} \psi(\hat{t}, \hat{x}) - \psi(\hat{t} - \tau, \hat{x}) & \text{for } \tau \in [0, \hat{t} - r] \times \mathbb{T}^d, \\ u(\hat{t}, \hat{x}) - u(\hat{t} - \tau, \hat{x}) & \text{for } \tau \in [\hat{t} - r, \hat{t}] \times \mathbb{T}^d. \end{cases}$$

We rewrite (2.6) as

$$(2.7) \quad I[v_r] \leq \frac{\Gamma(1 - \alpha)}{\alpha} \left(-H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) - \frac{u(\hat{t}, \hat{x}) - u(0, \hat{x})}{\hat{t}^\alpha \Gamma(1 - \alpha)} \right) =: C,$$

where

$$I[v_r](\hat{t}, \hat{x}) = \int_0^{\hat{t}} v_r(\tau) \frac{d\tau}{\tau^{\alpha+1}}.$$

From the relationship between u and ψ we see

$$(2.8) \quad \psi(\hat{t}, \hat{x}) - \psi(\hat{t} - \tau, \hat{x}) \leq u(\hat{t}, \hat{x}) - u(\hat{t} - \tau, \hat{x})$$

on $[0, \hat{t} - r]$. Then it suffices to prove that $I[v_r^+]$ exists for each small r and $\lim_{r \rightarrow 0} I[v_r^\pm] = I[v_0^\pm]$ exists as a finite number.

By the definition of v_r and (2.8), v_r^+ is monotone increasing with respect to r in the sense that $v_{r_1}^+ \leq v_{r_2}^+$ on $[0, \hat{t}]$ if $r_1 \geq r_2$. The monotone convergence theorem implies that $\lim_{r \rightarrow 0} I[v_r^+] = I[v_0^+]$. It is verified similarly as for v_r^+ that

v_r^- is monotone decreasing with respect to r in the sense that $v_{r_1}^- \leq v_{r_2}^-$ on $[0, \hat{t}]$ if $r_1 \leq r_2$. Thus we have from (2.7)

$$(2.9) \quad I[v_{r_1}^+](\hat{t}, \hat{x}) \leq I[v_{r_1}^-](\hat{t}, \hat{x}) + C \leq I[v_{r_2}^-](\hat{t}, \hat{x}) + C.$$

This implies that $I[v_r^+]$ exists for each small r and $I[v_0^+]$ exists (as a finite number) by passing to the limit $r_1 \rightarrow 0$. The monotone convergence theorem for $I[v_r^-]$ implies that $\lim_{r \rightarrow 0} I[v_r^-] = I[v_0^-]$. Therefore (2.9) ensures that $I[v_0^-]$ exists (as a finite number). The proof is now complete. \square

Proposition 2.10 (Consistency). *Assume that $u \in C^1(\overline{Q_T})$. Then u is a classical solution of (1.1)-(1.2) if and only if u is a viscosity solution of (1.1)-(1.2).*

Proof. Assume that u is a viscosity subsolution. We may take $\phi \equiv u$ so that $u - \phi$ attains a maximum at every point in Q_T . Since u is a viscosity subsolution of (1.1), Lemma 2.9 implies that

$$(2.10) \quad K_0[u](t, x) + H(t, x, u(t, x), Du(t, x)) \leq 0$$

for all $(t, x) \in Q_T$. Similarly, we have the reverse inequality of (2.10) from an inequality by viscosity supersolution. This shows that u is a classical solution since $K_0[u] = \partial_t^\alpha u$ by Proposition 2.3 (ii).

On the contrary we assume that u is a classical solution and that $u - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$, where $a, b \in (0, T]$ are constants and B is an open ball in \mathbb{R}^d . Since $(\partial_t^\alpha u)(\hat{t}, \hat{x}) = K_0[u](\hat{t}, \hat{x}) \geq J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u](\hat{t}, \hat{x})$, to combine the maximum principle in space implies that u is a viscosity subsolution. It is similar for viscosity supersolutions.

Since an initial condition is easily verified, we obtain the conclusion. \square

3. COMPARISON PRINCIPLE

Theorem 3.1 (Comparison principle). *Assume that (A1)-(A3). Let $u \in USC(\overline{Q_T})$ and $v \in LSC(\overline{Q_T})$ be a subsolution and a supersolution of (1.1), respectively. If $u(0, \cdot) \leq v(0, \cdot)$ on \mathbb{T}^d , then $u \leq v$ on $\overline{Q_T}$.*

We shall prepare one lemma for a proof of Theorem 3.1; see [8, Lemma 2], [17, Lemma 3.3] and [18, Lemma 1] for similar results for $\alpha = 1$. To do so we invoke a limit inferior/superior inequality of product of constant sequences that one of sequences is allowed to be negative. The statement looks fundamental and the proof is standard but we give for the reader's convenience.

Proposition 3.2. *Let $\{f_\varepsilon\}_{\varepsilon>0}$ and $\{g_\varepsilon\}_{\varepsilon>0}$ be constant sequences. Assume that g_ε is nonnegative, $\liminf_{\varepsilon \rightarrow 0} f_\varepsilon \geq f_0$ (resp. $\limsup_{\varepsilon \rightarrow 0} f_\varepsilon \leq f_0$) and $\lim_{\varepsilon \rightarrow 0} g_\varepsilon = g_0$ for some constants f_0 and g_0 . Then*

$$\liminf_{\varepsilon \rightarrow 0} f_\varepsilon g_\varepsilon \geq f_0 g_0, \quad (\text{resp. } \limsup_{\varepsilon \rightarrow 0} f_\varepsilon g_\varepsilon \leq f_0 g_0.)$$

Proof. It is enough to prove the case when $\liminf_{\varepsilon \rightarrow 0} f_\varepsilon \geq f_0$ since another case is proved by changing a sign of f_ε . Then for any $\delta > 0$ there exists $\varepsilon_\delta > 0$ such that $f_\varepsilon \geq f_0 - \delta$ for all $\varepsilon < \varepsilon_\delta$. It is fundamental that

$$\liminf_{\varepsilon \rightarrow 0} h_\varepsilon g_\varepsilon \geq \liminf_{\varepsilon \rightarrow 0} h_\varepsilon \liminf_{\varepsilon \rightarrow 0} g_\varepsilon$$

for nonnegative constants $h_\varepsilon, g_\varepsilon$. Applying this fact as $h_\varepsilon := f_\varepsilon - f_0 - \delta (\geq 0)$ we see

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} f_\varepsilon g_\varepsilon &\geq \liminf_{\varepsilon \rightarrow 0} (f_\varepsilon - f_0 - \delta) g_\varepsilon + \liminf_{\varepsilon \rightarrow 0} (f_0 + \delta) g_\varepsilon \\ &\geq -\delta g_0 + (f_0 + \delta) g_0 = f_0 g_0. \end{aligned}$$

□

Lemma 3.3. *Assume (A1). Let $u \in USC(\overline{Q_T})$ and $v \in LSC(\overline{Q_T})$ be a subsolution and a supersolution of (1.1), respectively. Assume that $(t, x, y) \mapsto u(t, x) - v(t, y) - \phi(t, x, y)$ attains a maximum at $(\hat{t}, \hat{x}, \hat{y}) \in (0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ over $[0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ for $\phi \in C^1([0, T] \times \mathbb{R}^d \times \mathbb{R}^d)$. Then*

$$K_0[u](\hat{t}, \hat{x}) - K_0[v](\hat{t}, \hat{y}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D_x \phi(\hat{t}, \hat{x}, \hat{y})) - H(\hat{t}, \hat{y}, v(\hat{t}, \hat{y}), -D_y \phi(\hat{t}, \hat{x}, \hat{y})) \leq 0.$$

Proof. We shall show that there exists a constant $C_r > 0$ such that $C_r \rightarrow 0$ as $r \rightarrow 0$ and

$$(3.1) \quad \begin{aligned} &-C_r + J_r[\phi](\hat{t}, \hat{x}, \hat{y}) + K_r[u](\hat{t}, \hat{x}) + K_r[v](\hat{t}, \hat{y}) \\ &+ H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D_x \phi(\hat{t}, \hat{x}, \hat{y})) - H(\hat{t}, \hat{y}, v(\hat{t}, \hat{y}), -D_y \phi(\hat{t}, \hat{x}, \hat{y})) \leq 0 \end{aligned}$$

for all $r \in (0, \hat{t})$. If this is clarified, passing to the limit $r \rightarrow 0$ in (3.1) yields the desired result by repeating the ‘only if’ in the proof of Lemma (2.9). Henceforth, let $r \in (0, \hat{t})$ fix arbitrarily.

For $\varepsilon > 0$ we consider a function $\Phi : [0, T] \times [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ defined by

$$\Phi(t, s, x, y) = u(t, x) - v(s, y) - \phi(t, x, y) - \frac{|t - s|^2}{2\varepsilon} - |t - \hat{t}|^2 - |x - \hat{x}|^2 - |y - \hat{y}|^2.$$

Since $\Phi \rightarrow -\infty$ as $|x|, |y| \rightarrow +\infty$ and Φ is bounded from above, it attains a maximum at a point $(t_\varepsilon, s_\varepsilon, x_\varepsilon, y_\varepsilon) \in [0, T] \times [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$. By following the standard argument of the theory of viscosity solutions we obtain

$$(3.2) \quad \begin{cases} (t_\varepsilon, s_\varepsilon, x_\varepsilon, y_\varepsilon) \rightarrow (\hat{t}, \hat{t}, \hat{x}, \hat{y}), \\ u(t_\varepsilon, x_\varepsilon) \rightarrow u(\hat{t}, \hat{x}) \text{ and } v(s_\varepsilon, y_\varepsilon) \rightarrow v(\hat{t}, \hat{y}) \end{cases}$$

as $\varepsilon \rightarrow 0$ by taking a subsequence if necessary; see [4, Theorem II.3.1] and [7, Lemma 3.1] for detail. Note that $t_\varepsilon > 0$ for sufficiently small ε since $\hat{t} > 0$.

For such a small parameter ε , $(t, x) \mapsto \Phi(t, s_\varepsilon, x, y_\varepsilon)$ attains a maximum at $(t_\varepsilon, x_\varepsilon) \in (0, T] \times \mathbb{R}^d$ over $[0, T] \times \mathbb{R}^d$ and $(s, y) \mapsto -\Phi(t_\varepsilon, s, x_\varepsilon, y)$ attains a minimum at $(s_\varepsilon, y_\varepsilon) \in (0, T] \times \mathbb{R}^d$ over $[0, T] \times \mathbb{R}^d$. Since u and v are respectively a subsolution and a supersolution of (1.1), Lemma 2.9 implies that $K_0[u](t_\varepsilon, x_\varepsilon), K_0[v](s_\varepsilon, y_\varepsilon)$ exist for each ε and

$$(3.3) \quad K_0[u](t_\varepsilon, x_\varepsilon) + H(t_\varepsilon, x_\varepsilon, u(t_\varepsilon, x_\varepsilon), D_x \phi(t_\varepsilon, x_\varepsilon, y_\varepsilon)) + 2(x_\varepsilon - \hat{x}) \leq 0,$$

$$(3.4) \quad K_0[v](s_\varepsilon, y_\varepsilon) + H(s_\varepsilon, y_\varepsilon, v(s_\varepsilon, y_\varepsilon), -D_y \phi(s_\varepsilon, x_\varepsilon, y_\varepsilon)) - 2(y_\varepsilon - \hat{y}) \geq 0.$$

Thus, by subtracting (3.4) from (3.3), we see

$$(3.5) \quad \begin{aligned} &K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](s_\varepsilon, y_\varepsilon) \\ &+ H(t_\varepsilon, x_\varepsilon, u(t_\varepsilon, x_\varepsilon), D_x \phi(t_\varepsilon, x_\varepsilon, y_\varepsilon)) + 2(x_\varepsilon - \hat{x}) \\ &- H(s_\varepsilon, y_\varepsilon, v(s_\varepsilon, y_\varepsilon), -D_y \phi(s_\varepsilon, x_\varepsilon, y_\varepsilon)) - 2(y_\varepsilon - \hat{y}) \leq 0 \end{aligned}$$

for each ε .

We shall pass to the limit $\varepsilon \rightarrow 0$ in (3.5). For Hamiltonians it is easily seen thanks to (A1), (3.2) and the smoothness of ϕ that

$$\begin{aligned} & H(t_\varepsilon, x_\varepsilon, u(t_\varepsilon, x_\varepsilon), D_x \phi(t_\varepsilon, x_\varepsilon, y_\varepsilon) + 2(x_\varepsilon - \hat{x})) \\ & - H(s_\varepsilon, y_\varepsilon, v(s_\varepsilon, y_\varepsilon), -D_y \phi(s_\varepsilon, x_\varepsilon, y_\varepsilon) - 2(y_\varepsilon - \hat{y})) \\ & \rightarrow H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), D_x \phi(\hat{t}, \hat{x}, \hat{y})) - H(\hat{t}, \hat{y}, v(\hat{t}, \hat{y}), -D_y \phi(\hat{t}, \hat{x}, \hat{y})) \end{aligned}$$

as $\varepsilon \rightarrow 0$. Let us focus on $K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](s_\varepsilon, y_\varepsilon)$. Assume hereafter that ε is so small that $r < \min\{t_\varepsilon, s_\varepsilon\}$ for all ε , which is possible since $(t_\varepsilon, s_\varepsilon) \rightarrow (\hat{t}, \hat{t})$ as $\varepsilon \rightarrow 0$ (see (3.2)). Set

$$\begin{aligned} I_{1,\varepsilon} &:= \frac{u(t_\varepsilon, x_\varepsilon) - u(0, x_\varepsilon)}{t_\varepsilon^\alpha \Gamma(1-\alpha)} - \frac{v(s_\varepsilon, y_\varepsilon) - v(0, y_\varepsilon)}{s_\varepsilon^\alpha \Gamma(1-\alpha)}, \\ I_{2,\varepsilon} &:= \int_0^r (u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \tau, x_\varepsilon)) \frac{d\tau}{\tau^{\alpha+1}} - \int_0^r (v(s_\varepsilon, y_\varepsilon) - v(s_\varepsilon - \tau, y_\varepsilon)) \frac{d\tau}{\tau^{1+\alpha}}, \end{aligned}$$

and

$$I_{3,\varepsilon} := \int_r^{t_\varepsilon} (u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \tau, x_\varepsilon)) \frac{d\tau}{\tau^{\alpha+1}} - \int_r^{s_\varepsilon} (v(s_\varepsilon, y_\varepsilon) - v(s_\varepsilon - \tau, y_\varepsilon)) \frac{d\tau}{\tau^{1+\alpha}}$$

so that $K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](s_\varepsilon, y_\varepsilon) = I_{3,\varepsilon} + \alpha(I_{1,\varepsilon} + I_{2,\varepsilon})/\Gamma(1-\alpha)$.

First, for $I_{1,\varepsilon}$, Proposition 3.2 with $f_\varepsilon := u(t_\varepsilon, x_\varepsilon) - u(0, x_\varepsilon) - v(s_\varepsilon, y_\varepsilon) + v(0, y_\varepsilon)$ and $g_\varepsilon := (t_\varepsilon^\alpha \Gamma(1-\alpha))^{-1}$ implies that

$$(3.6) \quad \liminf_{\varepsilon \rightarrow 0} I_{1,\varepsilon} \geq \frac{u(\hat{t}, \hat{x}) - u(0, \hat{x})}{\hat{t}^\alpha \Gamma(1-\alpha)} - \frac{v(\hat{t}, \hat{y}) - v(0, \hat{y})}{\hat{t}^\alpha \Gamma(1-\alpha)}.$$

Next, since

$$\begin{aligned} & u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \tau, x_\varepsilon) - (v(s_\varepsilon, y_\varepsilon) - v(s_\varepsilon - \tau, y_\varepsilon)) \\ & \geq \phi(t_\varepsilon, x_\varepsilon, y_\varepsilon) - \phi(t_\varepsilon - \tau, x_\varepsilon, y_\varepsilon) + |t_\varepsilon - \hat{t}|^2 - |t_\varepsilon - \tau - \hat{t}|^2 \end{aligned}$$

for all $\tau \in [0, r]$ by the inequality $\Phi(t_\varepsilon, s_\varepsilon, x_\varepsilon, y_\varepsilon) \geq \Phi(t_\varepsilon - \tau, s_\varepsilon - \tau, x_\varepsilon, y_\varepsilon)$, we see

$$\frac{\alpha}{\Gamma(1-\alpha)} I_{2,\varepsilon} \geq J_r[\phi](t_\varepsilon, x_\varepsilon, y_\varepsilon) + J_r[|t_\varepsilon - \hat{t} - \cdot|^2](t_\varepsilon).$$

Proposition 2.4 (iv) ensures that $\lim_{\varepsilon \rightarrow 0} J_r[\phi](t_\varepsilon, x_\varepsilon, y_\varepsilon) = J_r[\phi](\hat{t}, \hat{x}, \hat{y})$. Besides, $J_r[|t_\varepsilon - \hat{t} - \cdot|^2](t_\varepsilon)$ can be calculated precisely as

$$\begin{aligned} J_r[|t_\varepsilon - \hat{t} - \cdot|^2](t_\varepsilon) &= \frac{\alpha}{\Gamma(1-\alpha)} \int_0^r (|t_\varepsilon - \hat{t}|^2 - |t_\varepsilon - \hat{t} - \tau|^2) \frac{d\tau}{\tau^{1+\alpha}} \\ &= \frac{\alpha}{\Gamma(1-\alpha)} \int_0^r (2(t_\varepsilon - \hat{t})\tau - \tau^2) \frac{d\tau}{\tau^{1+\alpha}} \\ &= \frac{\alpha}{\Gamma(1-\alpha)} \left(\frac{2(t_\varepsilon - \hat{t})r^{1-\alpha}}{1-\alpha} - \frac{r^{2-\alpha}}{2-\alpha} \right). \end{aligned}$$

Hence

$$\lim_{\varepsilon \rightarrow 0} J_r[|t_\varepsilon - \hat{t} - \cdot|^2](t_\varepsilon) = -\frac{\alpha r^{2-\alpha}}{(2-\alpha)\Gamma(1-\alpha)} =: -C_r.$$

Note that $C_r \rightarrow 0$ as $r \rightarrow 0$. Therefore we know for $I_{2,\varepsilon}$ that

$$(3.7) \quad \liminf_{\varepsilon \rightarrow 0} \frac{\alpha}{\Gamma(1-\alpha)} I_{2,\varepsilon} \geq -C_r + J_r[\phi](\hat{t}, \hat{x}, \hat{y}).$$

Finally, for $I_{3,\varepsilon}$, we first see an existence of constants $C_1, C_2 > 0$ independent of ε such that

$$(3.8) \quad \begin{aligned} (u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \tau, x_\varepsilon))\mathbb{1}_{(r, t_\varepsilon)}(\tau) &\geq -C_1\mathbb{1}_{(r, T)}(\tau), \\ (v(s_\varepsilon, y_\varepsilon) - v(s_\varepsilon - \tau, y_\varepsilon))\mathbb{1}_{(r, s_\varepsilon)}(\tau) &\leq C_2\mathbb{1}_{(r, T)}(\tau) \end{aligned}$$

on $[0, T]$. Indeed, since $\lim_{\varepsilon \rightarrow 0} u(t_\varepsilon, x_\varepsilon) = u(\hat{t}, \hat{x})$, there is a constant $C > 0$ independent of ε

$$\begin{aligned} (u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \tau, x_\varepsilon))\mathbb{1}_{(r, t_\varepsilon)}(\tau) &\geq (u(\hat{t}, \hat{x}) - C - \frac{\max u}{Q_T})\mathbb{1}_{(r, t_\varepsilon)}(\tau) \\ &\geq -|u(\hat{t}, \hat{x}) - C - \frac{\max u}{Q_T}|\mathbb{1}_{(r, T)}. \end{aligned}$$

This shows the above one of (3.8) and another is proved similarly. Note that both right-hand sides of (3.8) multiplied by $\tau^{-\alpha-1}$ is integrable on $[0, T]$. Proposition 3.2 implies that

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} (u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \cdot, x_\varepsilon))\mathbb{1}_{(r, t_\varepsilon)}(\cdot) &\geq (u(\hat{t}, \hat{x}) - u(\hat{t} - \cdot, \hat{x}))\mathbb{1}_{(r, \hat{t})}(\cdot), \\ \limsup_{\varepsilon \rightarrow 0} (v(s_\varepsilon, y_\varepsilon) - v(s_\varepsilon - \cdot, y_\varepsilon))\mathbb{1}_{(r, s_\varepsilon)}(\cdot) &\leq (v(\hat{t}, \hat{x}) - v(\hat{t} - \cdot, \hat{x}))\mathbb{1}_{(r, \hat{t})}(\cdot) \end{aligned}$$

for each $\tau \in (0, T)$. Thus Fatou's lemma yields

$$(3.9) \quad \liminf_{\varepsilon \rightarrow 0} I_{3,\varepsilon} \geq \int_r^{\hat{t}} (u(\hat{t}, \hat{x}) - u(\hat{t} - \tau, \hat{x})) \frac{d\tau}{\tau^{1+\alpha}} - \int_r^{\hat{t}} (v(\hat{t}, \hat{x}) - v(\hat{t} - \tau, \hat{x})) \frac{d\tau}{\tau^{1+\alpha}}.$$

Summing up (3.6), (3.7) and (3.9) we reach

$$\liminf_{\varepsilon \rightarrow 0} (K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](s_\varepsilon, y_\varepsilon)) \geq -C_r + J_r[\phi](\hat{t}, \hat{x}, \hat{y}) + K_r[u](\hat{t}, \hat{x}) - K_r[v](\hat{t}, \hat{x}).$$

Consequently, taking the limit inferior to both sides of (3.5) yields the desired inequality (3.1). \square

Proof of Theorem 3.1. Suppose that the conclusion were false: $\max_{\overline{Q_T}}(u - v) =: \theta > 0$. For $\varepsilon > 0$ we consider a function $\Phi : [0, T] \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ defined by

$$\Phi(t, x, y) := u(t, x) - v(t, y) - \frac{|x - y|^2}{2\varepsilon}.$$

Let $(t_\varepsilon, x_\varepsilon, y_\varepsilon) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ be a maximum point of Φ . Then there is a point $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ such that

$$(3.10) \quad \begin{cases} (t_\varepsilon, x_\varepsilon, y_\varepsilon) \rightarrow (\hat{t}, \hat{x}, \hat{x}), \\ |x_\varepsilon - y_\varepsilon|^2/\varepsilon \rightarrow 0, \\ u(t_\varepsilon, x_\varepsilon) \rightarrow u(\hat{t}, \hat{x}) \text{ and } v(s_\varepsilon, y_\varepsilon) \rightarrow v(\hat{t}, \hat{x}). \end{cases}$$

as $\varepsilon \rightarrow 0$ by taking a subsequence if necessary; see, e.g., [4, Theorem II.3.1]. The above permits to use Lemma 3.3 and we know that $K_0[u](t_\varepsilon, x_\varepsilon), K_0[v](t_\varepsilon, y_\varepsilon)$ exists for each ε and

$$(3.11) \quad K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](t_\varepsilon, y_\varepsilon) + H(t_\varepsilon, x_\varepsilon, u(t_\varepsilon, x_\varepsilon), p_\varepsilon) - H(t_\varepsilon, y_\varepsilon, v(t_\varepsilon, y_\varepsilon), p_\varepsilon) \leq 0.$$

Here $p_\varepsilon = (x_\varepsilon - y_\varepsilon)/\varepsilon$.

Since $u(t_\varepsilon, x_\varepsilon) - u(t_\varepsilon - \cdot, x_\varepsilon) - v(t_\varepsilon, y_\varepsilon) + v(t_\varepsilon - \cdot, y_\varepsilon) \geq 0$ on $[0, t_\varepsilon]$ by the inequality $\Phi(t_\varepsilon, x_\varepsilon, y_\varepsilon) \geq \Phi(t_\varepsilon - \tau, x_\varepsilon, y_\varepsilon)$, the term of integration in $K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](t_\varepsilon, y_\varepsilon)$ is estimated from below by zero, that is,

$$K_0[u](t_\varepsilon, x_\varepsilon) - K_0[v](t_\varepsilon, y_\varepsilon) \geq \frac{u(t_\varepsilon, x_\varepsilon) - v(t_\varepsilon, y_\varepsilon) - u(0, x_\varepsilon) + v(0, y_\varepsilon)}{t_\varepsilon^\alpha \Gamma(1 - \alpha)}$$

Since $u(t_\varepsilon, x_\varepsilon) > v(t_\varepsilon, y_\varepsilon)$ by the inequality $\Phi(t_\varepsilon, x_\varepsilon, y_\varepsilon) \geq \theta > 0$, Hamiltonians in (3.11) are estimated as

$$\begin{aligned} & H(t_\varepsilon, x_\varepsilon, u(t_\varepsilon, x_\varepsilon), p_\varepsilon) - H(t_\varepsilon, y_\varepsilon, v(t_\varepsilon, y_\varepsilon), p_\varepsilon) \\ & \geq H(t_\varepsilon, x_\varepsilon, v(t_\varepsilon, y_\varepsilon), p_\varepsilon) - H(t_\varepsilon, y_\varepsilon, v(t_\varepsilon, y_\varepsilon), p_\varepsilon) \geq -\omega(|x_\varepsilon - y_\varepsilon|(1 + |p_\varepsilon|)) \end{aligned}$$

by (A2) and (A3). From these, (3.11) is led to

$$\frac{u(t_\varepsilon, x_\varepsilon) - v(t_\varepsilon, y_\varepsilon) - u(0, x_\varepsilon) + v(0, y_\varepsilon)}{t_\varepsilon^\alpha \Gamma(1 - \alpha)} \leq \omega(|x_\varepsilon - y_\varepsilon|(1 + |p_\varepsilon|)).$$

Taking the limit inferior $\varepsilon \rightarrow 0$ implies that

$$\frac{\theta - u(0, \hat{x}) + v(0, \hat{x})}{\hat{t}^\alpha \Gamma(1 - \alpha)} \leq 0$$

by Proposition 3.2. Since $u(0, \cdot) \leq v(0, \cdot)$ on \mathbb{T}^d and $\theta > 0$, this is a contradiction. \square

Corollary 3.4 (Uniqueness). *Assume (A1)-(A4). Let $u \in C(\overline{Q_T})$ and $v \in C(\overline{Q_T})$ be solutions of (1.1). Then*

$$(3.12) \quad \max_{(t,x) \in \overline{Q_T}} |u(t, x) - v(t, x)| \leq \max_{x \in \mathbb{T}^d} |u(0, x) - v(0, x)|.$$

Moreover, if u and v are solutions of (1.1)-(1.2), then $u \equiv v$ on $\overline{Q_T}$.

Proof. It suffices to prove (3.12). Set $C := \max_{x \in \mathbb{T}^d} |u(0, x) - v(0, x)|$. Then $v - C$ and $v + C$ are a subsolution and a supersolution of (1.1), respectively; see Remark 2.8. Moreover

$$v(0, \cdot) - C \leq u(0, \cdot) \leq v(0, \cdot) + C \quad \text{on } \mathbb{T}^d$$

by the definition of C . Thus, from Theorem 3.1, we have $|u - v| \leq C$ on $\overline{Q_T}$. The proof is complete by taking the maximum over $\overline{Q_T}$ to both sides. \square

For the reader's convenience we give a statement of the comparison principle for a general bounded domain Ω without a proof.

Theorem 3.5. *Let Ω be a bounded domain in \mathbb{R}^d . Let $u \in USC([0, T] \times \overline{\Omega}; \mathbb{R})$ and $v \in LSC([0, T] \times \overline{\Omega}; \mathbb{R})$ be a subsolution and a supersolution of (1.1) on $(0, T] \times \overline{\Omega}$, respectively. If $u \leq v$ on $(\{0\} \times \overline{\Omega}) \cup ([0, T] \times \partial\Omega)$, then $u \leq v$ on $[0, T] \times \overline{\Omega}$.*

4. EXISTENCE RESULT

Let denote by S^- and S^+ a set of upper semicontinuous subsolutions and lower semicontinuous supersolutions of (1.1), respectively. Note that $S^\pm \neq \emptyset$ as will be observed in Corollary 4.3 later.

Lemma 4.1 (Closedness under supremum/infimum operator). *Assume (A1). Let X be a nonempty subset of S^- (resp. S^+). Define*

$$u(t, x) := \sup\{v(t, x) \mid v \in X\} \quad (\text{resp. } \inf\{v(t, x) \mid v \in X\})$$

for $(t, x) \in \overline{Q_T}$. Assume that $u^* < +\infty$ (resp. $u_* > -\infty$) on $\overline{Q_T}$. Then u^* (resp. u_*) is a subsolution (resp. supersolution) of (1.1).

Proof. We only prove for a subsolution since the argument for a supersolution is similar. Fix $[a, b] \times B \subset (0, T] \times \mathbb{R}^d$ arbitrarily, where $a < b$ and B is an open in \mathbb{R}^d . Assume that $u^* - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (a, b] \times B$ over $[a, b] \times \overline{B}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$. Then we must show that

$$(4.1) \quad J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}) + K_{\hat{t}-a}[u^*](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u^*(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) \leq 0.$$

By Proposition 2.7 we may assume that (\hat{t}, \hat{x}) is a strict maximum point of $u^* - \phi$ such that $(u^* - \phi)(\hat{t}, \hat{x}) = 0$.

By arguing similarly as for $\alpha = 1$ we find sequences $\{(t_j, x_j)\}_{j \geq 0}$ and $\{v_j\}_{j \geq 0} \subset X$ such that, for each $j \geq 0$, $v_j - \phi$ attains a maximum at $(t_j, x_j) \in (a, b] \times B$ over $[a, b] \times \overline{B}$ and $(t_j, x_j, v_j(t_j, x_j)) \rightarrow (\hat{t}, \hat{x}, u^*(\hat{t}, \hat{x}))$ as $j \rightarrow \infty$. Indeed it is enough to translate slightly the proof of [12, Lemma 2.4.1] to the current situation. This is not difficult, so the detail is safely omitted. Since v_j is a subsolution of (1.1),

$$(4.2) \quad J_{t_j-a}[\phi](t_j, x_j) + K_{t_j-a}[v_j](t_j, x_j) + H(t_j, x_j, v_j(t_j, x_j), D\phi(t_j, x_j)) \leq 0$$

for each $j \geq 0$.

We shall pass to the limit $j \rightarrow \infty$ in (4.2). The continuity of Hamiltonian (A1) ensures that

$$\lim_{j \rightarrow \infty} H(t_j, x_j, v_j(t_j, x_j), D\phi(t_j, x_j)) = H(\hat{t}, \hat{x}, u^*(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})).$$

Proposition 2.4 implies that

$$\lim_{j \rightarrow \infty} J_{t_j-a}[\phi](t_j, x_j) = J_{\hat{t}-a}[\phi](\hat{t}, \hat{x}).$$

Henceforth, let us focus on $K_{t_j-a}[v_j](t_j, x_j)$. Since $v_j \leq u \leq u^*$ on $\overline{Q_T}$ by the definition of u and u^* , Proposition 3.2 implies that

$$(4.3) \quad \begin{aligned} \liminf_{j \rightarrow \infty} \frac{v_j(t_j, x_j) - v_j(0, x_j)}{t_j^\alpha \Gamma(1 - \alpha)} &\geq \liminf_{j \rightarrow \infty} \frac{v_j(t_j, x_j) - u^*(0, x_j)}{t_j^\alpha \Gamma(1 - \alpha)} \\ &\geq \frac{u^*(\hat{t}, \hat{x}) - u^*(0, \hat{x})}{\hat{t}^\alpha \Gamma(1 - \alpha)}. \end{aligned}$$

To handle the term of integration we first see the existence of a constant $C_2 > 0$ independent of j such that

$$(v_j(t_j, x_j) - v_j(t_j - \cdot, x_j))\mathbf{1}_{[t_j-a, t_j]}(\cdot) \geq -C_2\mathbf{1}_{[r, T]}(\cdot)$$

on $[0, T]$ for sufficiently large j , where $r := \min_{j \geq 0}(t_j - a) > 0$. Indeed, since $\sup_{j \geq 0} v_j \leq u \leq u^*$ and $u^* < +\infty$ on $\overline{Q_T}$, there is a constant $C_3 > 0$ such that $\sup_{j \geq 0} v_j \leq C_3$ on $\overline{Q_T}$. Since $v_j(t_j, x_j) \rightarrow u^*(\hat{t}, \hat{x})$ as $j \rightarrow \infty$, for a constant $C_4 > 0$ (independent of j), $v_j(t_j, x_j) \geq u^*(\hat{t}, \hat{x}) - C_4$ for large j . Thus, if we set $C_2 := |u^*(\hat{t}, \hat{x}) - C_4 - C_3|$, then

$$\begin{aligned} (v_j(t_j, x_j) - v_j(t_j - \cdot, x_j))\mathbf{1}_{[t_j-a, t_j]}(\cdot) &\geq (u^*(\hat{t}, \hat{x}) - C_4 - C_3)\mathbf{1}_{[t_j-a, t_j]}(\cdot) \\ &\geq -C_2\mathbf{1}_{[r, T]}(\cdot) \end{aligned}$$

on $[0, T]$ for sufficiently large j , which is the desired fact. Note that $-C_2 \mathbf{1}_{[r, T]}(\tau)/\tau^{\alpha+1}$ is integrable on $[0, T]$. Proposition 3.2 also implies that

$$\begin{aligned} & \liminf_{j \rightarrow \infty} (v_j(t_j, x_j) - v_j(t_j - \cdot, x_j)) \mathbf{1}_{[t_j - a, t_j]}(\cdot) \\ & \geq \liminf_{j \rightarrow \infty} (v_j(t_j, x_j) - u^*(t_j - \cdot, x_j)) \mathbf{1}_{[t_j - a, t_j]}(\cdot) \\ & \geq (u^*(\hat{t}, \hat{x}) - u^*(\hat{t} - \cdot, \hat{x})) \mathbf{1}_{[\hat{t} - a, \hat{t}]}(\cdot). \end{aligned}$$

Therefore Fatou's lemma can be applied and consequently

$$\liminf_{j \rightarrow \infty} K_{t_j - a}[v_j](t_j, x_j) \geq K_{\hat{t} - a}[u^*](\hat{t}, \hat{x})$$

by combining with (4.3).

Taking the limit inferior $j \rightarrow \infty$ to both sides in (4.2) yields (4.1). \square

Theorem 4.2 (Existence). *Assume (A1). Let $u^- \in USC(\overline{Q_T})$ and $u^+ \in LSC(\overline{Q_T})$ be a supersolution and a subsolution of (1.1) such that $(u^-)_* > -\infty$ and $(u^+)^* < +\infty$ on $\overline{Q_T}$. Suppose that $u^- \leq u^+$ in $\overline{Q_T}$. Then there exists a solution u of (1.1) that satisfies $u^- \leq u \leq u^+$ in $\overline{Q_T}$.*

Proof. Define

$$(4.4) \quad u(t, x) := \sup\{v(t, x) \mid v \in X\}$$

for $(t, x) \in \overline{Q_T}$, where

$$X := \{v \in S^- \mid v \leq u^+ \text{ on } \overline{Q_T}\}.$$

Note that $X \neq \emptyset$ since $u^- \in X$. Also, since $u^- \leq u \leq u^+$ on $\overline{Q_T}$ by the definition of u , $-\infty < (u^-)_* \leq u_* \leq u^* \leq (u^+)^* < +\infty$ on $\overline{Q_T}$. Our goal in this proof is to show that u defined by (4.4) is actually a solution of (1.1). Since we know that u^* is a subsolution of (1.1) from Lemma 4.1, so it suffices to show that u_* is a supersolution of (1.1).

Suppose by contradiction that u_* were not a supersolution of (1.1). Then there would be a function $\phi \in C^1([0, T] \times \mathbb{R}^d)$, a point $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ and a constant $\theta > 0$ such that $u_* - \phi$ attains a minimum at $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ over $(0, T] \times \mathbb{R}^d$ and

$$K_0[u_*](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u_*(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) < -2\theta.$$

Notice that $K_0[u_*](\hat{t}, \hat{x})$ may be $-\infty$, while $K_0[u_*](\hat{t}, \hat{x})$ makes sense and is bounded from above by Proposition 2.4.

Let $\rho > 0$ be a small parameter so that $\rho < \hat{t}$ and $\overline{B_{2\rho}(\hat{x})} \subset (\hat{x} - \frac{1}{2}, \hat{x} + \frac{1}{2})^d$. Define functions $w : (0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $U : [0, T] \times (\hat{x} - \frac{1}{2}, \hat{x} + \frac{1}{2})^d \rightarrow \mathbb{R}$ by

$$w(s, y) := \phi(s, y) + \frac{\rho^2}{2} - |s - \hat{t}|^2 - |y - \hat{x}|^2$$

and

$$U = \begin{cases} \max\{u^*, w\} & \text{in } ((\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})) \cap ([0, T] \times (\hat{x} - \frac{1}{2}, \hat{x} + \frac{1}{2})^d), \\ u^* & \text{in } ([0, T] \times (\hat{x} - \frac{1}{2}, \hat{x} + \frac{1}{2})^d) \setminus ((\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})), \end{cases}$$

respectively. We regard $B_{2\rho}(\hat{x})$ and, for each $s \in [0, T]$, $U(s, \cdot)$ as a open ball in \mathbb{T}^d and a function on \mathbb{T}^d by extending it periodically, respectively. We shall show that $U \in X$ and that there exists a point $(s, y) \in \overline{Q_T}$ such that $U(s, y) > u(s, y)$. Once these were proved, we would obtain a contradiction due to the maximality of u .

Set

$$\Omega := \left\{ (s, y) \in (\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x}) \mid |s - \hat{t}|^2 + |y - \hat{x}|^2 \leq \frac{\rho^2}{2} \right\} \subset \overline{Q_T}.$$

Then $\overline{\Omega} \subset (\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})$ and

$$(4.5) \quad u^*(s, y) \geq u_*(s, y) \geq \phi(s, y) = w(s, y) - \frac{\rho^2}{2} + |s - \hat{t}|^2 + |y - \hat{x}|^2 > w(s, y)$$

for all $(s, y) \in ((\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})) \setminus \Omega$. Thus U is upper semicontinuous on $\overline{Q_T}$ by its definition.

Assume that $U - \psi$ attains a maximum at $(\hat{s}, \hat{y}) \in (0, T] \times \mathbb{R}^d$ over $(0, T] \times \mathbb{R}^d$ for $\psi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$. We may assume that $(U - \psi)(\hat{s}, \hat{y}) = 0$.

Case 1: Suppose that $U(\hat{s}, \hat{y}) = u^*(\hat{s}, \hat{y})$. Then, since $U \geq u^*$ on $\overline{Q_T}$, it turns out that $u^* - \psi$ attains a maximum at (\hat{s}, \hat{y}) over $(0, T] \times \mathbb{R}^d$ and that

$$(4.6) \quad U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y}) \leq u^*(\hat{s}, \hat{y}) - u^*(\hat{s} - \tau, \hat{y})$$

for all $\tau \in [0, \hat{s}]$. Recall that u^* is a subsolution of (1.1), so that

$$K_0[u^*](\hat{s}, \hat{y}) + H(\hat{s}, \hat{y}, u(\hat{s}, \hat{y}), D\psi(\hat{s}, \hat{y})) \leq 0.$$

Proposition 2.4 (iii) with (4.6) ensures that $K_0[U](\hat{s}, \hat{y})$ exists and simultaneously $K_0[U](\hat{s}, \hat{y}) \leq K_0[u^*](\hat{s}, \hat{y})$. This implies that U is a subsolution of (1.1).

Case 2: Suppose that $U(\hat{s}, \hat{y}) = w(\hat{s}, \hat{y}) > u(\hat{s}, \hat{y})$. Then, from (4.5), we see $(\hat{s}, \hat{y}) \in \Omega$, which yields $\lim_{\rho \rightarrow 0}(\hat{s}, \hat{y}) = (\hat{t}, \hat{x})$. By employing the idea in [16, Theorem 3] for example, we shall show that

$$(4.7) \quad \limsup_{\rho \rightarrow 0} K_0[U](\hat{s}, \hat{y}) \leq K_0[u_*](\hat{t}, \hat{x}).$$

Since $U \geq u^* \geq u_*$ on $\overline{Q_T}$ the non-integration term is estimated as

$$\frac{U(\hat{s}, \hat{y}) - U(0, \hat{y})}{\hat{s}^\alpha \Gamma(1 - \alpha)} \leq \frac{w(\hat{s}, \hat{y}) - u_*(0, \hat{y})}{\hat{s}^\alpha \Gamma(1 - \alpha)}.$$

Recalling that $\lim_{\rho \rightarrow 0} w(\hat{s}, \hat{y}) = \phi(\hat{t}, \hat{x}) = u_*(\hat{t}, \hat{x})$ we see

$$\limsup_{\rho \rightarrow 0} \frac{U(\hat{s}, \hat{y}) - U(0, \hat{y})}{\hat{s}^\alpha \Gamma(1 - \alpha)} \leq \frac{u_*(\hat{t}, \hat{x}) - u_*(0, \hat{x})}{\hat{t}^\alpha \Gamma(1 - \alpha)}.$$

To handle the term of integration let us divide the term of integration in $K_0[U](\hat{s}, \hat{y})$ multiplied by $\Gamma(1 - \alpha)/\alpha$ into two integrations as follows:

$$I_{1,\rho}[U] := \int_0^{\rho^2} (U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y})) \frac{d\tau}{\tau^{\alpha+1}}$$

and

$$I_{2,\rho}[U] := \int_{\rho^2}^{\hat{s}} (U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y})) \frac{d\tau}{\tau^{\alpha+1}}.$$

By definitions of U and w we have

$$\begin{aligned} U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y}) &\leq w(\hat{s}, \hat{y}) - w(\hat{s} - \tau, \hat{y}) \\ &= \phi(\hat{s}, \hat{y}) - \phi(\hat{s} - \tau, \hat{y}) + \tau^2 - 2(\hat{s} - \hat{y})\tau \end{aligned}$$

for all $\tau \in [0, \rho^2]$. Hence $I_{1,\rho}[U] \leq I_{1,\rho}[\phi] + C_\rho$ with a constant C_ρ such that $\lim_{\rho \rightarrow 0} C_\rho = 0$. By Proposition 2.4 (iv), we see that $\lim_{\rho \rightarrow 0} I_{1,\rho}[\phi] = 0$, so that

$$\limsup_{\rho \rightarrow 0} I_{1,\rho}[U] \leq 0.$$

Since $U \geq u_*$ on $\overline{Q_T}$,

$$(4.8) \quad \begin{aligned} U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y}) &\leq w(\hat{s}, \hat{y}) - u_*(\hat{s} - \tau, \hat{y}) \\ &\leq \phi(\hat{s}, \hat{y}) - u_*(\hat{s} - \tau, \hat{y}) + \frac{\rho^2}{2} \end{aligned}$$

on $[\rho^2, \hat{s}]$. Moreover, the relationship between u_* and ϕ yields

$$\phi(\hat{s}, \hat{y}) - u_*(\hat{s} - \tau, \hat{y}) + \frac{\rho^2}{2} \leq \phi(\hat{s}, \hat{y}) - \phi(\hat{s} - \tau, \hat{y}) + \frac{\rho^2}{2}.$$

Since $\phi(\cdot, x)$ is continuous on $[0, T]$, we are able to find a large constant $C_1 > 0$ such that

$$\phi(\hat{s}, \hat{y}) - \phi(\hat{s} - \tau, \hat{y}) \leq C_1 \tau$$

for all $\tau \in [\rho^2, \hat{s}]$. In addition we may assume that C_1 does not depend on ρ . Notice that there exists a constant $C_2 > 0$ such that $C_1 \tau^2 + \rho^2/2 \leq C_2 \tau$ for all $\tau \in [\rho^2, \hat{s}]$. Consequently (4.8) is lead to

$$U(\hat{s}, \hat{y}) - U(\hat{s} - \tau, \hat{y}) \leq C_2 \tau$$

on $[\rho^2, \hat{s}]$. The right-hand side with $\tau^{-\alpha-1}$ is integrable on $[0, T]$, so that Fatou's lemma yields

$$\limsup_{\rho \rightarrow 0} I_{2,\rho}[U](\hat{s}, \hat{y}) \leq I_{2,0}[u_*](\hat{t}, \hat{x}).$$

The above ensures (4.7) and thus we see

$$K_0[U](\hat{s}, \hat{y}) - K_0[u_*](\hat{t}, \hat{x}) \leq \theta$$

for sufficiently small ρ . Notice that this means $K_0[U]$ and $K_0[u_*]$ actually exist.

Since the maximizer (\hat{s}, \hat{y}) of $U - \psi$ is of $w - \psi$ (on Ω) as well, the classical maximum principle for $w - \psi$ implies that $D\phi(\hat{s}, \hat{y}) - 2(\hat{y} - \hat{x}) = D\psi(\hat{s}, \hat{y})$. Hence we see $\lim_{\rho \rightarrow 0} D\psi(\hat{s}, \hat{y}) = D\phi(\hat{t}, \hat{x})$. Moreover, $\lim_{\rho \rightarrow 0} U(\hat{s}, \hat{y}) = \lim_{\rho \rightarrow 0} w(\hat{s}, \hat{y}) = \phi(\hat{t}, \hat{x}) = u_*(\hat{t}, \hat{x})$. Therefore

$$H(\hat{s}, \hat{y}, U(\hat{s}, \hat{y}), D\psi(\hat{s}, \hat{y})) - H(\hat{t}, \hat{x}, u_*(\hat{t}, \hat{s}), D\phi(\hat{t}, \hat{x})) \leq \theta$$

if ρ is sufficiently small. Summing up the above we obtain for sufficiently small ρ that

$$\begin{aligned} &K_0[U](\hat{s}, \hat{y}) + H(\hat{s}, \hat{y}, U(\hat{s}, \hat{y}), D\psi(\hat{s}, \hat{y})) \\ &\leq -2\theta + K_0[U](\hat{s}, \hat{y}) - K_0[u_*](\hat{t}, \hat{x}) \\ &\quad + H(\hat{s}, \hat{y}, U(\hat{s}, \hat{y}), D\psi(\hat{s}, \hat{y})) - H(\hat{t}, \hat{x}, u_*(\hat{t}, \hat{s}), D\phi(\hat{t}, \hat{x})) \leq 0, \end{aligned}$$

which shows that U is a subsolution of (1.1).

Theorem 3.1 implies that $U \leq u^+$. Let $\{(t_j, x_j)\}_{j \geq 0}$ be a sequence such that $(t_j, x_j, u(t_j, x_j)) \rightarrow (\hat{t}, \hat{x}, u_*(\hat{t}, \hat{x}))$ as $j \rightarrow \infty$. Then

$$\liminf_{j \rightarrow \infty} (U(t_j, x_j) - u(t_j, x_j)) \geq \lim_{j \rightarrow \infty} (w(t_j, x_j) - u(t_j, x_j)) = \frac{\rho^2}{2} > 0.$$

In other words there exists a point (s, y) such that $U(s, y) > u(s, y)$. Therefore the proof is complete. \square

Corollary 4.3 (Unique existence for (1.1)-(1.2)). *Assume (A1)-(A4). Then there exists at most one solution u of (1.1)-(1.2).*

Proof. The uniqueness of a solution is guaranteed by Theorem 3.1. Henceforth, it is enough to construct u^- and u^+ in Theorem 4.2 so that u defined by (4.4) satisfies $u(0, \cdot) = u_0$ on \mathbb{T}^d .

Set $\omega(\ell) := \sup\{|u_0(\zeta) - u_0(\eta)| \mid \zeta, \eta \in \mathbb{T}^d, |\zeta - \eta| \leq \ell\}$ for $\ell \geq 0$ and $f_y(x) := \sum_{i=1}^d (1 - \cos(2\pi(x_i - y_i)))$ for $x, y \in \mathbb{T}^d$, where x_i and y_i are i -th components of each variable. Then for each $\varepsilon > 0$ there exists a constant $C_\varepsilon > 0$ such that $\omega(|x - y|) \leq \varepsilon + C_\varepsilon f_y(x)$ for all $x, y \in \mathbb{T}^d$. For $\varepsilon \in (0, 1)$ and $y \in \mathbb{T}^d$ we define a function $u_{\varepsilon, y}^- : \overline{Q_T} \rightarrow \mathbb{R}$ by

$$u_{\varepsilon, y}^-(t, x) := u_0(y) - \varepsilon - C_\varepsilon f_y(x) - \frac{Ct^\alpha}{\Gamma(1 + \alpha)},$$

where $C > 0$ is a large constant. Then $u_{\varepsilon, y}^- \in \mathcal{C}^1(\overline{Q_T})$. Moreover $u_{\varepsilon, y}^- \leq u_0(y)$ by the non-negativity of f_y and $|Du_{\varepsilon, y}^-|$ is bounded on Q_T . It is well-known that

$$(4.9) \quad \frac{1}{\Gamma(1 - \alpha)} \int_a^t \frac{[(s - a)^\beta]^\alpha}{(t - s)^\alpha} ds = \frac{\Gamma(\beta + 1)}{\Gamma(\beta - \alpha + 1)} (t - a)^{\beta - \alpha}$$

for given constants $a \in \mathbb{R}$ and $\beta \in (0, 1)$; see [35, (2.56)] for the proof. From this formula with $(a, \beta) = (0, \alpha)$ and the above, we see

$$-C + H(t, x, u_{\varepsilon, y}^-(t, x), Du_{\varepsilon, y}^-(t, x)) \leq 0$$

for all $(t, x) \in Q_T$. Thus Proposition 2.10 implies that $u_{\varepsilon, y}^-$ is a (viscosity) subsolution of (1.1).

We also see

$$u_{\varepsilon, y}^-(t, x) \leq u_0(x) + \omega(|x - y|) - \varepsilon - C_\varepsilon f_y(x) - \frac{Ct^\alpha}{\Gamma(1 - \alpha)} \leq u_0(x)$$

for all $(t, x) \in \overline{Q_T}$. Therefore, Lemma 4.1 ensures that

$$u^-(t, x) := (\sup\{u_{\varepsilon, y}^-(t, x) \mid \varepsilon \in (0, 1), y \in \mathbb{T}^d\})^*$$

is a subsolution of (1.1) and satisfies $u^-(t, x) \leq u_0(x)$ for all $(t, x) \in \overline{Q_T}$. The definition of u^- yields $u^-(0, \cdot) \geq u_0$ on \mathbb{T}^d , which guarantees that $(u^-)_* > -\infty$ on $\overline{Q_T}$ and $u^-(0, \cdot) = u_0$ on \mathbb{T}^d . Similarly, a supersolution with desired properties is constructed. Moreover, it turns out that u^\pm satisfy $u_0(x) = \lim_{(t, y) \rightarrow (x, 0)} u^\pm(t, y)$ but we leave the verification to the reader; cf. [12]. With u^\pm above, we obtain a solution u by Theorem 4.2, and it satisfies $u(0, \cdot) = u_0$ on \mathbb{T}^d . The proof is now complete. \square

5. SOME STABILITY RESULTS

Two main theorems in this section are in what follows:

Theorem 5.1 (Stability I). *Let H_ε and H satisfy (A1)-(A3), where $\varepsilon > 0$. Let $u_\varepsilon \in USC(\overline{Q_T})$ (resp. $LSC(\overline{Q_T})$) be a subsolution (resp. supersolution) of*

$$\partial_t^\alpha u_\varepsilon + H_\varepsilon(t, x, u_\varepsilon, Du_\varepsilon) = 0 \quad \text{in } Q_T.$$

Assume that H_ε converges to H as $\varepsilon \rightarrow 0$ locally uniformly in $(0, T] \times \mathbb{T}^d \times \mathbb{R} \times \mathbb{R}^d$. Assume that $\{u_\varepsilon\}_{\varepsilon>0}$ is locally uniformly bounded. Then $u := \limsup^* u_\varepsilon$ (resp. $\liminf^* u_\varepsilon$) is a subsolution (resp. supersolution) of

$$\partial_t^\alpha u + H(t, x, u, Du) = 0 \quad \text{in } Q_T.$$

Here $\limsup^* u_\varepsilon$ appears above is the *upper relaxed limit* defined by

$$(5.1) \quad (\limsup_{\varepsilon \rightarrow 0}^* u_\varepsilon)(t, x) := \limsup_{\delta \searrow 0} \{u_\varepsilon(s, y) \mid (s, y) \in Q_T \cap \overline{B_\delta(t, x)}, 0 < \varepsilon < \delta\}$$

for $(t, x) \in \overline{Q_T}$ and $\liminf^* u_\varepsilon := -\limsup^*(-u_\varepsilon)$ is the *lower relaxed limit*.

Theorem 5.2 (Stability II). *Assume that (A1)-(A4). Let $u_\alpha \in C(\overline{Q_T})$ be a solution of (1.1)-(1.2) whose time-derivative's order is $\alpha \in (0, 1)$. Then u_α converges to u_β locally uniformly in Q_T as $\alpha \rightarrow \beta$, where u_β is a solution of (1.1)-(1.2) whose time-derivative's order is $\beta \in (0, 1]$.*

A same idea as for the proof of [4, Theorem V.1.7] is used for Theorem 5.1 and Theorem 5.2. A deal for the term of time-derivative is the only difference between Theorem 5.1 and [4, Theorem V.1.7] but it is similar between Theorem 5.1 and Theorem 5.2. For this reason we only prove Theorem 5.2.

Proof. As the analogy of the upper/lower relaxed limits, for $\beta \in (0, 1]$, we define functions u^\sharp and u_\sharp by

$$u^\sharp(t, x) := \limsup_{\delta \searrow 0} \{u_\alpha(s, y) \mid (s, y) \in \overline{B_\delta(t, x)} \cap Q_T, \alpha \in (\beta - \delta, \beta + \delta) \cap (0, 1)\}$$

and $u_\sharp := -(-u)^\sharp$ on $\overline{Q_T}$. In Remark 6.3 later we will mention that $\{u_\alpha\}_{\alpha \in (0, 1]}$ of (1.1)-(1.2) is uniformly bounded on $\overline{Q_T}$. Hence u^\sharp and u_\sharp are bounded on $\overline{Q_T}$. Note also that u^\sharp is an upper semicontinuous function, so u_\sharp is a lower semicontinuous function.

We shall show that u^\sharp and u_\sharp are a subsolution and a supersolution of (1.1)-(1.2) whose time-derivative's order is β . It suffices to show that u^\sharp is a subsolution of (1.1) since the similar argument is applied for u_\sharp and it is clear that $u^\sharp(0, \cdot) \leq u_0$ and $u_\sharp(0, \cdot) \geq u_0$ on \mathbb{T}^d .

Fix $[a, b] \times B \subset (0, T] \times \mathbb{R}^d$ arbitrarily, where $a < b$ and B is an open set in \mathbb{R}^d . Assume that $u^\sharp - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ for $\phi \in \mathcal{C}^1([0, T] \times \mathbb{R}^d)$. Let $\{\alpha_j\}_{j \geq 0}$ and $\{(t_j, x_j)\}_{j \geq 0}$ be sequences such that $u_{\alpha_j} - \phi$ attains a maximum at $(t_j, x_j) \in (a, b) \times B$ over $[a, b] \times \overline{B}$ and

$$(\alpha_j, t_j, x_j, u_{\alpha_j}(t_j, x_j)) \rightarrow (\beta, \hat{t}, \hat{x}, u^\sharp(\hat{t}, \hat{x}))$$

as $j \rightarrow \infty$. A proof of existence of such sequences is essentially same as for [4, Lemma V.1.6] and not difficult, so we omit it.

Case 1: $\beta \neq 1$. Since u_{α_j} is a subsolution of (1.1),

$$(5.2) \quad J_{t_j-a}^{\alpha_j}[\psi_j](t_j, x_j) + K_{t_j-a}^{\alpha_j}[u_{\alpha_j}](t_j, x_j) + H(t_j, x_j, u_{\alpha_j}(t_j, x_j), D\phi(t_j, x_j)) \leq 0.$$

Here $J_r^{\alpha_j}$ and $K_r^{\alpha_j}$ are associated with $\alpha = \alpha_j$. By similar arguments in previous sections it can be turns out that

$$(5.3) \quad \liminf_{j \rightarrow \infty} (J_{t_j-a}^{\alpha_j}[\psi_j](t_j, x_j) + K_{t_j-a}^{\alpha_j}[u_{\alpha_j}](t_j, x_j)) \geq J_{\hat{t}-a}^\beta[\phi](\hat{t}, \hat{x}) + K_{\hat{t}-a}^\beta[u^\sharp](\hat{t}, \hat{x}).$$

Since

$$\lim_{j \rightarrow \infty} H(t_j, x_j, u_{\alpha_j}(t_j, x_j), D\phi(t_j, x_j)) = H(\hat{t}, \hat{x}, u^\sharp(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})),$$

we find that u^\sharp is a subsolution of (1.1).

Case 2: $\beta = 1$. There are similar sequences $\{\alpha_j\}_j$ and $\{(t_j, x_j)\}_j$ even for $\varphi \in C^1((0, T] \times \mathbb{R}^d)$ instead of $\phi \in C^1([0, T] \times \mathbb{R}^d)$. Since $(t_j, x_j) \rightarrow (\hat{t}, \hat{x}) \in (a, b) \times B$ as $j \rightarrow \infty$, we may assume that $\{(t_j, x_j)\}_j \subset (\hat{t} - 3\delta, \hat{t}] \times \overline{B_{2\delta}(\hat{x})}$ by considering large j , where $\delta > 0$ is a constant such that $[\hat{t} - 3\delta, \hat{t}] \times \overline{B_{2\delta}(\hat{x})} \subset (a, b) \times B$. Let $\xi_1, \xi_2 : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ be C^∞ functions such that $\xi_1 + \xi_2 = 1$ on $[0, T] \times \mathbb{R}^d$, $\xi_1 = 1$ on $[\hat{t} - \delta, \hat{t}] \times B_\delta(\hat{x})$ and $\xi_2 = 1$ on $([0, T] \times \mathbb{R}^d) \setminus ([\hat{t} - 2\delta, \hat{t}] \times B_{2\delta}(\hat{x}))$. Since $\{u_\alpha\}_{\alpha \in (0, 1]}$ is uniformly bounded on $\overline{Q_T}$, there is a constant $C > 0$ such that $\max_{\overline{Q_T}} |u_{\alpha_j}| \leq C$ for all j . Set $\psi = \xi_1 \phi + \xi_2 M$, where $M := C + 1$. Then $\psi \in C^1([0, T] \times \mathbb{R}^d) \subset C^1([0, T] \times \mathbb{R}^d)$ and $u_{\alpha_j} - \psi$ attains a maximum at $(t_j, x_j) \in (0, T] \times \mathbb{R}^d$. Thus we have

$$(5.4) \quad K_0[u_{\alpha_j}](t_j, x_j) + H(t_j, x_j, u_{\alpha_j}(t_j, x_j), D\psi(t_j, x_j)) \leq 0.$$

Note that $K_0[u_{\alpha_j}](t_j, x_j) \geq \partial_t^{\alpha_j} \psi(t_j, x_j)$. According to [9, Theorem 2.10] we notice that

$$\lim_{j \rightarrow \infty} \partial_t^{\alpha_j} \psi(t_j, x_j) = \partial_t \psi(\hat{t}, \hat{x}).$$

Thus estimating $K_0[u_{\alpha_j}](t_j, x_j)$ by $\partial_t^{\alpha_j} \psi(t_j, x_j)$ in (5.4) and then passing to the limit $j \rightarrow \infty$ implies that

$$\partial_t \psi(\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u^\sharp(\hat{t}, \hat{x}), D\psi(\hat{t}, \hat{x})) \leq 0.$$

Since $\partial_t \psi(\hat{t}, \hat{x}) = \partial_t \phi(\hat{t}, \hat{x})$ and $D\psi(\hat{t}, \hat{x}) = D\phi(\hat{t}, \hat{x})$, u^\sharp is a subsolution of (1.1) with $\alpha = 1$.

The comparison principle implies that $u^\sharp \leq u_\#$ on $\overline{Q_T}$ but $u_\# \leq u^\sharp$ by their definition. We hence see that $u := u^\sharp = u_\#$ is a solution of (1.1) and $u(0, \cdot) = u_0$ on \mathbb{T}^d . Corollary 3.4 ensures that $u = u_\beta$, a conclusion. \square

6. REGULARITY RESULTS

Let consider one-dimensional transport equations of the form

$$\partial_t^\alpha u + \partial_x u = 0 \quad \text{in } (0, \infty) \times \mathbb{R}$$

with prescribed initial value $u|_{t=0} = u_0 \in C(\mathbb{R})$. In [33] for instance, a solution of this equation was given as

$$(6.1) \quad u(t, x) = \frac{1}{t^\alpha} \int_0^\infty W_{-\alpha, 1-\alpha} \left(-\frac{z}{t^\alpha} \right) u_0(x - z) dz$$

through the Laplace and the inverse Laplace transformation. Here $W_{-\alpha, 1-\alpha}$ is Wright function defined by

$$W_{-\alpha, 1-\alpha}(z) := \sum_{j=0}^{\infty} \frac{z^j}{j! \Gamma(-\alpha j + 1 - \alpha)}.$$

For properties and formulae for Wright function, see [35] and references therein. It can be verified that this solution is indeed a unique viscosity solution. We leave the detail of calculations to the reader.

Let us assume that

(A4') u_0 is a Lipschitz continuous function with the Lipschitz constant $\text{Lip}[u_0]$.

We shall prove a continuity of solutions with respect to each variable. In space, since $W_{-\alpha,1-\alpha} \geq 0$ on $(0, \infty)$ and

$$\int_0^\infty W_{-\alpha,1-\alpha}(-z)dz = 1,$$

we have

$$\begin{aligned} |u(t, x) - u(t, y)| &\leq \frac{1}{t^\alpha} \int_0^\infty W_{-\alpha,1-\alpha}\left(-\frac{z}{t}\right) |u_0(x-z) - u_0(y-z)| dz \\ &\leq \frac{1}{t^\alpha} \int_0^\infty W_{-\alpha,1-\alpha}\left(-\frac{z}{t}\right) dz \text{Lip}[u_0] |x-y| = \text{Lip}[u_0] |x-y| \end{aligned}$$

for $(t, x, y) \in [0, T] \times \mathbb{R} \times \mathbb{R}$. In time, since

$$\int_0^\infty W_{-\alpha,1-\alpha}(-z)z dz = \frac{1}{\Gamma(\alpha+1)}$$

and u given by (6.1) is rewritten as

$$u(t, x) = \int_0^\infty W_{-\alpha,1-\alpha}(-z)u_0(x-t^\alpha z)dz,$$

we have

$$\begin{aligned} |u(t, x) - u(s, x)| &\leq \int_0^\infty W_{-\alpha,1-\alpha}(-z) |u_0(x-t^\alpha z) - u_0(x-s^\alpha z)| dz \\ &\leq \int_0^\infty W_{-\alpha,1-\alpha}(-z)z dz \text{Lip}[u_0] |t^\alpha - s^\alpha| \leq \frac{\text{Lip}[u_0]}{\Gamma(\alpha+1)} |t-s|^\alpha \end{aligned}$$

for $(t, s, x) \in [0, T] \times [0, T] \times \mathbb{R}$. These regularity results hold for solutions of (1.1)-(1.2) with some general Hamiltonians H under some Lipschitz continuity in x for H as for $\alpha = 1$.

Lemma 6.1 (Lipschitz preserving). *Assume (A1), (A3), (A4') and that there exist constants $L_1 \geq 0$ and $L_2 > 0$ such that*

$$|H(t, x, r, p) - H(t, y, r, p)| \leq L_1|x-y| + L_2|x-y||p|$$

for all $(t, x, y, r, p) \in (0, T] \times \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$. Let $u \in C(\overline{Q_T})$ be a solution of (1.1)-(1.2). Then $|u(t, x) - u(t, y)| \leq L(t)|x-y|$ for all $(t, x, y) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ with

$$L(t) = \left(\text{Lip}[u_0] + \frac{L_1}{L_2} \right) E_\alpha(L_2 t^\alpha) - \frac{L_1}{L_2}.$$

Proof. For δ we set

$$\Phi_\delta(t, x, y) := u(t, x) - u(t, y) - L_\delta(t)|x-y|,$$

where

$$L_\delta(t) = \left(\text{Lip}[u_0] + \frac{L_1 + \delta}{L_2} \right) E_\alpha(L_2 t^\alpha) - \frac{L_1 + \delta}{L_2}.$$

Note that $L_\delta \in C^1((0, T]) \cap C([0, T])$ and $L'_\delta \in L^1(0, T)$. Suppose by contradiction that there would exit $\delta > 0$ such that $\max_{[0, T] \times \mathbb{R}^d \times \mathbb{R}^d} \Phi_\delta > 0$.

Let $(\hat{t}, \hat{x}, \hat{y}) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$ be a maximum point of Φ . Note that $\hat{t} > 0$ and $\hat{x} \neq \hat{y}$; otherwise $0 < \Phi(\hat{t}, \hat{x}, \hat{x}) = 0$ or $0 < \Phi(0, \hat{x}, \hat{y}) \leq 0$ since $u(0, \cdot) = u_0$ is Lipschitz continuous and $E_\alpha(0) = 1$. Moreover $u(\hat{t}, \hat{x}) \geq u(\hat{t}, \hat{y})$ from $\Phi_\delta(\hat{t}, \hat{x}, \hat{y}) > 0$.

Since u is a solution of (1.1), we have from Lemma 3.3

$$(6.2) \quad K_0[u](\hat{t}, \hat{x}) - K_0[u](\hat{t}, \hat{y}) + H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), L_\delta(\hat{t})\hat{p}) - H(\hat{t}, \hat{y}, u(\hat{t}, \hat{y}), L_\delta(\hat{t})\hat{p}) \leq 0,$$

where $\hat{p} := (\hat{x} - \hat{y})/|\hat{x} - \hat{y}|$. It follows from the definition of Φ that

$$K_0[u](\hat{t}, \hat{x}) - K_0[u](\hat{t}, \hat{y}) \geq K_0[L_\delta](\hat{t})|\hat{x} - \hat{y}| = (\partial_t^\alpha L_\delta)(\hat{t})|\hat{x} - \hat{y}|.$$

Hamiltonians are estimates as

$$\begin{aligned} & H(\hat{t}, \hat{x}, u(\hat{t}, \hat{x}), L_\delta(\hat{t})\hat{p}) - H(\hat{t}, \hat{y}, u(\hat{t}, \hat{y}), L_\delta(\hat{t})\hat{p}) \\ & \geq H(\hat{t}, \hat{x}, u(\hat{t}, \hat{y}), L_\delta(\hat{t})\hat{p}) - H(\hat{t}, \hat{y}, u(\hat{t}, \hat{y}), L_\delta(\hat{t})\hat{p}) \\ & \geq -L_1|\hat{x} - \hat{y}| - L_2L_\delta(\hat{t})|\hat{x} - \hat{y}| \end{aligned}$$

by (A3) and (A2') with $|\hat{p}| = 1$. Thus (6.2) is led to

$$[(\partial_t^\alpha L_\delta)(\hat{t}) - L_2L_\delta(\hat{t}) - L_1]|\hat{x} - \hat{y}| \leq 0.$$

Since L_δ satisfies

$$\partial_t^\alpha L_\delta - L_2L_\delta - L_1 = \delta$$

according to [9, Theorem 7.2, Remark 7.1], we obtain

$$\delta|\hat{x} - \hat{y}| \leq 0,$$

a contradiction.

Consequently for any $\delta > 0$ we see that $|u(t, x) - u(t, y)| \leq L_\delta|x - y|$ for all $(t, x, y) \in [0, T] \times \mathbb{R}^d \times \mathbb{R}^d$. Letting $\delta \rightarrow 0$ yields the conclusion. \square

Lemma 6.2. *Assume (A1), (A2), (A3) and (A4'). Let $u \in C(\overline{Q_T})$ be a solution of (1.1)-(1.2). Then there exists a constant $M > 0$ depending only on H, u_0, α and T such that*

$$|u(t, x) - u_0(x)| \leq Mt^\alpha$$

for all $(t, x) \in \overline{Q_T}$.

Proof. We find that $u^-(t, x) := u_0(x) - Mt^\alpha$ and $u^+(t, x) := u_0(x) + Mt^\alpha$ are a subsolution and a supersolution of (1.1)-(1.2), respectively, where the constant M is chosen so large that

$$M \geq \sup\{\Gamma^{-1}(\alpha + 1)|H(t, x, \max_{\mathbb{T}^d} |u_0|, p)| \mid (t, x) \in Q_T, |p| \leq \text{Lip}[u_0], \alpha \in (0, 1)\}.$$

In fact, if $u^- - \phi$ attains a maximum at $(\hat{t}, \hat{x}) \in (0, T] \times \mathbb{R}^d$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$, then $|D\phi(\hat{t}, \hat{x})| \leq \text{Lip}[u_0]$ and $K_0[u^-](\hat{t}, \hat{x}) = -M\partial_t^\alpha t^\alpha|_{t=\hat{t}} = \Gamma(1 + \alpha)M$ by using the formula $\partial_t^\alpha t^\alpha = \Gamma(1 + \alpha)$ derived from (4.9). Therefore

$$K_0[u^-](\hat{t}, \hat{x}) + H(\hat{t}, \hat{x}, u^-(\hat{t}, \hat{x}), D\phi(\hat{t}, \hat{x})) \leq -\Gamma(1 + \alpha)M + H(\hat{t}, \hat{x}, \max_{\mathbb{T}^d} |u_0|, D\phi(\hat{t}, \hat{x})) \leq 0.$$

by (A3) and the choice of M since $u^-(t, x) = u_0(x) - Mt^\alpha \leq \max_{\mathbb{T}^d} |u_0|$. Similarly, it is verified that u^+ is a supersolution of (1.1).

Theorem 3.1 (comparison principle) yields to

$$u_0(x) - Mt^\alpha = u^-(t, x) \leq u(t, x) \leq u^+(t, x) = u_0(x) + Mt^\alpha$$

on $\overline{Q_T}$, which is noting but the desired estimate. \square

Remark 6.3. Let u_α be a solution of (1.1)-(1.2) whose time-derivative's order is $\alpha \in (0, 1]$. For $u_0 \in C(\mathbb{T}^d)$ not necessarily Lipschitz continuous, we observe that

$$(6.3) \quad \sup\{|u_\alpha(t, x)| \mid (t, x) \in \overline{Q_T}, \alpha \in (0, 1]\} \leq \max_{\mathbb{T}^d} |u_0| + C \max\{1, T\}.$$

Here $C > 0$ is a large constant so that

$$C \geq \sup\{\Gamma^{-1}(\alpha + 1)|H(t, x, \max_{\mathbb{T}^d} |u_0|, 0)| \mid (t, x) \in \overline{Q_T}, \alpha \in (0, 1]\}.$$

Indeed, $\max_{\mathbb{T}^d} |u_0| - Ct^\alpha$ and $-\max_{\mathbb{T}^d} |u_0| + Ct^\alpha$ are a (classical) subsolution and a (classical) supersolution of (1.1)-(1.2), respectively. Thus the comparison principle implies (6.3) once one realizes that $t^\alpha \leq T^\alpha \leq \max\{1, T\}$ for all t, T and $\alpha \in (0, 1]$.

Lemma 6.4 (Temporal Hölder continuity). *Assume (A1), (A2), (A3) and (A4'). Let $u \in C(\overline{Q_T})$ be a solution of (1.1)-(1.2). Then for the same constant $M > 0$ as in Lemma 6.2*

$$(6.4) \quad |u(t, x) - u(s, x)| \leq M|t - s|^\alpha$$

for all $(t, s, x) \in [0, T] \times [0, T] \times \mathbb{T}^d$.

Proof. Let X be a set of subsolutions v of (1.1)-(1.2) such that v satisfy (6.4) and $u_0 - Mt^\alpha \leq v \leq u_0 + Mt^\alpha$ on $\overline{Q_T}$. Notice that $X \neq \emptyset$ since $u_0(x) - Mt^\alpha \in X$ due to Lemma 6.2. Define $u = \sup\{v \mid v \in X\}$. We show by Perron's method that u is a solution of (1.1)-(1.2) satisfying (6.4). In view of Corollary 4.3 it is enough to prove that u is a solution of (1.1) satisfying (6.4). In this proof we use same notations associated to the above u as in Theorem 4.2.

It is not hard to see that

$$(6.5) \quad |u(t, x) - u(s, x)| \leq \sup\{|v(t, x) - v(s, x)| \mid v \in X\}$$

for all $t, s \in [0, T]$ and $x \in \mathbb{T}^d$. Since M does not depend unknown functions and v satisfies (6.4), we see that u satisfies (6.4). Let us prove that u is a solution of (1.1).

To do so we must show that the function U satisfies (6.4). All processes except for this step work to the current situation. We first show that the function w satisfies (6.4) near (\hat{t}, \hat{x}) . Expanding w by Taylor formula, we have

$$\begin{aligned} w(s, y) - w(\hat{t}, \hat{x}) &= \phi(s, y) - \phi(\hat{t}, \hat{x}) - |s - \hat{t}|^2 - |y - \hat{x}|^2 \\ &= a(s - \hat{t}) + p \cdot (y - \hat{x}) + o(|s - \hat{t}| + |y - \hat{x}|) - |s - \hat{t}|^2 - |y - \hat{x}|^2 \end{aligned}$$

for $Q_T \ni (s, y) \rightarrow (\hat{t}, \hat{x})$, where $a = \partial_t \phi(\hat{t}, \hat{x})$ and $p = D\phi(\hat{t}, \hat{x})$. For every $\eta > 0$ there exists $\delta > 0$ such that

$$|w(s, y) - w(\hat{t}, \hat{x})| \leq ((|a| + \eta)|s - \hat{t}|^{1-\alpha} + |s - \hat{t}|^{2-\alpha})|s - \hat{t}|^\alpha + (|p| + \eta)|y - \hat{x}| + |y - \hat{x}|^2$$

for all $(s, y) \in \overline{B_\delta(\hat{t}, \hat{x})}$. Fix such $\eta > 0$. For δ so that $(|a| + \eta)\delta^{1-\alpha} + \delta^{2-\alpha} \leq M$, w satisfies (6.4) on $\overline{B_\delta(\hat{t}, \hat{x})}$.

Let ρ be taken so small that $\rho \leq \sqrt{2}\delta$. Then $\Omega \subset B_\delta(\hat{t}, \hat{x})$. Since u and w satisfy (6.4) in $((\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})) \cap B_\delta(\hat{t}, \hat{x})$, it turns out by a similar inequality for the function $\max\{u^*, w\}$ as (6.5) that $\max\{u^*, w\}$ satisfies (6.4) in the same region. Since $u^* > w$ in $((\hat{t} - \rho, \hat{t} + \rho) \times B_{2\rho}(\hat{x})) \setminus \Omega$, in consequence, U satisfies (6.4) in $\overline{Q_T}$, a conclusion. \square

7. ANOTHER POSSIBLE DEFINITION

In this section, for simplicity, we only treat Hamiltonians independent of t and r , i.e., $H = H(x, p)$ and assume (A1)-(A2). Following to the usual style for viscosity solutions we are also able to define weak solutions of (1.1) as follows:

Definition 7.1 (Provisional solutions). For a function $u \in C(\overline{Q_T})$ we call u a *provisional subsolution* (resp. *provisional supersolution*) of (1.1) if

$$(\partial_t^\alpha \phi)(\hat{t}, \hat{x}) + H(\hat{x}, D\phi(\hat{t}, \hat{x})) \leq 0 \quad (\text{resp. } \geq 0)$$

whenever $u - \phi$ attains a maximum (resp. minimum) at $(\hat{t}, \hat{x}) \in Q_T$ over $\overline{Q_T}$ for $\phi \in C^1([0, T] \times \mathbb{R}^d)$. If $u \in C(\overline{Q_T})$ is a both provisional sub- and supersolution of (1.1), then we call u a *provisional solution* of (1.1).

It is no difficulties to prove that provisional solutions of (1.1) are consistent with classical solutions of (1.1) if they belong to $C^1(\overline{Q_T})$; cf. Proposition 2.10 and [30, Theorem 1]. Although Definition 7.1 looks good, there are some technical difficulties to handle provisional solutions. We conclude this paper by sharing a main part of such difficulties.

They occurs in a proof of comparison principle. Let u and v be respectively a provisional subsolution and a provisional supersolution of (1.1) such that $u(0, \cdot) \leq v(0, \cdot)$ on \mathbb{T}^d . Suppose that $\max_{\overline{Q_T}}(u - v) > 0$ and aim to derive a contradiction.

There is a small constant $\eta > 0$ such that

$$\max_{(t,x) \in Q_T} ((u - v)(t, x) - \eta t^\alpha) =: \theta > 0.$$

For $\varepsilon > 0$ and $\delta > 0$ we consider the function

$$\Phi(t, s, x, y) := u(t, x) - v(s, y) - \frac{|x - y|^2}{2\varepsilon} - \frac{|t - s|^2}{2\delta} - \eta t^\alpha.$$

on $[0, T]^2 \times \mathbb{T}^{2d}$. Let $(\bar{t}, \bar{s}, \bar{x}, \bar{y})$ be a maximum point of Φ . From inequalities for provisional sub- and supersolutions, we have

$$(7.1) \quad \left(\partial_t^\alpha \frac{|\cdot - \bar{s}|^2}{2\delta} \right) (\bar{t}, \bar{s}) + \left(\partial_s^\alpha \frac{|\bar{t} - \cdot|^2}{2\delta} \right) (\bar{t}, \bar{s}) + \eta \Gamma(1 + \alpha) + H(\bar{x}, \bar{p}) - H(\bar{y}, \bar{p}) \leq 0.$$

Here $\bar{p} := (\bar{x} - \bar{y})/\varepsilon$. A similar argument is found in Section 3 of this paper. The third term comes from the last term of Φ , i.e. ηt^α . Let us focus on the first and second terms. A direct calculation implies that

$$\partial_t^\alpha |t - s|^2 = \frac{2(t - (2 - \alpha)s)t^{1-\alpha}}{\Gamma(3 - \alpha)}$$

by the formula (4.9). By changing the role of t and s , consequently, we get

$$(7.2) \quad \begin{aligned} \left(\partial_t^\alpha \frac{|\cdot - \bar{s}|^2}{2\delta} \right) (\bar{t}) + \left(\partial_s^\alpha \frac{|\bar{t} - \cdot|^2}{2\delta} \right) (\bar{s}) &= \frac{(\bar{t} - (2 - \alpha)\bar{s})\bar{t}^{1-\alpha} + (\bar{s} - (2 - \alpha)\bar{t})\bar{s}^{1-\alpha}}{\delta \Gamma(1 - \alpha)} \\ &= \frac{\bar{t}^{2-\alpha} + \bar{s}^{2-\alpha} - (2 - \alpha)(\bar{s}\bar{t}^{1-\alpha} + \bar{t}\bar{s}^{1-\alpha})}{\delta \Gamma(3 - \alpha)}. \end{aligned}$$

When $\alpha = 1$, (7.2) vanishes. Thus estimating Hamiltonians suitably (see the proof of Theorem 3.1 in this paper) and then passing to the limit $\varepsilon, \delta \rightarrow 0$ in (7.1) yields the contradiction thanks to the third term. On the other hand, the situation for $\alpha \in (0, 1)$ is completely different. Indeed, (7.2) possibly does not vanish and it

is hard to control (\bar{t}, \bar{s}) so that (7.2) is sufficiently small comparing to $\eta\Gamma(1 + \alpha)$ as $\delta \rightarrow 0$. There is a possibility that (7.2) diverges as $\delta \rightarrow 0$ as well.

To solve above difficulties let us consider the following problem:

Problem 7.2. Find a function $\psi \in C^1((0, T]^2; \mathbb{R}) \cap C([0, T]^2)$ satisfying $\partial_t \psi(\cdot, s) \in L^1(0, T)$ for every $s \in [0, T]$ and $\partial_s \psi(t, \cdot) \in L^1(0, T)$ for every $t \in [0, T]$ such that

$$(7.3) \quad \begin{cases} \partial_t^\alpha \psi(\cdot, s) + \partial_s^\alpha \psi(t, \cdot) \geq 0 & \text{on } (0, T)^2, \\ \psi = 0 & \text{on } \{t = s\} \text{ and} \\ \psi > 0 & \text{on } [0, T]^2 \setminus \{t = s\}. \end{cases}$$

If we could find such a function, then the contradiction would be obtained by handling

$$\Psi(t, s, x, y) := u(t, x) - v(s, y) - \frac{|x - y|^2}{2\varepsilon} - \frac{\psi(t, s)}{2\delta} - \eta t^\alpha$$

instead of Φ . However, such a modification unfortunately does not overcome the difficulty yet.

Proposition 7.3. *There is no function ψ solving Problem 7.2.*

Proof. Suppose by contradiction that there is a function ψ solves Problem 7.2. Then ψ should satisfy

$$(\partial_t^\alpha \psi)(t, t) + (\partial_s^\alpha \psi)(t, t) \geq 0,$$

that is,

$$(7.4) \quad \int_0^t \frac{(\partial_t \psi)(\tau, t)}{(t - \tau)^\alpha} d\tau + \int_0^t \frac{(\partial_s \psi)(t, \tau)}{(t - \tau)^\alpha} d\tau \geq 0.$$

Since $\psi(t, t) = 0$ and $\psi(\cdot, t) \in C^1(0, T)$, integration by parts implies that

$$\int_0^t \frac{(\partial_t \psi)(\tau, t)}{(t - \tau)^\alpha} d\tau = -\frac{\psi(0, t)}{t^\alpha} - \alpha \int_0^t \frac{\psi(t, \tau)}{(t - \tau)^{\alpha+1}} d\tau.$$

Thus (7.4) is rewritten as

$$\frac{\psi(0, t) + \psi(t, 0)}{t^\alpha} + \alpha \int_0^t \frac{\psi(t, \tau) + \psi(\tau, t)}{(t - \tau)^{\alpha+1}} d\tau \leq 0.$$

However the left-hand side is positive since $\psi > 0$ on $[0, T]^2 \setminus \{t = s\}$, a contradiction. \square

Acknowledgements. The authors are grateful to Professor Masahiro Yamamoto and members of his group including Ms. Anna Suzuki for valuable information on this topic. The authors would like to thank Professor Adam Kubica for careful reading of the manuscript and beneficial discussions. The first author is partly supported by the Japan Society for the Promotion of Science (JSPS) through grant No. 26220702 (Kiban S) and No. 16H03948 (Kiban B). The second author is supported by Grant-in-aid for Scientific Research of JSPS Fellows No. 16J03422 and the Program for Leading Graduate Schools, MEXT, Japan.

REFERENCES

- [1] Achdou, Y., Barles, G., Ishii, H., Litvinov, G. L. (2013). Hamilton-Jacobi equations: approximations, numerical analysis and applications. Lecture Notes in Mathematics, 2074. Heidelberg: Springer.
- [2] Alibaud, N., Imbert, C. (2008). Fractional semi-linear parabolic equations with unbounded data. *Trans. Amer. Math. Soc.* 361:2527–2566.
- [3] Allen, M. A nondivergence parabolic problem with a fractional time derivative, preprint, arXiv:1507.04324 [math.AP].
- [4] Bardi, M., Capuzzo Dolcetta, I. (1997). Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations. With appendices by Maurizio Falcone and Pierpaolo Soravia. *Systems & Control: Foundations & Applications*. Boston: Birkhäuser.
- [5] Barles, G., Imbert, C. (2008). Second-order elliptic integro-differential equations: viscosity solutions' theory revisited. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 25:567–585.
- [6] Caffarelli, L., Silvestre, L. (2009). Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.* 62:597–638.
- [7] Crandall, M. G., Ishii, H., Lions, P.-L. (1992). User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc. (N.S.)* 27:1–67.
- [8] Crandall, M. G., Lions P.-L. (1983). Viscosity solutions of Hamilton-Jacobi equations. *Trans. Amer. Math. Soc.* 277:1–42.
- [9] Diethelm, K. (2010). The analysis of fractional differential equations: An application-oriented exposition using differential operators of Caputo type. *Lecture Notes in Mathematics*, 2004. Berlin: Springer-Verlag.
- [10] Fomin, S. A., Chugunov, V. A., Hashida, T. (2011). Non-Fickian mass transport in fractured porous media. *Adv. Water Resour.* 34:205–214.
- [11] Foufoula-Georgiou, E., Ganti, V., Dietrich, W. (2010). A nonlocal theory of sediment transport on hillslopes. *J. Geophys. Res.* 115.
- [12] Giga, Y. (2006). Surface evolution equations: A level set approach. *Monographs in Mathematics*, 99. Basel: Birkhäuser Verlag.
- [13] Gorenflo, R., Kilbas, A. A., Mainardi, F., Rogosin, S. V. (2014). Mittag-Leffler functions, related topics and applications. *Springer Monographs in Mathematics*. Heidelberg: Springer.
- [14] Hamamuki, N. (2014). Asymptotically self-similar solutions to curvature flow equations with prescribed contact angle and their applications to groove profiles due to evaporation-condensation. *Adv. Differential Equations* 19:317–358.
- [15] Harvey, F. R., Lawson, H. B. (2013). The equivalence of viscosity and distributional subsolutions for convex subequations - a strong Bellman principle. *Bull. Braz. Math. Soc. (N.S.)* 44:621–652.
- [16] Imbert, C. (2005). A non-local regularization of first order Hamilton-Jacobi equations. *J. Differential Equations* 211:218–246.
- [17] Ishii, H. (1983). Remarks on Existence of Viscosity Solutions of Hamilton-Jacobi Equations. *Bul. Fac. Sci. Engrg. Chuo Univ.* 26:5–24.
- [18] Ishii, H. (1986). Existence and uniqueness of solutions of Hamilton-Jacobi equations. *Funkcial Ekvac.* 29:167–188.
- [19] Ishii, H. (1995). On the equivalence of two notions of weak solutions, viscosity solutions and distribution solutions. *Funkcial. Ekvac.* 38:101–120.
- [20] Juutinen, P., Lindqvist, P., Manfredi, J. J. (2001). On the equivalence of viscosity solutions and weak solutions for a quasi-linear equation. *SIAM J. Math. Anal.* 33:699–717.
- [21] Juutinen, P., Lukkari, T., Parviainen, M. (2010). Equivalence of viscosity and weak solutions for the $p(x)$ -Laplacian. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 27:1471–1487.
- [22] Kilbas, A. A., Srivastava, H. M., Trujillo, J. J. (2006). *Theory and applications of fractional differential equations*. North-Holland Mathematics Studies, 204. Amsterdam: Elsevier Science B.V..
- [23] Klages, R., Radons, G., Sokolov, I. M. (2008). *Anomalous Transport: Foundations and Applications*. Weinheim: WILEY-VCH Verlag GmbH & Co..
- [24] Koike, S. (2004). *A Beginner's Guide to the Theory of Viscosity Solutions*. MSJ Memoirs, 13. Tokyo: Math. Soc. Japan.
- [25] Kolokoltsovi, V. N., Veretennikova, M. A. (2014). Well-posedness and regularity of the Cauchy problem for nonlinear fractional in time and space equations. *Fractional Differ. Calc.* 4:1–30.

- [26] Kolokoltsovi, V. N., Veretennikova, M. A. (2014). A fractional Hamilton Jacobi Bellman equation for scaled limits of controlled continuous time random walks. *Commun. Appl. Ind. Math.* 6.
- [27] Kubica, A. personal communication.
- [28] Kubica, A., Rybka, P., Ryszewska, K. Weak solutions of fractional differential equations in non cylindrical domain. preprint, arXiv:1605.06775 [math.AP].
- [29] Li, Z., Liu, Y., Yamamoto, M. (2015). Initial-boundary-value problems for the generalized multi-term time-fractional diffusion equation. *Appl. Math. Comput.* 257:381–397.
- [30] Luchko, Y. (2009). Maximum principle for the generalized time-fractional diffusion equation. *J. Math. Anal. Appl.* 351:218–223.
- [31] Luchko, Y. (2010). Some uniqueness and existence results for the initial-boundary-value problems for the generalized time-fractional diffusion equation. *Comput. Math. Appl.* 59:1766–1772.
- [32] Luchko, Y., Rundell, W., Yamamoto, M., Zuo, L. (2013). Uniqueness and reconstruction of an unknown semilinear term in a time-fractional reaction-diffusion equation. *Inverse Problems* 29:065019.
- [33] Mainardi, F., Mura, A., Pagnini, G. (2010). The M-Wright function in time-fractional diffusion processes: A tutorial survey. *Int. J. Differ. Equations* 2010:1-29.
- [34] Montroll, E. W., Weiss, G. H. (1965). Random walks on lattices. II. *J. Math. Phys.* 6:167–181.
- [35] Podlubny, I. (1999). Fractional differential equations: An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. *Mathematics in Science and Engineering*, 198. San Diego: Academic Press.
- [36] Sakamoto, K., Yamamoto, M. (2011). Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.* 382:426–447.
- [37] Samko, S. G., Kilbas, A. A., Marichev, O. I. (1993). Fractional integrals and derivatives: Theory and applications. Yverdon: Gordon and Breach Science Publishers.
- [38] Schumer, R., Meerschaert, M. M., Baeumer, B. (2009). Fractional advection-dispersion equations for modeling transport at the Earth surface. *J. Geophys. Res.* 114.
- [39] Soner, H. M. (1986). Optimal control with state-space constraint. II. *SIAM J. Control Optim.* 24:1110–1122.
- [40] Voller, V. R. (2014). Fractional Stefan problems. *Int. J. Heat Mass Transfer* 74:269–277.
- [41] Zacher, R. (2009). Weak Solutions of Abstract Evolutionary Integro-Differential Equations in Hilbert Spaces. *Funkcial. Ekvac.* 52:1–18.
- [42] Zacher, R. (2012). Global strong solvability of a quasilinear subdiffusion problem. *J. Evol. Equ.* 12:813–831.
- [43] Zacher, R. (2013). A weak Harnack inequality for fractional evolution equations with discontinuous coefficients. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* 12:903–940.
- [44] Zhou, Y. (2014). Basic theory of fractional differential equations. Hackensack: World Scientific Publishing Co. Pte. Ltd..

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914 JAPAN

E-mail address: labgiga@ms.u-tokyo.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914 JAPAN

E-mail address: namba@ms.u-tokyo.ac.jp

UTMS

- 2015–1 Guanyu ZHOU and Norikazu SAITO: *The Navier-Stokes equations under a unilateral boundary condition of Signorini's type.*
- 2015–2 Toru YOSHIYASU: *On Lagrangian embeddings into the complex projective spaces.*
- 2015–3 Norikazu SAITO, Yoshiki SUGITANI and Guanyu ZHOU: *Unilateral problem for the Stokes equations: the well-posedness and finite element approximation.*
- 2015–4 Koya SAKAKIBARA: *Analysis of the dipole simulation method for two-dimensional Dirichlet problems in Jordan regions with analytic boundaries .*
- 2015–5 Junjiro NOGUCHI: *A scalar associated with the inverse of some Abelian integrals on open Riemann surfaces and a ramified Riemann domain .*
- 2015–6 Tomoya KEMMOCHI: *Discrete maximal regularity for abstract Cauchy problems.*
- 2015–7 Tatsu-Hiko MIURA: *Zero width limit of the heat equation on moving thin domains.*
- 2016–1 Norikazu SAITO and Takiko SASAKI: *Finite difference approximation for nonlinear Schrödinger equations with application to blow-up computation.*
- 2016–2 Takiko SASAKI: *Regularity and singularity of blow-up curve for $\partial_t^2 u - \partial_x^2 u = |\partial_t u|^p$.*
- 2016–3 Atsushi KAWAMOTO: *Lipschitz stability estimates in inverse source problems for a fractional diffusion equation of half order in time by Carleman estimates.*
- 2016–4 Yoshikazu GIGA and Naoto KAJIWARA: *On a resolvent estimate for bidomain operators and its applications.*
- 2016–5 Yoshikazu GIGA and Tokinaga NAMBA : *Well-posedness of Hamilton-Jacobi equations with Caputo's time-fractional derivative.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012