

ON THE MINIMAL MODEL THEORY OF NUMERICAL KODAIRA DIMENSION ZERO

YOSHINORI GONGYO

ABSTRACT. We prove the existence of good log minimal models for dlt pairs of numerical log Kodaira dimension 0.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Log minimal model program with scaling	6
4. divisorial Zariski decomposition	8
5. Existence of minimal model in the case where $\nu = 0$	9
6. Abundance theorem in the case where $\nu = 0$	12
References	13

1. INTRODUCTION

Throughout this article, we work over $\mathbb{C}$, the complex number field. We will make use of the standard notation and definitions as in [KM] and [KMM]. The minimal model conjecture for smooth varieties is the following:

Conjecture 1.1 (Minimal model conjecture). *Let X be a smooth projective variety. Then there exists a minimal model or a Mori fiber space of X .*

This conjecture is true in dimension 3 and 4 by Kawamata, Kollár, Mori, Shokurov and Reid (cf. [KMM], [KM] and [Sho2]). In the case where K_X is numerically equivalent to some effective divisor in dimension 5, this conjecture is proved by Birkar (cf. [Bi1]). When X is of general type or K_X is not pseudo-effective, Birkar, Cascini, Hacon

Date: 2011/1/22, version 5.06.

2010 Mathematics Subject Classification. 14E30.

Key words and phrases. minimal model, the abundance conjecture, numerical Kodaira dimension.

and McKernan prove Conjecture 1.1 for arbitrary dimension ([BCHM]). Moreover if X has maximal Albanese dimension, Conjecture 1.1 is true by [F2]. In this article, among other things, we consider Conjecture 1.1 in the case where $\nu(K_X) = 0$ (for the definition of ν , see Definition 2.5):

Theorem 1.2. *Let (X, Δ) be a projective $\mathbb{Q}$ -factorial dlt pair such that $\nu(K_X + \Delta) = 0$. Then there exists a minimal model (X_m, Δ_m) of (X, Δ) .*

Actually the author heard from Vladimir Lazić that Theorem 1.2 for klt pairs has known by Druel (cf. [D]) after he finished this work. In this article, moreover, we give the generalization of his result for dlt pairs by using the sophisticated Birkar–Cascini–Hacon–McKernan’s results and Druel’s way. Essentially our method seems to be same as Druel’s. However, by expanding this result to dlt pairs we give the different proof of the abundance theorem for log canonical pairs in the case where $\nu = 0$ from [CKP] and [Ka2]:

Theorem 1.3 (=Theorem 6.1). *Let X be a normal projective variety and Δ an effective $\mathbb{Q}$ -divisor. Suppose that (X, Δ) is a log canonical pair such that $\nu(K_X + \Delta) = 0$. Then $K_X + \Delta$ is abundant, i.e. $\nu(K_X + \Delta) = \kappa(K_X + \Delta)$.*

First, Nakayama proved Theorem 1.3 when (X, Δ) is klt. Nakayama’s proof is independent of Simpson’s results. Simpson’s results is used to approach the abundance conjecture in [CPT] in the first. Campana–Perternell–Toma prove Theorem 1.3 when X is smooth and $\Delta = 0$. Siu also gave the analytic proof of it (cf. [Siu]). The results of [CKP], [Ka2] and [Siu] depend on [Sim] and [Bu]. In this article, we show Theorem 1.3 by a different way from [CPT], [CKP], [Ka2] and [Siu]. In our proof of Theorem 1.3, we do not need results of [Sim] and [Bu]. Our proof depends on [BCHM] and [G].

We summarize the contents of this article. In Section 2, we define several base loci and the *numerical Kodaira dimension*, and collect some properties of these. In Section 3, we review a *log minimal model program with scaling* and works of Birkar–Cascini–Hacon–McKernan. In Section 4, we introduce the *divisorial Zariski decomposition* and collect some properties of it. Section 5 is devoted to the proof of the existence of minimal models in the case where $\nu = 0$. In Section 6, we prove Theorem 1.3.

Notation and Definition 1.4. Let $\mathbb{K}$ be the real number field $\mathbb{R}$ or the rational number field $\mathbb{Q}$. We set $\mathbb{K}_{>0} = \{x \in \mathbb{K} | x > 0\}$.

Let X be a normal variety and let Δ be an effective $\mathbb{K}$ -divisor such that $K_X + \Delta$ is $\mathbb{K}$ -Cartier. Then we can define the *discrepancy* $a(E, X, \Delta) \in \mathbb{K}$ for every prime divisor E over X . If $a(E, X, \Delta) \geq -1$ (resp. > -1) for every E , then (X, Δ) is called *log canonical* (resp. *kawamata log terminal*). We sometimes abbreviate log canonical (resp. kawamata log terminal) to *lc* (resp. *klt*).

Assume that (X, Δ) is log canonical. If E is a prime divisor over X such that $a(E, X, \Delta) = -1$, then $c_X(E)$ is called a *log canonical center* (*lc center*, for short) of (X, Δ) , where $c_X(E)$ is the closure of the image of E on X . For the basic properties of log canonical centers, see [F1, Section 9].

Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and D a $\mathbb{Z}$ -Cartier divisor on X . We set the complete linear system $|D/S| = \{E|D \sim_{\mathbb{Z}, S} E \geq 0\}$ of D over S . The base locus of the linear system $|D/S|$ is denoted by $\text{Bs}|D/S|$. When $S = \text{Spec } \mathbb{C}$, we denote simply $|D|$ and $\text{Bs}|D|$.

Acknowledgments. The author wishes to express his deep gratitude to Professor Osamu Fujino for various comments and discussions. He would like to thank his supervisor Professor Hiromichi Takagi for carefully reading a preliminary version of this article and valuable comments. He also wishes to thank Professor Caucher Birkar, Professor Noboru Nakayama and Professor Frédéric Campana for valuable comments. Moreover he would like to thank Professor Vyacheslav V. Shokurov for serious remarks about the Kodaira dimensions of $\mathbb{R}$ -divisors and Doctor Vladimir Lazić for teaching him [D]. He is partially supported by the Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists. Lastly, he would like to thank Professor Stéphane Druel, Professor Nobuo Hara, Professor Yujiro Kawamata, Professor Yoichi Miyaoka, Professor Keiji Ogusio and Professor Mihai Păun for the encouragement and comments.

2. PRELIMINARIES

Definition 2.1 (Classical Iitaka dimension, cf. [N, II, 3.2, Definition]). Let X be a normal projective variety and D an $\mathbb{R}$ -Cartier divisor on X . If $|\lfloor mD \rfloor| \neq \emptyset$, we put a dominant rational map

$$\phi_{|\lfloor mD \rfloor|} : X \dashrightarrow W_m,$$

with respect to the complete linear system of $\lfloor mD \rfloor$. We define the *Classical Iitaka dimension* $\kappa(D)$ of D as the following:

$$\kappa(D) = \max\{\dim W_m\}$$

if $H^0(X, \lfloor mD \rfloor) \neq 0$ for some positive integer m or $\kappa(D) = -\infty$ otherwise.

Lemma 2.2. *Let Y be a normal projective variety, $\varphi : Y \rightarrow X$ a projective birational morphism onto a normal projective variety, and let D be an $\mathbb{R}$ -Cartier divisor on X . Then it holds the following:*

- (1) $\kappa(\varphi^*D) = \kappa(\varphi^*D + E)$ for a φ -exceptional effective $\mathbb{R}$ -divisor E , and
- (2) $\kappa(\varphi^*D) = \kappa(D)$.

Proof. (1) and (2) follows from [N, II, 3.11, Lemma]. □

The following is remarked by Shokurov:

Remark 2.3. In general, $\kappa(D)$ may not coincide with $\kappa(D')$ if $D \sim_{\mathbb{R}} D'$. For example, let X be the $\mathbb{P}^1$, P and Q closed points in X such that $P \neq Q$ and a irrational number. Set $D = a(P - Q)$. Then $\kappa(D) = -\infty$ in spite of the fact that $D \sim_{\mathbb{R}} 0$.

However, fortunately, $\kappa(D)$ coincides with $\kappa(D')$ if D and D' are effective divisors such that $D \sim_{\mathbb{R}} D'$ ([Ch, Corollary 2.1.4]). Hence it seems reasonable that we define the following as the *Iitaka (Kodaira) dimension* for $\mathbb{R}$ -divisors.

Definition 2.4 (Invariant Iitaka dimension, [Ch, Definition 2.2.1], cf. [CS, Section 7]). Let X be a normal projective variety and D an $\mathbb{R}$ -Cartier divisor on X . We define the *invariant Iitaka dimension* $\kappa(D)$ of D as the following:

$$K(D) = \kappa(D')$$

if there exists an effective divisor D' such that $D \sim_{\mathbb{R}} D'$ or $K(D) = -\infty$ otherwise. Let (X, Δ) be a log canonical. Then we call $K(K_X + \Delta)$ the *log Kodaira dimension* of (X, Δ) .

Definition 2.5 (Numerical Iitaka dimension). Let X be a normal projective variety, D an $\mathbb{R}$ -Cartier divisor and A an ample Cartier divisor on X . We set

$$\nu(D) = \max\{k \in \mathbb{Z}_{\geq 0} \mid \limsup_{m \rightarrow \infty} m^{-k} \dim H^0(X, \lfloor mD \rfloor + A) > 0\}$$

if $H^0(X, \lfloor mD \rfloor + A) \neq 0$ for infinitely many $m \in \mathbb{N}$ or $\nu(D) = -\infty$ otherwise. Remark that $\nu(D)$ is independent of the choice of φ . Let (X, Δ) be a log pair. Then we call $\nu(K_X + \Delta)$ the *numerical log Kodaira dimension* of (X, Δ) . If $\Delta = 0$, we simply say $\nu(K_X)$ is the *numerical Kodaira dimension* of X .

Lemma 2.6. *Let Y be a normal projective variety, $\varphi : Y \rightarrow X$ a projective birational morphism onto a normal projective variety, and let D be an $\mathbb{R}$ -Cartier divisor on X . Then it holds the following:*

- (1) $\nu(\varphi^*D) = \nu(\varphi^*D + E)$ for a φ -exceptional effective $\mathbb{R}$ -divisor E ,
- (2) $\nu(\varphi^*D) = \nu(D)$ when X is smooth, and
- (3) $\nu(D) = \max\{k \in \mathbb{Z}_{\geq 0} \mid D^k \not\equiv 0\}$ when D is nef.

Proof. See [N, V, 2.7, Proposition]. □

Lemma 2.7 ([N, V, 2.7, Proposition, (1)]). *Let X be a projective variety and D and D' $\mathbb{R}$ -Cartier divisors on X such that $D \equiv D'$. Then $\nu(D) = \nu(D')$.*

Remark 2.8. $\nu(D)$ is denoted as $\kappa_\sigma(D)$ in [N, V, § 2]. Moreover Nakayama also defined $\kappa_\sigma^-(D)$, $\kappa_\sigma^+(D)$ and $\kappa_\nu(D)$ as some numerical Itaka dimensions. In this article we mainly treat in the case where $\nu(D) = 0$, i.e. $\kappa_\sigma(D) = 0$. Then it holds that $\kappa_\sigma^-(D) = \kappa_\sigma^+(D) = \kappa_\nu(D) = 0$ (cf. [N, V, 2.7, Proposition (8)]). Moreover, if a log canonical pair (X, Δ) has a *weakly log canonical model* in the sense of Shokurov, then $\nu(K_X + \Delta)$ coincides with the *numerical log Kodaira dimension* in the sense of Shokurov by Lemma 2.6 (cf. [Sho1, 2.4, Proposition]).

Definition 2.9. Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and D an $\mathbb{R}$ -Cartier divisor on X . We set

$$\mathbf{B}(D/S) = \bigcap_{D \sim_{S, \mathbb{R}} E \geq 0} \text{Supp } E, \quad \mathbf{B}_\equiv(D/S) = \bigcap_{D \equiv_S E \geq 0} \text{Supp } E,$$

and

$$\mathbf{B}_+(D/S) = \bigcap_{\epsilon \in \mathbb{R}_{>0}} \mathbf{B}(D - \epsilon A/S),$$

where A be a π -ample divisor. We remark that these definitions are independent of the choice of A . When $S = \text{Spec } \mathbb{C}$, we denote simply $\mathbf{B}(D)$, $\mathbf{B}_\equiv(D)$ and $\mathbf{B}_+(D)$.

Lemma 2.10. *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and D a π -big $\mathbb{R}$ -Cartier divisor on X . Let $Z_1, Z_2, \dots, Z_k$ be Zariski closed subsets of X . Suppose that $Z_i \not\subseteq \mathbf{B}_+(D/S)$ for all i . Then there exist a π -ample $\mathbb{Q}$ -divisor A and an effective $\mathbb{R}$ -divisor B such that $Z_i \not\subseteq \text{Supp } B$ and*

$$A + B \sim_{\mathbb{R}, S} D.$$

Proof. By [ELMNP, Proposition 1.5], we can take a π -ample $\mathbb{R}$ -divisor A' and a sufficiently small positive number ϵ such that $D - \epsilon A'$ is a $\mathbb{Q}$ -Cartier divisor and

$$\mathbf{B}_+(D/S) = \mathbf{B}(D - \epsilon A'/S).$$

Moreover, from [BCHM, Lemma 3.5.3], it holds that

$$\mathbf{B}(D - \epsilon A'/S) = \bigcap_{m \geq 1} \text{Bs}|md(D - \epsilon A')/S|,$$

where d is a positive integer such that $d(D - \epsilon A')$ is an integral divisor. Therefore we can take $m \gg 0$ such that $Z_i \not\subseteq \text{Bs}|md(D - \epsilon A')/S|$ for all i . Thus a general member $B' \in |md(D - \epsilon A')/S|$ satisfies that $Z_i \not\subseteq \text{Supp } B'$ for any i . We decompose

$$A'' + A \sim_{\mathbb{R}, S} \epsilon A'$$

such that A is a π -ample $\mathbb{Q}$ -divisor and A'' is an effective $\mathbb{R}$ -divisor which satisfies $Z_i \not\subseteq \text{Supp } A''$ for any i . When we put

$$B = \frac{1}{md} B' + A'',$$

Lemma 2.10 follows. $\square$

We introduce a *dlt blow-up*. The following theorem is originally proved by Hacon:

Theorem 2.11 (Dlt blow-up, [F1, Theorem 10.4], [KK, Theorem 3.1]). *Let X be a normal quasi-projective variety and Δ an effective $\mathbb{Q}$ -divisor on X such that $K_X + \Delta$ is $\mathbb{Q}$ -Cartier. Suppose that (X, Δ) is log canonical. Then there exists a projective birational morphism $\varphi : Y \rightarrow X$ from a normal quasi-projective variety with the following properties:*

- (i) Y is $\mathbb{Q}$ -factorial,
- (ii) $a(E, X, \Delta) = -1$ for every φ -exceptional divisor E on Y , and
- (iii) for

$$\Gamma = \varphi_*^{-1} \Delta + \sum_{E: \varphi\text{-exceptional}} E,$$

it holds that (Y, Γ) is dlt and $K_Y + \Gamma = \varphi^(K_X + \Delta)$.*

The above theorem is very useful for studying log canonical singularities (cf. [F1], [G], [KK]).

3. LOG MINIMAL MODEL PROGRAM WITH SCALING

In this section, we review a *log minimal model program with scaling* and introduce works by Birkar–Cascini–Hacon–McKernan.

Lemma 3.1 (cf. [Bi1, Lemma 2.1] and [F1, Theorem 18.9]). *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and (X, Δ) a $\mathbb{Q}$ -factorial projective log canonical pair such that Δ is a $\mathbb{Q}$ -divisor. Let H be an effective $\mathbb{Q}$ -divisor such that $K_X + \Delta + H$ is π -nef and $(X, \Delta + H)$ is log canonical. Suppose that $K_X + \Delta$ is not π -nef. We put*

$$\lambda = \inf\{\alpha \in \mathbb{R} \mid K_X + \Delta + \alpha H \text{ is } \pi\text{-nef}\}.$$

Then $\lambda \in \mathbb{K}_{>0}$ and there exists an extremal ray $R \subseteq \overline{NE}(X/S)$ such that $(K_X + \Delta).R < 0$ and $(K_X + \Delta + \lambda H).R = 0$.

Definition 3.2 (Log minimal model program with scaling). Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and (X, Δ) a $\mathbb{Q}$ -factorial projective divisorial log terminal pair such that Δ is a $\mathbb{K}$ -divisor. Let H be an effective $\mathbb{K}$ -divisor such that $K_X + \Delta + H$ is π -nef and $(X, \Delta + H)$ is divisorial log terminal. We put

$$\lambda_1 = \inf\{\alpha \in \mathbb{R} \mid K_X + \Delta + \alpha H \text{ is } \pi\text{-nef}\}.$$

If $K_X + \Delta$ is not π -nef, then $\lambda_1 > 0$. By Lemma 3.1, there exists an extremal ray $R_1 \subseteq \overline{NE}(X/S)$ such that $(K_X + \Delta).R_1 < 0$ and $(K_X + \Delta + \lambda_1 H).R_1 = 0$. We consider an extremal contraction with respect to this R_1 . If it is a divisorial contraction or a flipping contraction, let

$$(X, \Delta) \dashrightarrow (X_1, \Delta_1)$$

be the divisorial contraction or its flip. Since $K_{X_1} + \Delta_1 + \lambda_1 H_1$ is π -nef, we put

$$\lambda_2 = \inf\{\alpha \in \mathbb{R} \mid K_{X_1} + \Delta_1 + \alpha H_1 \text{ is } \pi\text{-nef}\},$$

where H_1 is the strict transform of H on X_1 . Then we find an extremal ray R_2 by the same way as the above. We may repeat the process. We call this program a *log minimal model program with scaling of H* over S . When this program runs as the following:

$$(X_0, \Delta_0) = (X, \Delta) \dashrightarrow (X_1, \Delta_1) \dashrightarrow \cdots \dashrightarrow (X_i, \Delta_i) \cdots,$$

then

$$\lambda_1 \geq \lambda_2 \geq \lambda_3 \cdots,$$

where $\lambda_i = \inf\{\alpha \in \mathbb{R} \mid K_{X_{i-1}} + \Delta_{i-1} + \alpha H_{i-1} \text{ is } \pi\text{-nef}\}$ and H_{i-1} is the strict transform of H on X_{i-1} .

The following theorems are slight generalizations of [BCHM, Corollary 1.4.1] and [BCHM, Corollary 1.4.2]. These seem to be well-known for the experts.

Theorem 3.3 (cf. [BCHM, Corollary 1.4.1]). *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and (X, Δ) be a $\mathbb{Q}$ -factorial projective divisorial log terminal pair such that Δ is an $\mathbb{R}$ -divisor. Suppose that $\varphi : X \rightarrow Y$ is a flipping contraction of (X, Δ) . Then there exists the log flip of φ .*

Proof. Since $-(K_X + \Delta)$ is φ -ample, so is $-(K_X + \Delta - \epsilon_L \Delta_L)$ for a sufficiently small $\epsilon > 0$. Because $\rho(X/Y) = 1$, it holds that $K_X + \Delta \sim_{\mathbb{R}, Y} c(K_X + \Delta - \epsilon_L \Delta_L)$ for some positive number c . By [BCHM, Corollary 1.4.1], there exists the log flip of $(X, \Delta - \epsilon_L \Delta_L)$. This log flip is also the log flip of (X, Δ) since $K_X + \Delta \sim_{\mathbb{R}, Y} c(K_X + \Delta - \epsilon_L \Delta_L)$. $\square$

Theorem 3.4 (cf. [BCHM, Corollary 1.4.2]). *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and (X, Δ) be a $\mathbb{Q}$ -factorial projective divisorial log terminal pair such that Δ is an $\mathbb{R}$ -divisor. Suppose that there exists a π -ample $\mathbb{R}$ -divisor A on X such that $\Delta \geq A$. Then any sequences of log flips starting from (X, Δ) with scaling of H over S terminate, where H satisfies that $(X, \Delta + H)$ is divisorial log terminal and $K_X + \Delta + H$ is π -nef.*

The above theorem is proved by the same argument as the proof of [BCHM, Corollary 1.4.2] because [BCHM, Theorem E] holds on the above setting.

4. DIVISORIAL ZARISKI DECOMPOSITION

In this section, we introduce the *divisorial Zariski decomposition* for a pseudo-effective divisor.

Definition 4.1 (cf. [N, III, 1.13, Definition] and [Ka1]). Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and D an $\mathbb{R}$ -Cartier divisor. We call that D is a *limit of movable $\mathbb{R}$ -divisors* over S if $[D] \in \overline{\text{Mov}}(X/S) \subseteq N^1(X/S)$ where $\overline{\text{Mov}}(X/S)$ is the closure of the convex cone spanned by classes of fixed part free $\mathbb{Z}$ -Cartier divisors over S . When $S = \text{Spec } \mathbb{C}$, we denote simply $\overline{\text{Mov}}(X)$.

Lemma 4.2. *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and D an $\mathbb{R}$ -Cartier divisor. Suppose that $\mathbf{B}(D/S)$ does not coincide with X and has no codimension 1 components. Then there exist fixed part free $\mathbb{Z}$ -Cartier divisors $\{Z_i\}$ over S and positive*

numbers $\{\alpha_i\}$ such that

$$[D] = \sum_i \alpha_i [Z_i] \in N^1(X/S).$$

Proof. This follows from [BCHM, Proposition 3.5.4]. $\square$

Definition 4.3 (cf. [N, III, 1.6, Definition and 1.12, Definition]). Let X be a smooth projective variety and B an $\mathbb{R}$ -big divisor. We define

$$\sigma_\Gamma(B) = \inf\{\text{mult}_\Gamma B' \mid B \equiv B' \geq 0\}$$

for a prime divisor Γ . Let D be a pseudo-effective divisor. Then we define the following:

$$\sigma_\Gamma(D) = \lim_{\epsilon \rightarrow 0+} \sigma_\Gamma(D + \epsilon A)$$

for some ample divisor A . We remark that $\sigma_\Gamma(D)$ is independent of the choice of A . Moreover the above two definitions coincide for a big divisor because a function $\sigma_\Gamma(\cdot)$ on $\text{Big}(X)$ is continuous where $\text{Big}(X) := \{[B] \in N^1(X) \mid B \text{ is big}\}$ (cf. [N, III, 1.7, Lemma]). We set

$$N(D) = \sum_{\Gamma: \text{prime divisor}} \sigma_\Gamma(D) \Gamma \text{ and } P(D) = D - N(D).$$

We remark that $N(D)$ is a finite sum. We call the decomposition $D \equiv P(D) + N(D)$ the *divisorial Zariski decomposition* of D . We say that $P(D)$ (resp. $N(D)$) is the *positive part* (resp. *negative part*) of D .

Remark that the decomposition $D \equiv P(D) + N(D)$ is called several names: the *sectional decomposition* ([Ka1]), the σ -*decomposition* ([N]), the *divisorial Zariski decomposition* ([Bo]), and the *numerical Zariski decomposition* ([Ka2]).

Proposition 4.4. *Let X be a smooth projective variety and D a pseudo-effective $\mathbb{R}$ -divisor on X . Then it holds the following:*

- (1) $\sigma_\Gamma(D) = \lim_{\epsilon \rightarrow 0+} \sigma_\Gamma(D + \epsilon E)$ for a pseudo-effective divisor E ,
and
- (2) $\nu(D) = 0$ if and only if $D \equiv N(D)$.

Proof. (1) follows from [N, III, 1.4, Lemma]. (2) follows from [N, V, 2.7, Proposition (8)]. $\square$

5. EXISTENCE OF MINIMAL MODEL IN THE CASE WHERE $\nu = 0$

Lemma 5.1 (cf. [BCHM, Lemma 3.7.3]). *Let $\pi : X \rightarrow S$ be a projective morphism of normal quasi-projective varieties and (X, Δ) a $\mathbb{Q}$ -factorial projective divisorial log terminal pair such that Δ is an $\mathbb{R}$ -divisor, and H an effective π -big $\mathbb{R}$ -divisor such that $(X, \Delta + H)$ is*

divisorial log terminal. Suppose that no lc centers of (X, Δ) are contained in $\mathbf{B}_+(H/S)$. Then there exist an effective π -ample $\mathbb{Q}$ -divisor A and an effective $\mathbb{R}$ -divisor B such that $H \sim_{\mathbb{R}, S} B + A$ and $(X, \Delta + B + A)$ is divisorial log terminal.

Proof. Since no lc centers of (X, Δ) are contained in $\mathbf{B}_+(H/S)$, there exist a π -ample $\mathbb{Q}$ -divisor A' and an effective $\mathbb{R}$ -divisor B' such that $\text{Supp } B'$ contains no lc centers of (X, Δ) and

$$A' + B' \sim_{\mathbb{R}, S} H$$

from Lemma 2.10. We take a general effective $\mathbb{Q}$ -divisor $A'' \sim_{\mathbb{Q}, S} A'$ and a sufficiently small number $\delta \in \mathbb{Q}_{>0}$ such that $(X, \Delta + (1 - \delta)H + \delta(A'' + B'))$ is dlt. When we put

$$A = \delta A'' \text{ and } B = (1 - \delta)H + \delta B',$$

Lemma 5.1 holds. □

Theorem 5.2 (cf. [D, Corollaire 3.4]). *Let X be a $\mathbb{Q}$ -factorial projective variety and Δ an effective $\mathbb{R}$ -divisor such that (X, Δ) is divisorial log terminal. Suppose that $\nu(K_X + \Delta) = 0$. Then any log minimal model programs starting from (X, Δ) with scaling of H terminate, where H satisfies that $H \geq A$ for some effective $\mathbb{Q}$ -ample divisor A and $(X, \Delta + H)$ is divisorial log terminal and $K_X + \Delta + H$ is nef.*

Proof. Let $(X, \Delta) \dashrightarrow (X_1, \Delta_1)$ be a divisorial contraction or a log flip. Remark that it holds that

$$\nu(K_X + \Delta) = \nu(K_{X_1} + \Delta_1)$$

from Lemma 2.6 (1) and the negativity lemma. If some lc centers of (X, Δ) are contained in $\text{Supp}(H - A)$, then $(X, \Delta + H)$ is not log canonical. Hence, by the assumption, no lc centers of (X, Δ) are contained in $\mathbf{B}_+(H)$. By the negativity lemma, no lc centers of (X_1, Δ_1) are contained in $\mathbf{B}_+(H_1)$, where H_1 is the strict transform of H on X_1 (cf. [BCHM, Lemma 3.10.11]). From Lemma 5.1, there exist an effective $\mathbb{Q}$ -ample divisor A_1 and an effective $\mathbb{R}$ -divisor B_1 such that $H_1 \sim_{\mathbb{R}} B_1 + A_1$, $(X_1, \Delta_1 + B_1 + A_1)$ is divisorial log terminal and $K_{X_1} + \Delta_1 + A_1 + B_1$ is nef. We may replace (X, Δ, H) with $(X_1, \Delta_1, A_1 + B_1)$. Therefore it suffices to show termination of such log flips. Assume that there exists a sequence of log flips

$$(X_i, \Delta_i) \dashrightarrow (X_{i+1}, \Delta_{i+1})$$

with scaling of H does not terminate where $(X_0, \Delta_0) = (X, \Delta)$. Let $\{\lambda_i\}$ be as in Definition 3.2. We set

$$\lambda = \lim_{i \rightarrow \infty} \lambda_i.$$

If $\lambda \neq 0$, the sequence is composed by $(K_X + \Delta + \frac{1}{2}\lambda H)$ -log flips. Thus the sequence terminates by Theorem 3.4. We may assume that $\lambda = 0$. Then we see the following:

Claim 5.3. $K_X + \Delta$ is a limit of movable $\mathbb{R}$ -divisors.

Proof of Claim 5.3. Let $(X, \Delta) = (X_0, \Delta_0) \dashrightarrow (X_i, \Delta_i)$ be a composition of the log flips. We take a log resolution $p_i : W_i \rightarrow X$ of $(X, \Delta + H)$ and a log resolution $q_i : W_i \rightarrow X_i$ of $(X_i, \Delta_i + H_i)$ such that the following diagram commutes, where H_i is the strict transform of H on X_i :

$$\begin{array}{ccc} & W_i & \\ p_i \swarrow & & \searrow q_i \\ X & \cdots \cdots \cdots & X_i. \end{array}$$

Let G_i be an ample divisor on X_i and ϵ_i a sufficiently small rational number. Since $K_{X_i} + \Delta_i + \lambda_i H_i$ is nef, we see that $K_{X_i} + \Delta_i + \lambda_i H_i + \epsilon_i G_i$ is ample. By the negativity lemma, it holds that

$$p_i^*(K_X + \Delta + \lambda_i H + \epsilon_i G_{i,X}) = q_i^*(K_{X_i} + \Delta_i + \lambda_i H_i + \epsilon_i G_i) + E_i$$

where $G_{i,X}$ is the strict transform of G_i on X and E_i is an effective p_i -exceptional and q_i -exceptional divisor. Thus it holds that

$$\mathbf{B}(p_i^*(K_X + \Delta + \lambda_i H + \epsilon_i G_{i,X})) = \text{Supp } E_i.$$

Since $\mathbf{B}(K_X + \Delta + \lambda_i H + \epsilon_i G_{i,X}) \subseteq p_i(\text{Supp } E_i)$, we see that $K_X + \Delta + \lambda_i H + \epsilon_i G_{i,X}$ is movable from Lemma 4.2. Hence, if we take $\{\epsilon_i\}$ such that $\epsilon_i \rightarrow 0$, Claim 5.3 follows. $\square$

Let $\varphi : Y \rightarrow X$ be a log resolution of (X, Δ) . We consider the σ -decomposition

$$\varphi^*(K_X + \Delta) \equiv P(\varphi^*(K_X + \Delta)) + N(\varphi^*(K_X + \Delta))$$

(Definition 4.3). Since

$$\nu(\varphi^*(K_X + \Delta)) = \nu(K_X + \Delta) = 0,$$

we see $P(\varphi^*(K_X + \Delta)) = 0$ by Proposition 4.4 (2). In particular, this is the divisorial Zariski decomposition. Moreover we see the following claim:

Claim 5.4. $N(\varphi^*(K_X + \Delta))$ is a φ -exceptional divisor.

Proof of Claim 5.4. Let G be an ample divisor on X and ϵ a sufficiently small positive number. By Proposition 4.4 (1), it holds that

$$\text{Supp } N(\varphi^*(K_X + \Delta)) \subseteq \text{Supp } N(\varphi^*(K_X + \Delta + \epsilon G)).$$

If it holds that $\varphi_*(N(\varphi^*(K_X + \Delta))) \neq 0$, we see that $\mathbf{B}_\equiv(K_X + \Delta + \epsilon G)$ has codimension 1 components. This is a contradiction to Claim 5.3. Thus $N(\varphi^*(K_X + \Delta))$ is a φ -exceptional divisor. $\square$

Hence $K_X + \Delta \equiv 0$, in particular, $K_X + \Delta$ is nef. This is a contradiction to the assumption. $\square$

Corollary 5.5. *Let X be a $\mathbb{Q}$ -factorial projective variety and Δ an effective $\mathbb{R}$ -divisor such that (X, Δ) is divisorial log terminal. Suppose that $\nu(K_X + \Delta) = 0$. Then there exists a log minimal model of (X, Δ) .*

Remark 5.6. These results are on the absolute setting. It may be difficult to extend these to the relative settings. See [F3].

6. ABUNDANCE THEOREM IN THE CASE WHERE $\nu = 0$

In this section, we prove the abundance theorem in the case where $\nu = 0$ for an $\mathbb{R}$ -divisor:

Theorem 6.1. *Let X be a normal projective variety and Δ an effective $\mathbb{R}$ -divisor. Suppose that (X, Δ) is a log canonical pair such that $\nu(K_X + \Delta) = 0$. Then $\nu(K_X + \Delta) = K(K_X + \Delta)$. Moreover, if Δ is a $\mathbb{Q}$ -divisor, then $\nu(K_X + \Delta) = \kappa(K_X + \Delta) = K(K_X + \Delta)$.*

First we extend [G, Theorem 1.2] to an $\mathbb{R}$ -divisor.

Lemma 6.2 (cf. [FG, Theorem 3.1]). *Let X be a normal projective variety and Δ an effective $\mathbb{K}$ -divisor. Suppose that (X, Δ) is a log canonical pair such that $K_X + \Delta \equiv 0$. Then $K_X + \Delta \sim_{\mathbb{K}} 0$.*

Proof. If $\mathbb{K} = \mathbb{Q}$, then the statement is nothing but [G, Theorem 1.2]. From now on, we assume that $\mathbb{K} = \mathbb{R}$. Let $\sum_i B_i$ be the irreducible decomposition of $\text{Supp } \Delta$. We put $V = \bigoplus_i \mathbb{R}B_i$. Then it is well known that

$$\mathcal{L} = \{\Delta \in V \mid (X, \Delta) \text{ is log canonical}\}$$

is a rational polytope in V . We can also check that

$$\mathcal{N} = \{\Delta \in \mathcal{L} \mid K_X + \Delta \text{ is nef}\}$$

is a rational polytope and $\Delta \in \mathcal{N}$ (cf. [Bi2, Proposition 3.2] and [Sho1, 6.2 First Main theorem]). We note that $\mathcal{N}$ is known as Shokurov's polytope. Therefore, we can write

$$K_X + \Delta = \sum_{i=1}^k r_i (K_X + \Delta_i)$$

such that

- (i) Δ_i is an effective $\mathbb{Q}$ -divisor such that $\Delta_i \in \mathcal{N}$ for every i ,
- (ii) (X, Δ_i) is log canonical for every i , and
- (iii) $0 < r_i < 1$, $r_i \in \mathbb{R}$ for every i , and $\sum_{i=1}^k r_i = 1$.

Since $K_X + \Delta$ is numerically trivial and $K_X + \Delta_i$ is nef for every i , $K_X + \Delta_i$ is numerically trivial for every i . By [G, Theorem 1.2], we see that $K_X + \Delta \sim_{\mathbb{R}} 0$. □

Proof of Theorem 6.1. By taking a dlt blow-up (Theorem 2.11), we may assume that (X, Δ) is a $\mathbb{Q}$ -factorial dlt pair. By Corollary 5.5, there exists a log minimal model (X_m, Δ_m) of (X, Δ) . From Lemma 2.6 (3), it holds that $K_{X_m} + \Delta_m \equiv 0$. By Lemma 6.2, it holds that $\kappa(K_{X_m} + \Delta_m) = 0$. Lemma 2.2 implies that $K(K_X + \Delta) = 0$. If Δ is a $\mathbb{Q}$ -divisor, then there exists an effective $\mathbb{Q}$ -divisor E such that $K_X + \Delta \sim_{\mathbb{Q}} E$ by Corollary 5.5 and Lemma 6.2. Thus we see that $\kappa(K_X + \Delta) = K(K_X + \Delta)$. We finish the proof of Theorem 6.1. □

Corollary 6.3. *Let $\pi : X \rightarrow S$ be a projective surjective morphism of normal quasi-projective varieties, and let (X, Δ) be a projective log canonical pair such that Δ is an effective $\mathbb{K}$ -divisor. Suppose that $\nu(K_F + \Delta_F) = 0$ for a general fiber F , where $K_F + \Delta_F = (K_X + \Delta)|_F$. Then there exists an effective $\mathbb{K}$ -divisor D such that $K_X + \Delta \sim_{\mathbb{K}, \pi} D$.*

Proof. This follows from Theorem 6.1 and [BCHM, Lemma 3.2.1]. □

REFERENCES

- [Bi1] C. Birkar, On existence of log minimal models, *Compos. Math.* 146 (2010), no. 4, 919-928.
- [Bi2] ———, On existence of log minimal models II, preprint, arXiv:0907.4170, to appear in *J. Reine Angew. Math.*
- [BCHM] C. Birkar, P. Cascini, C. D. Hacon and J. McKernan, Existence of minimal models for varieties of log general type, *J. Amer. Math. Soc.* 23 (2010), 405-468.
- [Bo] S. Boucksom, Divisorial Zariski decomposition on compact complex manifolds, *Ann. Sci. Ecole Norm. Sup.* 37 (2004), no.4, 45-76.

- [Bu] N. Budur, Unitary local systems, multiplier ideals, and polynomial periodicity of Hodge numbers, *Adv. Math.* 221 (2009), no. 1, 217–250.
- [CKP] F. Campana, V. Koziarz and M. Păun, Numerical character of the effectivity of adjoint line bundles, preprint, arXiv:1004.0584, to appear in *Ann. Inst. Fourier*.
- [CPT] F. Campana, T. Peternell and M. Toma, Geometric stability of the cotangent bundle and the universal cover of a projective manifold, preprint, math/0405093.
- [Ch] S. R. Choi, The geography of log models and its applications, Ph.D thesis. Johns Hophkin University, 2008.
- [CS] S. R. Choi and V. V. Shokurov, The geography of log models and its applications, arXiv:0909.0288.
- [D] S. Druel, Quelques remarques sur la décomposition de Zariski divisorielle sur les variétés dont la première classe de Chern est nulle, to appear in *Math. Z.*
- [ELMNP] L. Ein, R. Lazarsfeld, M. Mustață, M. Nakamaye, M. Popa, Asymptotic invariants of base loci, *Ann. Inst. Fourier (Grenoble)* 56 (2006), no. 6, 1701–1734.
- [F1] O. Fujino, Fundamental theorems for the log minimal model program, preprint, to appear in *Publ. Res. Inst. Math. Sci.*
- [F2] ———, On maximal Albanese dimensional varieties. preprint, arXiv:0911.2851.
- [F3] ———, Semi-stable minimal model program for varieties with trivial canonical divisor, preprint, arXiv:1010.2577.
- [FG] O. Fujino and Y. Gongyo, On canonical bundle formulae and subadjunctions, preprint, arXiv:1009.3996.
- [G] Y. Gongyo, Abundance theorem for numerically trivial log canonical divisors of semi-log canonical pairs, preprint, arXiv:1005.2796v2.
- [Ka1] Y. Kawamata, Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces. *Ann. of Math.* 127(1988), 93–163.
- [Ka2] Y. Kawamata, On the abundance theorem in the case of $\nu = 0$, preprint, arXiv:1002.2682.
- [KMM] Y. Kawamata, K. Matsuda and K. Matsuki, *Introduction to the minimal model problem*, Algebraic geometry, Sendai, 1985, 283–360, *Adv. Stud. Pure Math.*, 10, North-Holland, Amsterdam, 1987.
- [KK] J. Kollár and S. J. Kovács, Log canonical singularities are Du Bois, preprint, arXiv:0902.0648.
- [KM] J. Kollár and S. Mori. *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., 134 (1998).
- [N] N. Nakayama, *Zariski decomposition and abundance*, MSJ Memoirs, 14. Mathematical Society of Japan, Tokyo, 2004.
- [Sho1] V. V. Shokurov, 3-fold log models, *Algebraic geometry*, 4. *J. Math. Sci.* 81 (1996), no. 3, 2667–2699.
- [Sho2] ———, Prelimiting flips, *Tr. Mat. Inst. Steklova* 240 (2003), Biratsion. Geom. Linein. Sist. Konechno Porozhdennye Algebry, 82–219; translation in *Proc. Steklov Inst. Math.* 2003, no. 1 (240), 75–213.

- [Sim] C. Simpson, Subspaces of moduli spaces of rank one local systems, Ann. Sci. École Norm. Sup. (4) 26 (1993), no. 3, 361–401.
- [Siu] Y. T. Siu, Abundance conjecture, preprint, arXiv:0912.0576v3.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,
3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914, JAPAN.

E-mail address: gongyo@ms.u-tokyo.ac.jp