

In this talk we define the pro-étale site of a scheme, discuss the case of a field in detail, see that in general the pro-étale topos is replete.

1 The pro-étale site


Recall that a morphism of schemes $f : Y \rightarrow X$ of finite presentation is étale, if it is flat and the diagonal $Y \rightarrow Y \times_X Y$ is flat.¹

Definition 1 (Def.4.1.1). *A map $f : Y \rightarrow X$ of schemes is weakly étale if it is flat, and the diagonal $\Delta : Y \rightarrow Y \times_X Y$ is flat. The category of weakly étale X -schemes is denoted $X_{\text{proét}}$.*

Example 2.

1. Suppose $A \rightarrow B$ is an ind-étale morphism of rings (so $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is pro-étale). Then we saw previously that $A \rightarrow B$ is a weakly étale morphism of rings [Prop.2.3.3], so $\text{Spec}(B) \rightarrow \text{Spec}(A)$ is a weakly étale morphism of schemes.
2. Bhatt, Scholze choose to work with weakly étale maps instead of pro-étale morphisms of schemes in general because pro-étale morphisms of schemes are not so well-behaved, cf. [Exa.4.1.12] reproduced below.
3. [Exa.4.1.9] Given a scheme X , and a profinite set $S = \varprojlim_i S_i$, the morphism $X \otimes S := \varprojlim_{i \in I} (\sqcup_{s \in S_i} X) \rightarrow X$ is pro-étale. This defines a functor

$$\text{ProFinSet} \times X_{\text{proét}} \rightarrow X_{\text{proét}}; \quad (S, Y) \mapsto S \otimes Y$$




4. [Exa.4.1.12] Consider the set²

$$S = \{e^{\pi i(1-\frac{1}{2^n})} : n \in \mathbb{Z}, n \geq 0\} \cup \{e^{\pi i(2^n-1)} : n \in \mathbb{Z}, n \leq 0\} \cup \{-1\} \subseteq \mathbb{C}$$


¹The diagonal being flat is one of a number of equivalent definitions for a finite presentation morphism to be unramified.

²I.e., the one point compactification of \mathbb{Z} considered as a discrete set.


equipped with the translation function T induced by $n \mapsto n + 1$ on the image of \mathbb{Z} , and sending $-1 \in \mathbb{C}$ to -1 .




Note that this is a profinite set



Now let $X_1, X_2 \subseteq \mathbb{A}_{\mathbb{C}}^2$ be two smooth curves meeting transversally at points p and q , and $X = X_1 \cup X_2$.



Then consider the X -scheme Y which is $S \otimes X_1$ glued to $S \otimes X_2$ using the identity at p and the translation function T at q .



Then $Y \rightarrow X$ is locally pro-étale. Indeed, away from p (or q), it is just $S \otimes (X \setminus \{p\}) \rightarrow (X \setminus \{p\})$. However, $Y \rightarrow X$ is not globally pro-étale. By closely considering the topology on Y , one can see that it cannot be written as $Y = \varprojlim Y_i$ for étale X schemes Y_i .

5. [Exa.4.1.4] If k is a field, then a morphism $\text{Spec}(R) \rightarrow \text{Spec}(k)$ is weakly étale if and only if $k \rightarrow R$ is ind-étale.³
6. [Exa.4.1.5] For any scheme X , point $x \in X$, and geometric point $\bar{x} \rightarrow X$, the morphisms

$$\text{Spec}(\mathcal{O}_{X,x}) \rightarrow X, \quad \text{Spec}(\mathcal{O}_{X,x}^h) \rightarrow X, \quad \text{Spec}(\mathcal{O}_{X,\bar{x}}^h) \rightarrow X$$

are all weakly étale.

Recall that flat morphisms are preserved by base change and composition.

Exercise 1 ([Lem.4.1.6]). *Weakly étale morphisms are preserved by base change.* Show that if $f : Y \rightarrow X$ is weakly étale then $X' \times_X Y \rightarrow X'$ is weakly étale for any morphism $X' \rightarrow X$.


Exercise 2 ([Lem.4.1.6], [Lem.4.1.7]). *Weakly étale morphisms are preserved by composition, and all morphisms in $X_{\text{proét}}$ are weakly étale.* Let $g : W \rightarrow Y$ and $f : Y \rightarrow X$, $f' : Y' \rightarrow X$ be weakly étale morphisms, and $h : Y' \rightarrow Y$ any X -morphism.

³By [BS, Thm.2.3.4], for every weakly étale morphism $A \rightarrow B$ there is a faithfully flat ind-étale morphism $B \rightarrow C$ such that $A \rightarrow C$ is ind-étale. In particular, B is a sub- A -algebra of an ind-étale A -algebra. But for fields k , every sub- k -algebra B of an ind-étale k -algebra C is again an ind-étale k -algebra: Indeed, write $C = \varinjlim_{i \in I} C_i$ where the C_i are étale k -algebras, and recall that this means that each C_i is a finite product of finite separable k -field extensions. Replacing each C_i with its image in C , we can assume all morphisms $C_i \rightarrow C$ are injective. Then taking $B_i = C_i \cap B$, we produce a system $(B_i)_{i \in I}$ such that each B_i is an étale k -algebra and $B = \varinjlim_{i \in I} B_i$.

1. Use the fact that $W \cong (Y \times_X W) \times_{(Y \times_X Y)} Y$ to show that $W \rightarrow Y \times_X W$ is flat.
2. Use part (1), the fact that $Y \times_X W \times_X W \cong (Y \times_X W) \times_Y (Y \times_X W)$, and a clever factorisation of $W \rightarrow W \times_X W$ to show that $f \circ g : W \rightarrow X$ is weakly étale.
3. As in part (1), show that $Y' \rightarrow Y \times_X Y'$ is flat.
4. Use part (3) and the fact that $Y' \cong (Y' \times_Y Y') \times_{(Y' \times_X Y')} (Y')$ to show that $Y' \rightarrow Y$ is pro-étale.

Exercise 3 ([Lem.4.1.8]). Use Exercise 1 and Exercise 2 to show that $X_{\text{proét}}$ has fibre products. Deduce that $X_{\text{proét}}$ has all finite limits. (Recall, that an exercise earlier in the course was to show that a category has all finite limits if and only if it has fibre products and a terminal object).

Definition 3. A family $\{Y_i \rightarrow Y\}_{i \in I}$ in $X_{\text{proét}}$ is a covering, if for every open affine $U \subseteq Y$, there is a finite subset $J \subseteq I$ and open affines $V_j \subseteq Y_j$ for each $j \in J$ such that $\coprod_{j \in J} V_j \rightarrow U$ is surjective.



The finiteness in the above definition is important, and affects the topology:


Example 4 ([Exa.4.1.13]). Consider $\text{Spec}(\mathbb{Z})$. If p_1, \dots, p_n are finitely many primes, then

$$\left\{ \text{Spec}(\mathbb{Z}_{(p_1)}^{sh}), \dots, \text{Spec}(\mathbb{Z}_{(p_n)}^{sh}), \text{Spec}(\mathbb{Z}[\frac{1}{p_1}, \dots, \frac{1}{p_n}]) \right\}$$

is a weakly étale cover. However,

$$\left\{ \text{Spec}(\mathbb{Z}_{(p)}^{sh}) : p \text{ is prime} \right\}$$

is *not* a weakly étale cover.



2 The pro-étale site of a field

Example 5. Suppose that k is a separably closed field, and R is an ind-étale algebra

$$k \rightarrow R = \varinjlim R_\lambda.$$

So each R_λ is a *finite* product $R_\lambda = \prod_{i \in I_\lambda} k$. Moreover, every morphism $\prod_{i \in I_\lambda} k = R_\lambda \rightarrow R'_\lambda = \prod_{j \in I'_\lambda} k$ is induced by morphisms of sets $\phi_{\lambda, \lambda'} : I'_\lambda \rightarrow I_\lambda$. Since the underlying topological space of Spec of a filtered colimit of rings is the inverse limit of the underlying topological spaces, [EGAIV, §8], we see that the underlying topological space of $\text{Spec}(R)$ is the profinite set $I = \varprojlim I_\lambda$.

$$\text{Spec}(R)_{\text{top}} = I.$$

For any profinite set $\varprojlim S_\lambda$, and any discrete set X , one can see that we have⁴ $\text{hom}_{\text{cont.}}(\varprojlim S_\lambda, X) = \varinjlim \text{hom}(S_\lambda, X)$. Hence,

$$\Gamma(\text{Spec}(R), \mathcal{O}_{\text{Spec}(R)}) = \text{hom}_{\text{cont.}}(I, k)$$

is the set of continuous morphisms where k is given the discrete topology. Moreover, for any open subset of the form $U_{\lambda, i} = \phi_\lambda^{-1}(i)$ where $i \in I_\lambda$ and $\phi_\lambda : I \rightarrow I_\lambda$ is the canonical projection, $U_{\lambda, i}$ is again ind-étale, and so $\Gamma(U_{\lambda, i}, \mathcal{O}_{\text{Spec}(R)}) = \text{hom}_{\text{cont.}}(U_{\lambda, i}, k)$. Finally, if $U \subseteq I$ is any open subset, and $\{U_i \subset U\}$ an open covering, we have

$$\Gamma(U, \mathcal{O}_{\text{Spec}(R)}) = \text{Eq} \left(\prod_{i \in I} \Gamma(U_i, \mathcal{O}_{\text{Spec}(R)}) \rightrightarrows \prod_{i, j \in I} \Gamma(U_i \cap U_j, \mathcal{O}_{\text{Spec}(R)}) \right)$$

⁴ If $f : \varprojlim S_\lambda \rightarrow X$ is a continuous morphism, then for every point $x \in X$, $f^{-1}x$ is open, and by definition of the limit topology, admits a covering of the form $\mathcal{U}_x = \{U_{\lambda, s} : \lambda \in \Lambda_x, s \in S_\lambda\}$ where $U_{\lambda, s} = \phi_\lambda^{-1}(s)$ is the preimage of s under the canonical projection $\phi : S \rightarrow S_\lambda$. Since S is profinite, it is quasicompact, so the covering family $\cup_{x \in X} \mathcal{U}_x$ admits a finite subcovering $\{V_{\lambda_i, s_i} : 1 \leq i \leq n, s_i \in S_{\lambda_i}\}$. Then by construction any λ with $\lambda \leq \lambda_1, \dots, \lambda_n$, has the property that f is constant on the fibres of $S \rightarrow S_\lambda$. Hence, f factors as $S \rightarrow S_\lambda \rightarrow X$.

By definition, every open of I is covered by opens of the form $U_{\lambda,i}$, so we deduce that in general,

$$\Gamma(U, \mathcal{O}_{\text{Spec}(R)}) = \text{hom}_{\text{cont.}}(U, k).$$

Given any point $x \in I$, and any open $x \in U$, there is a smaller open containing x of the form $U_{\lambda,i}$. For any continuous function $f : U \rightarrow k$, there is a refinement $x \in U_{\lambda,i} \subseteq U$. But $U_{\lambda,i}$ is profinite, and therefore quasicompact, so the image $f(U_{\lambda,i})$ is finite, so there is a further refinement $x \in V \subseteq U_{\lambda,i}$ such that $f : V \rightarrow k$ is constant. It follows that all local rings of $\text{Spec}(R)$ are isomorphic to k .

$$\mathcal{O}_{\text{Spec}(R),x} \cong k.$$

Conversely, if (X, \mathcal{O}_X) is a locally ringed space such that $X = \varprojlim X_\lambda$ is profinite, and $\mathcal{O}_X(U) = \text{hom}_{\text{cont.}}(U, k)$ for some separable closed field k , then $(X, \mathcal{O}_X) \cong \text{Spec}(\varinjlim \prod_{X_\lambda} k)$, i.e., (X, \mathcal{O}_X) is the affine scheme associated to an ind-étale k -algebra.

Proposition 6. *Suppose k is a separable closed field and $X \in \text{Spec}(k)_{\text{proét}}$. The following are equivalent.*

1. X is affine.
2. X is the spectrum of an ind-étale algebra.
3. X is qcqs.
4. X is of the form $\text{Spec}(k) \otimes S$ for a profinite set S .

Proof. (1 \iff 2) We have seen, Exa.2(5) that the affine schemes in $\text{Spec}(k)_{\text{proét}}$ are precisely the spectra of ind-étale k -algebras.

(2 \iff 3) All affine schemes are qcqs, so consider the other direction. Suppose that X is qcqs. A scheme is qcqs if and only if it admits a finite open affine cover $\{U_i \rightarrow X\}_{i=1}^n$ such that each $U_i \cap U_j$ for $1 \leq i, j \leq n$ is also affine. Since affines in $\text{Spec}(k)_{\text{proét}}$ have profinite underlying topological space (i.e., compact, Hausdorff, totally disconnected topological space), it follows that any qcqs X also has profinite underlying topological space (see the lemma below). Moreover, the structure sheaf of X has the form $V \mapsto \text{hom}_{\text{cont.}}(V, k)$ since those of the U_i and $U_i \cap U_j$ have this form. Hence, it follows from Example 5 that if X is qcqs, it is the spectrum of an ind-étale algebra.

(2 \iff 4) This follows from the definition of $- \otimes S$ and the equivalence between the category of finite sets and the category of étale k -algebras. \square

Lemma 7. *Suppose that X is a topological space admitting a finite open cover $\{U_i \rightarrow X\}_{i=1}^n$ such that all U_i and $U_i \cap U_j$ are compact, Hausdorff, totally disconnected topological spaces. Then show that X is also compact, Hausdorff, and totally disconnected.*

Proof. X is compact: Suppose that $\{V_j \rightarrow X\}_{j \in J}$ is an open covering. then each $\{V_j \cap U_i\}$ is an open covering. But each U_i is compact, so for each $i = 1, \dots, n$,

there is a finite subset $J_i \subseteq J$ such that $U_i = \cup_{j \in J_i} U_i \cap V_j$. It follows that $X = \cup_{i=1}^n \cup_{j \in J_i} V_j$.

X is Hausdorff: Suppose that $x \neq y \in X$ are two points. Choose i_x, i_y such that $x \in U_{i_x}$ and $y \in U_{i_y}$ and set $U_x = U_{i_x}, U_y = U_{i_y}, U_{xy} = U_{i_x} \cap U_{i_y}$. If, say, $y \in U_{xy} \subseteq U_x$, then since U_x is Hausdorff, we can find opens $x \in V, y \in W$ such that $V \cap W = \emptyset$. So suppose that $x, y \notin U_{xy}$. Since U_{xy} is compact, and U_x, U_y are both profinite, U_{xy} is both closed and open in both U_x and U_y . In particular, $V = (U_x \setminus U_{xy}) \subseteq U_x$ and $W = (U_y \setminus U_{xy}) \subseteq U_y$ are also both closed and open in U_x, U_y respectively. This means that V and W are both open in X . By construction, $x \in V$ and $y \in W$ and $V \cap W = \emptyset$, so we are done.

X is totally disconnected: First recall that a subset $W \subseteq X$ is open (resp. closed) if and only if $W \cap U_i$ is open (resp. closed) for all i . Let us write $Y \Subset W$ to indicate that Y is both open and closed in W . Suppose that $W \subseteq X$ is a subset containing more than one point. We want to find a proper nonempty $Y \Subset W$. If $W \cap U_i$ has a single point, say w , for some i , then $\{w\}$ is open in W . But all U_i are totally disconnected, so $\{w\}$ is closed in all U_i , and therefore closed in X , and therefore closed in W . Hence, $Y = \{w\} \Subset W$ works.

So suppose each $W \cap U_i$ has more than one point. Since the U_i are totally disconnected, for each i there is some proper nonempty $Y_i \Subset W \cap U_i$. For any other j , we then have that $Y_i \cap U_j \Subset (W \cap U_i) \cap U_j$. Now as above, since $U_i \cap U_j$ is quasicompact, $U_i \cap U_j \Subset U_j$, so, $W \cap U_i \cap U_j \Subset W \cap U_j$, and we find that in fact, $Y_i \cap U_j \Subset W \cap U_i \cap U_j \Subset W \cap U_j$. Now define T_i inductively by setting $T_0 = W$. If one of $T_{i-1} \cap Y_i$ or $T_{i-1} \cap (W \cap U_i \setminus Y_i)$ are nonempty then choose one and set T_i to be this nonempty intersection. If both are empty, then define $T_i^a = T_{i-1}^a$. Now note that since $Y_i \cap U_j \Subset W \cap U_j$ for every i, j , it follows that each $T_i \Subset W \cap U_j$ for every $1 \leq j \leq i$. In particular, $T_n \Subset W \cap U_j$ for all j , and therefore $T_n \Subset W$. It is nonempty and proper by construction. \square

Corollary 8. *If k is a separably closed field, there is an equivalence of categories*

$$\begin{aligned} \text{ProFinSet} &\cong \text{Spec}(k)_{\text{proét}}^{\text{aff}} \\ S &\mapsto \text{Spec}(k) \otimes S \\ X(k) &\leftarrow X \end{aligned}$$

Under this identification, coverings of $\text{Spec}(k) \otimes S$ are precisely the jointly surjective families of profinite sets $\{S_i \rightarrow S\}_{i \in I}$ that admit a jointly surjective finite subfamily $\{S_{i_j} \rightarrow S\}_{j=1}^n$.

Example 9. If S is any nonfinite profinite set then the family $\{s \rightarrow S\}_{s \in S}$ of inclusions of its points is *not* a covering family.

The following is basically a version of the equivalence we saw in Galois theory between étale k -algebras and finite G -sets.

Proposition 10. *Let k be any field, choose a separable closure k^{sep}/k , and let $G = \text{Gal}(k^{\text{sep}}/k)$. There is an equivalence of categories between profinite sets*

equipped with a continuous G -action and the affine objects in $\mathrm{Spec}(k)_{\mathrm{pro\acute{e}t}}$.

$$G\text{-ProFinSet} \cong \mathrm{Spec}(k)_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$$

Under this identification, coverings of $\mathrm{Spec}(k) \otimes S$ are precisely the jointly surjective families of profinite sets $\{S_i \rightarrow S\}_{i \in I}$ that admit a jointly surjective finite subfamily $\{S_{i_j} \rightarrow S\}_{j=1}^n$.

Sketch of proof. In one direction, we use the functor

$$\mathrm{Spec}(k)_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}} \xrightarrow{k^{sep} \otimes_k -} \mathrm{Spec}(k^{sep})_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$$

and the equivalence

$$\mathrm{Spec}(k^{sep})_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}} \cong \mathrm{ProFinSet}$$

The G -action is induced by the canonical G -action on $\mathrm{Spec}(k^{sep})$. In the other direction, given a pro-finite set S equipped with a continuous G -action, we take $\mathrm{Spec}(\mathrm{hom}_{\mathrm{cont}}(S, k^{sep})^G)$, i.e., the spectrum of the ring of those continuous functions which are invariant for the action of G acting via its action on S . \square

3 The pro-étale topos

Definition 11 ([Def.4.2.1]). *Let X be a scheme. An object $U \in X_{\mathrm{pro\acute{e}t}}$ is called a pro-étale affine if it is of the form $U = \varprojlim U_i$ for some small filtered diagram $(U_i)_{i \in I}$ of (absolutely) affine schemes $U_i = \mathrm{Spec}(A_i)$ in X_{et} . The expression $U = \varprojlim U_i$ is called a presentation of U . The full subcategory of $X_{\mathrm{pro\acute{e}t}}$ spanned by pro-étale affines is denoted $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$. We make it a site by saying a family in $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$ is a covering in $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$ if it is a covering in $X_{\mathrm{pro\acute{e}t}}$.*

Lemma 12. *For X a scheme, every scheme $Y \in X_{\mathrm{pro\acute{e}t}}$ admits a pro-étale covering $\{Y_i \rightarrow Y\}$ such that each Y_i is in $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$.*

Proof. Choose an open affine covering $\{\mathrm{Spec}(A_i) \rightarrow X\}_{i \in I}$ of X , and for each i , choose an open affine covering $\{\mathrm{Spec}(B_{ij}) \rightarrow \mathrm{Spec}(A_i) \times_X Y\}_{i \in I, j \in J_i}$ of the preimage of $\mathrm{Spec}(A_i)$ in Y . Now by [Thm.2.3.4], since the morphisms $A_i \rightarrow B_{ij}$ are weakly étale for each i, j , there is a faithfully flat ind-étale morphism $B_{ij} \rightarrow C_{ij}$ such that $A_i \rightarrow C_{ij}$ is ind-étale. Consequently, $\{\mathrm{Spec}(C_{ij}) \rightarrow Y\}_{i \in I, j \in J_i}$ is a covering of the desired form. \square

Corollary 13 ([Lem.4.2.4, Rem.4.2.5]). *For any scheme X , the canonical restriction functor induces an equivalence of categories of sheaves*

$$\mathrm{Shv}(X_{\mathrm{pro\acute{e}t}}) \xrightarrow{\sim} \mathrm{Shv}(X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}).$$

Proof. This is a general fact about Grothendieck sites. Consider any site (C, τ) and full subcategory $D \subseteq C$ equipped with the induced topology. If every object of C has a covering by objects of D , then there is an equivalence $\mathrm{Shv}(C) \cong \mathrm{Shv}(D)$. \square

Proposition 14 ([Prop.4.2.8]). *For any scheme X , the topos $\mathrm{Shv}(X_{\mathrm{pro\acute{e}t}})$ is locally weakly contractible [Def.3.2.1]. In particular, it is replete [Def.3.1.1], and so $D(X_{\mathrm{pro\acute{e}t}})$ is left-complete [Def.3.3.1].*

Proof. [Prop.3.2.3] says that a locally weakly contractible topos is replete. [Prop.3.3.3] says that the derived category of a replete topos is left-complete. It suffices to show that for every scheme $Y \in X_{\mathrm{pro\acute{e}t}}$ there is a covering $\{Y_i \rightarrow Y\}_{i \in I}$ with each Y_i coherent and locally weakly contractible. Lemma 12 says that every scheme admits a pro-étale affine covering. So it remains only to see that affine schemes are coherent objects. This follows from the fact that affine schemes are qcqs. \square

On the pro-étale site, one can define interesting “constant” sheaves associated to topological spaces.

Lemma 15 ([Lem.4.2.12]). *Suppose X is a scheme and T is a topological space. Then the presheaf*

$$F_T : X_{\mathrm{pro\acute{e}t}}^{\mathrm{op}} \rightarrow \mathrm{Set}; \quad U \mapsto \mathrm{Map}_{\mathrm{cont}}(U, T)$$

which sends a scheme U to the set of continuous maps from the underlying topological space of U to T is a sheaf.

Sketch of proof. This uses [Lem.4.2.6] which we did not do. It says that a presheaf F on $X_{\mathrm{pro\acute{e}t}}$ is a sheaf if and only if it satisfies the sheaf condition for Zariski covers, and surjective maps in $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$. In the category of topological spaces, any representable presheaf is a sheaf for the topology generated by usual open coverings of topological spaces, and surjective morphisms $Y \rightarrow X$ such that X has the quotient topology induced from Y . Hence, in our setting, it suffices to check that for any surjective morphism $f : \mathrm{Spec}(B) \rightarrow \mathrm{Spec}(A)$ in $X_{\mathrm{pro\acute{e}t}}^{\mathrm{aff}}$ a subset $U \subseteq \mathrm{Spec}(A)$ is open if and only if f^{-1} is open. This is proved in a really neat way using the constructible topology, and the fact that a subset of a scheme is open if and only if it is constructible and closed under generisation. \square

4 Addendum

We did not have time for the following comments. There are of course many more details in Bhatt, Scholze.

Let k be a field, k^{sep} a separable closure, and $G = \mathrm{Gal}(k^{\mathrm{sep}}/k)$. Recall that we had an equivalence of categories

$$\mathrm{Shv}(k_{\mathrm{et}}, \mathrm{Ab}) \cong G\text{-mod}$$

between the category of étale sheaves on k , and discrete G -modules. A consequence of this was that for any discrete G -module M with associated sheaf F_M , the group cohomology of M is isomorphic to the étale sheaf cohomology of F_M ,

$$H_{\mathrm{et}}^n(k, F_M) \cong H^n(G, M).$$

The pro-étale site allows us to upgrade this, although things become more technical and complicated.

Recall that we have already seen an equivalence of categories

$$k_{\text{proét}}^{\text{aff}} \cong G\text{-ProFinSet}$$

between the subcategory of affine objects in $k_{\text{proét}}$ and the category of profinite sets equipped with a continuous action. The covering families in the left side are just surjective families.

Definition 16. *Given an arbitrary profinite group G , we define a topology on the category $G\text{-ProFinSet}$ whose covering families are surjective families.*

Definition 17. *Let $G\text{-Spc}$ be the category of topological spaces equipped with a continuous G -action. Let $G\text{-Spc}_{\text{cg}} \subseteq G\text{-Spc}$ be the full subcategory of $X \in G\text{-Spc}$ whose underlying topological space can be written as a quotient of a disjoint union of compact Hausdorff spaces. These spaces are called compactly generated.*

Lemma 18 ([Lem.4.3.2]). *The association $T \mapsto \text{hom}_{\text{cont},G}(-, T)$ produces a functor $G\text{-Spc} \rightarrow \text{Shv}(G\text{-ProFinSet})$. This functor is fully faithful on $G\text{-Spc}_{\text{cg}}$, admits a left adjoint (everywhere), and its essential image generates $\text{Shv}(G\text{-ProFinSet})$ under colimits.*

Definition 19. *We write $G\text{-Mod}$ for the category of topological abelian groups equipped with a continuous G -action. We write $G\text{-Mod}_{\text{cg}}$ for the full subcategory whose underlying space is compactly generated (i.e., lies in $G\text{-Spc}_{\text{cg}}$).*

As above, given $M \in G\text{-Mod}$, we get an abelian sheaf $F_M : X \mapsto \text{hom}_{\text{cont},G}(-, M)$ on $G\text{-ProFinSet}$.

We did not define continuous cohomology, but the main result about it is the following.

Lemma 20 ([Lem.4.3.9]). *For a large class of “nice” $M \in G\text{-Mod}$ we have*

$$H_{\text{cont}}^n(G, M) \cong H_{\text{proét}}^n(G\text{-ProFinSet}, F_M).$$