

0 Introduction

In the introduction to the pro-étale topology we argued that adding (filtered) limits to the site made \mathbb{Z}_l - and \mathbb{Q}_l -sheaves work better. One of the technical reasons that the category works better is that schemes morally become “locally contractible”. More concretely, every scheme X admits a pro-étale covering $\{U_i \rightarrow X\}_{i \in I}$ such that each U_i is *weakly contractible* in the sense that for every pro-étale covering $\{V_{ij} \rightarrow U_i\}$ the morphism $\coprod_j V_{ij} \rightarrow U_i$ has a section $\coprod_j V_{ij} \xrightarrow{\sim} U_i$.

The goal of this lecture is to show the following:

Today’s Goal. Every affine scheme X admits a surjective pro-étale morphism $U \rightarrow X$ such that U is affine and weakly contractible.

This happens in roughly four steps:

1. (Zariski case) Build a surjective pro-Zariski morphism $X^Z \rightarrow X$ with X^Z weakly contractible for the Zariski topology.
2. (Profinite set case) Build a surjective morphism $T \rightarrow (X^Z)^c$ to the set $(X^Z)^c$ of closed points of X^Z from a profinite set T which is weakly contractible as a compact Hausdorff topological space.
3. (Dimension zero scheme case) Give T a structure of affine scheme X_0 such that all residue fields are separably closed.
4. Henselise along $X_0 \rightarrow X$ to produce the desired $U \rightarrow X$.

If X has finitely many points (e.g., $X = \text{Spec}(R)$ with R a discrete valuation ring, or more generally, a localisation of a Dedekind domain at finitely many primes) then X^Z in Step 1 is just the disjoint union $\coprod_{x \in X} \text{Spec}(\mathcal{O}_{X,x})$ of the localisations at each point of X , Step 2 is redundant, Step 3 just chooses separable closures for each $k(x)$, and Step 4 produces the disjoint union of the strict henselisations $\coprod_{x \in X} \text{Spec}(\mathcal{O}_{X,x}^{sh}) \rightarrow X$.

The general case is not so easy because, for example, $\coprod_{x \in X} x$ is not affine if X has infinitely many points (all affine schemes are quasi-compact). However, instead of just making things more complicated, things actually become very, very interesting. The scheme $X^Z \rightarrow X$ that is produced is in a precise sense the “smallest” Zariski covering of X . As a set, X^Z is the disjoint union $\coprod_{x \in X} \text{Spec}(\mathcal{O}_{X,x})$ of the localisations of X but the topology is coarser: the subset $(X^Z)^c$ of closed points of X^Z is also affine, and has the curious property of being homeomorphic to the set of points of X equipped with the constructible topology.

Step 2 is clearly necessary for the following reason: strictly hensel local schemes X have the property that given any étale morphism $U \rightarrow X$, any lift of the closed point $X^c \rightarrow U \rightarrow X$ can be extended to a section $X \rightarrow U \rightarrow X$. So if our set of closed points is not weakly contractible, there is not much hope for X to be.

Step 3 and Step 4 are classical, but Bhatt-Scholze’s approach to separably closing the residue fields in Step 3 quite amusing; they take the fibre product of *all* finite presentation surjective étale morphisms. Of course, this produces something huge, but certainly also produces something which has all residue fields separably closed.

1 Pro-Zariski covers of affine schemes

In this section, we consider the question: is there a “smallest” open cover, and what does it look like?

The category of *spectral spaces* is the image of the functor

$$(\mathrm{Spec}(-))_{\mathrm{top}} : (\mathrm{Ring})^{\mathrm{op}} \rightarrow \mathrm{Top}$$

which sends a ring to its set of primes equipped with the Zariski topology. Of course, different rings can give rise to the same space (e.g, fields, discrete valuation rings, noetherian dimension one schemes, ...) but none-the-less, if we are only interested in the Zariski topology, all we are concerned with is the underlying topological space.

Let \mathcal{S} be the category of spectral spaces (i.e., spaces of the form $\mathrm{Spec}(R)$ for some ring R), with spectral maps (i.e., maps of the form $\mathrm{Spec}(R) \rightarrow \mathrm{Spec}(S)$ for some ring homomorphism $S \rightarrow R$).

Definition 1 (Def.2.1.1). *A spectral space X is w -local if it satisfies:*

1. *All open covers split, i.e., for every open cover $\{U_i \hookrightarrow X\}$, the map $\sqcup_i U_i \rightarrow X$ has a section.*
2. *The subspace $X^c \subset X$ of closed points is closed.*

A map $f : X \rightarrow Y$ of w -local spaces is w -local if f is spectral and $f(X^c) \subset Y^c$. Let $i : \mathcal{S}^{wl} \hookrightarrow \mathcal{S}$ be the subcategory of w -local spaces with w -local maps.

Exercise 1 (Exa.2.1.2). Show that the following spectral spaces are w -local.

1. Any profinite set (with the profinite topology).
2. The topological space $\mathrm{Spec}(\mathcal{O}_{X,x})$ underlying any local ring of any scheme, i.e., a topological space with a unique closed point.
3. Any finite disjoint union of w -local spaces.

Exercise 2. Show that if X is w -local then every connected component has at most one closed point.

Remark 2. In fact, if X is w -local, then every connected component was exactly one closed point. Indeed: Suppose Z has two closed points $x, y \in Z$. Every connected component is closed, so x, y are closed in X as well. Then $X \setminus \{x\} \amalg X \setminus \{y\} \rightarrow X$ has a section because X is w -local. This induces a decomposition

of Z into two non-empty clopen subsets, contradicting the assumption that Z is connected. On the other hand, since X is spectral, Z is also spectral, and every spectral space has at least one closed point (cf. every ring has at least one maximal ideal).

----- picture: triangles over the cantor set $C \times \text{Spec}(\mathcal{O}_{X,x})$ -----

Definition 3 (Def.2.1.12). *A map $f : W \rightarrow V$ of spectral spaces is a Zariski localization if $W = \sqcup_{j=1}^n U_j$ with the $U_j \rightarrow V$ open immersions (automatically quasi-compact because W is spectral). A pro-(Zariski localization) is a cofiltered limit of such maps.*

Example 4.

1. For any profinite set $I = \varprojlim I_\lambda$ the map $\varprojlim \prod_{I_\lambda} V \rightarrow V$ is a pro-(Zariski localisation).

----- picture: cantor set of copies of some space -----

2. For any point $v \in V$, the map $\bigcap_{v \in U} U \rightarrow V$ is a pro-(Zariski localisation), where the intersection is over quasicompact opens containing v .

----- picture: triangle representing a localisation at a point -----

3. We can combine the above two to describe every pro-(Zariski localisation). Suppose that $W \rightarrow V$ is a pro-(Zariski localisation), so $W = \varprojlim_{i \in I} \sqcup_{j=1}^{n_i} U_{ij}$ with each $U_{ij} \rightarrow V$ a quasicompact open immersion. Then $W \subseteq \varprojlim_{i \in I} \sqcup_{j=1}^{n_i} V$. As a set, $\varprojlim_{i \in I} \sqcup_{j=1}^{n_i} V$ is a disjoint union of copies of V indexed by the profinite set $\varprojlim_{i \in I} \{1, \dots, n_i\}$. Given an element $(j_i)_{i \in I} \in \varprojlim_{i \in I} \{1, \dots, n_i\}$, the intersection W with the $(j_i)_{i \in I}$ th copy of V is $\bigcap_{i \in I} U_{ij_i}$.

----- picture: $U \subseteq \mathbb{N}^* \times \text{Spec } \mathbb{Z}$ -----

Lemma 5 (Lem.2.1.10). *The inclusion $i : \mathcal{S}^{wl} \rightarrow \mathcal{S}$ admits a right adjoint $(-)^Z : \mathcal{S} \rightarrow \mathcal{S}^{wl}$. The counit $X^Z \rightarrow X$ is a surjective pro-(Zariski localisation) for all X , and the composite $(X^Z)^c \rightarrow X$ is a homeomorphism for the constructible topology on X .*

The Z probably stands for “Zariski”.

Remark 6. The family of *constructible* subsets of an affine scheme $\text{Spec}(A)$ is the smallest family closed under finite intersection, finite union, complement, and containing the closed subsets $V(I)$ for every *finitely generated* ideal I (a ring is noetherian if and only if every ideal is finitely generated; we will often need non-noetherian rings). The constructible topology is a (usual) topology on X whose opens are the constructible subsets.

Exercise 3. Describe the constructible subsets of $\text{Spec}(\mathbb{Z})$ and $\text{Spec}(\mathbb{C}[x, y])$.

Sketch of proof. The idea is that every spectral space is an inverse limit of finite spectral spaces, and for finite spectral spaces, X^Z is the disjoint union of the localisations at each point $X^Z = \coprod_{x \in X} X_x$.

— — — picture: X^Z for a dvr — — —

□

Exercise 4. Prove that if X is a finite spectral space, then any map $Y \rightarrow X$ from a w-local space factors through $\coprod_{x \in X} X_x \rightarrow X$. Hint: use the fact that each $X_x \rightarrow X$ is an open immersion, and in fact, $\{X_x \rightarrow X\}_{x \in X}$ is an open covering.

Remark 7 (Omitted from lecture). In fact, every spectral space is the inverse limit of its constructible partitions: Let $X = \cup_{i \in I} X_i$ be a partition of X into constructible sets, and consider the canonical projection $X \rightarrow I$ to the *components* of the partition (so $x \in X$ is sent to the i such that $x \in X_i$). We equip I with the coarsest topology which makes this map continuous (so a subset $J \subset I$ is open iff its preimage is). Then the topological space of X is the inverse limit over these projections.

Remark 8 (Rem.2.1.11, Omitted from lecture). The space X^Z can be alternatively described as:

$$X^Z = \varprojlim_{\{X_i \hookrightarrow X\}} \sqcup_i \widetilde{X}_i,$$

where the limit is indexed by the cofiltered category of constructible stratifications $\{X_i \hookrightarrow X\}$, and \widetilde{X}_i denotes the set of all points of X specializing to a point of X_i . As a set, X^Z is the disjoint union of the localisations at each point $X^Z = \coprod_{x \in X} X_x$, but the topology is coarser.

Example 9. Let X be the topological space associated to a curve (e.g., $\text{Spec}(\mathbb{Z})$ or $\text{Spec}(k[t])$), so X has one generic point η , every other point is closed, and the non-empty proper closed subsets are finite sets of closed points. Then X^Z as a set is $\{\eta\} \sqcup \coprod_{x \in X^c} \{x, \eta_x\}$. The sets $\{x, \eta_x\}$ and $\{\eta_x\}$ are open, as well as $\{\eta\} \sqcup \coprod_{x \in S} \{x, \eta_x\}$ for any cofinite set $S \subseteq X^c$ of closed points. These open sets generate the topology. Note that $\{\eta\}$ is closed, but not open. Note also that the topology induced on $\{\eta\} \cup X^c$ is the constructible topology of X .

From a ring point of view,

$$\text{Spec}(\mathbb{Z})^Z = \text{Spec} \left(\varinjlim_{\text{primes } p} \mathbb{Z}_{(2)} \times \mathbb{Z}_{(3)} \times \mathbb{Z}_{(5)} \times \cdots \times \mathbb{Z}_{(p)} \times (\mathbb{Z}[\frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{p}]) \right)$$

The transition morphisms are the identities on the $\mathbb{Z}_{(p)}$ factors, and the diagonal $\mathbb{Z}[\frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{p}] \rightarrow \mathbb{Z}_{(p')} \times \mathbb{Z}[\frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{p}, \frac{1}{p'}]$ on the end.

The following lemma is used in Lem.2.2.8 to transfer a scheme structure from T to S .

Lemma 10 (Lem.2.1.14). *Any map $f : X \rightarrow Y$ in \mathcal{S}^{wl} admits a canonical factorization $X \rightarrow Z \rightarrow Y$ in \mathcal{S}^{wl} with $Z \rightarrow Y$ a pro-(Zariski localization) and $X \rightarrow Z$ inducing a homeomorphism $X^c \simeq Z^c$.*

Proof. (Omitted from lecture). Take $Z = \pi_0(X) \times_{\pi_0(Y)} Y$. Here $\pi_0(X), \pi_0(Y)$ are the sets of connected components of X, Y . The morphism $\pi_0(X) \rightarrow \pi_0(Y)$ is a morphism of profinite sets, and therefore a pro-(Zariski localisation). Hence, its pullback $\pi_0(X) \times_{\pi_0(Y)} Y \rightarrow Y$ is also a pro-(Zariski localisation). The map $X \rightarrow \pi_0(X)$ induces a homeomorphism $X^c \rightarrow \pi_0(X)$ because each connected component of X has a unique closed point. For the same reason $(\pi_0(X) \times_{\pi_0(Y)} Y)^c \rightarrow \pi_0(\pi_0(X) \times_{\pi_0(Y)} Y)$ is a homeomorphism. But $\pi_0(\pi_0(X) \times_{\pi_0(Y)} Y) \cong \pi_0(X)$, so it follows that $X^c \rightarrow Z^c$ is a homeomorphism. \square

2 Rings

Definition 11 (Def.2.2.1). *Fix a ring A .*

1. A is w -local if $\text{Spec}(A)$ is w -local.
2. A is w -strictly local if A is w -local, and every faithfully flat étale map $A \rightarrow B$ has a section.
3. A map $f : A \rightarrow B$ of w -local rings is w -local if $\text{Spec}(f)$ is w -local.
4. A map $f : A \rightarrow B$ is called a Zariski localization if $B = \prod_{i=1}^n A[\frac{1}{f_i}]$ for some $f_1, \dots, f_n \in A$. An ind-(Zariski localization) is a filtered colimit of Zariski localizations.
5. A map $f : A \rightarrow B$ is called ind-étale if it is a filtered colimit of étale A -algebras.

Example 12.

1. We have already seen an example of an ind-(Zariski localisation) above:

$$\varinjlim_{\text{primes } p} \mathbb{Z}_{(2)} \times \mathbb{Z}_{(3)} \times \mathbb{Z}_{(5)} \times \cdots \times \mathbb{Z}_{(p)} \times (\mathbb{Z}[\frac{1}{2}, \frac{1}{3}, \dots, \frac{1}{p}])$$

2. Let k be a field, and k^{sep} a separable closure. Then $k \rightarrow k^{sep}$ is étale if and only if k^{sep}/k is a finite extension, but it is always ind-étale.
3. Any field k and any profinite set $T = \varprojlim_{i \in I} T_i$, gives rise to a k -algebra $\varinjlim_{i \in I} \prod_{t \in T_i} k \subseteq \prod_{t \in T} k$. The algebra $\varinjlim_{i \in I} \prod_{t \in T_i} k$ has the property that each residue field is isomorphic to k , and it's topological space is homeomorphic to the profinite set T .

Recall that a ring B is called *absolutely flat* if B is reduced with Krull dimension 0 (or, equivalently, that B is reduced with $\text{Spec}(B)$ Hausdorff). Absolutely flat rings are also characterised as those rings such that every module is flat [Stacks project, 092F]. An example is $\bar{k} \otimes_k \bar{k}$ for any perfect field k . Note that $\text{Spec}(\bar{k} \otimes_k \bar{k}) \cong \text{Gal}(\bar{k}/k)$ as *topological spaces*.

So $\text{Spec}(B)$ of an absolutely flat ring has a profinite set as its topological space,¹ and all local rings are fields.² So it is some kind of “profinite product” of fields.

Lemma 13 (Lem.2.2.3). *If A is w -local, then the Jacobson radical $I_A (= \bigcap_{\text{maximal ideals } \mathfrak{m}} \mathfrak{m})$ cuts out $\text{Spec}(A)^c \subset \text{Spec}(A)$ with its reduced structure. The quotient A/I_A is an absolutely flat ring.*

Proof. (Omitted from lecture). Let $J \subset A$ be the (radical) ideal cutting out $\text{Spec}(A)^c \subset \text{Spec}(A)$ with the reduced structure. Then $J \subset \mathfrak{m}$ for each $\mathfrak{m} \in \text{Spec}(A)^c$, so $J \subset I_A$. Hence, $\text{Spec}(A/I_A) \subset \text{Spec}(A)^c$ is a closed subspace; we want the two spaces to coincide. If they are not equal, then there exists a maximal ideal \mathfrak{m} such that $I_A \not\subset \mathfrak{m}$, which is impossible. \square

Lemma 14 (Lem.2.2.4). *The inclusion of the category w -local rings and maps inside all rings admits a left adjoint $A \mapsto A^Z$. The unit $A \rightarrow A^Z$ is a faithfully flat ind-(Zariski localization), and $\text{Spec}(A)^Z = \text{Spec}(A^Z)$ over $\text{Spec}(A)$.*

Proof. (Omitted from lecture). This follows from Remark 2.1.11 (above). In more details, let $X = \text{Spec } A$, and define a ringed space $X^Z \rightarrow X$ by equipping $(\text{Spec } A)^Z$ with the pullback of the structure sheaf from X . Then Remark 2.1.11 presents X^Z as an inverse limit of affine schemes, so that $X^Z = \text{Spec}(A^Z)$ is itself affine. \square

Lemma 15 (Lem.2.2.6). *Any w -local map $f : A \rightarrow B$ of w -local rings admits a canonical factorization $A \xrightarrow{a} C \xrightarrow{b} B$ with C w -local, $a : A \rightarrow C$ a w -local ind-(Zariski localization), and $b : C \rightarrow B$ a w -local map inducing $\pi_0(\text{Spec}(B)) \simeq \pi_0(\text{Spec}(C))$.*

Lemma 16 (Lem.2.2.7). *For any absolutely flat ring A , there is an ind-étale faithfully flat map $A \rightarrow \bar{A}$ with \bar{A} w -strictly local and absolutely flat. For a map $A \rightarrow B$ of absolutely flat rings, we can choose such maps $A \rightarrow \bar{A}$ and $B \rightarrow \bar{B}$ together with a map $\bar{A} \rightarrow \bar{B}$ of A -algebras.*

¹Recall that profinite sets are precisely the quasi-compact Hausdorff totally disconnected spaces. Every affine scheme is quasi-compact. We observed above that absolutely flat rings have Hausdorff spectrum. Suppose $\text{Spec}(B/I) \subseteq \text{Spec}(B)$ is a connected component. We can assume $I = \sqrt{I}$, and so $\text{Spec}(B/I)$ is again reduced of Krull dimension zero, so we can in fact, assume $B = B/I$. Since $\text{Spec}(B)$ is Hausdorff, for any two points $\mathfrak{m}_0, \mathfrak{m}_1$, there are disjoint opens containing each of them. Shrinking, we can assume the opens are of the form $\text{Spec}(B_{f_0}), \text{Spec}(B_{f_1})$. The complement of union of these two opens is the closed subset of the ideal $\langle f_0, f_1 \rangle$. Since every module is flat however, the support of this ideal is open, and we deduce that $\text{Spec}(B)$ is a disjoint union of three opens, two of which are nonempty by virtue of containing the closed points $\mathfrak{m}_0, \mathfrak{m}_1$ from the beginning. This contradicts $\text{Spec}(B)$ being connected.

²Indeed, since all modules are flat, for any maximal ideal \mathfrak{m} , the module A/\mathfrak{m} is flat. It follows that $A/\mathfrak{m} = A_{\mathfrak{m}}$.

Proof. The following fact is used without further comment below: any ind-étale algebra over an absolutely flat ring is also absolutely flat. Choose a set I of isomorphism classes of faithfully flat étale A -algebras, and set $\overline{A} = \otimes_I A_i$, where the tensor product takes place over $A_i \in I$, i.e., $\overline{A} = \varinjlim_{J \subset I} \otimes_{j \in J} A_j$, where the (filtered) colimit is indexed by the poset of finite subsets of I . Then one checks that \overline{A} is absolutely flat, and that any faithfully flat étale \overline{A} -algebra has a section, so \overline{A} is w-strictly local as $\text{Spec}(\overline{A})$ is profinite. For the second part, simply set \overline{B} to be a w-strictly local faithfully flat ind-étale algebra over $\overline{A} \otimes_A B$. \square

Remark 17. Note that one can construct the algebraic closure of a field k using this method. Take the tensor product over all finite algebraic extensions of k . This will give some complicated ring but every residue field will be an algebraic closure of k .

Note that if A is w -local then $\pi_0(\text{Spec}(A))$ is canonically homeomorphic to the set $\text{Spec}(A)^c$ of closed points of $\text{Spec}(A)$.

Lemma 18 (Lem.2.2.8). *For any ring A and a map $T \rightarrow \pi_0(\text{Spec}(A))$ of profinite sets, there is an ind-(Zariski localization) $A \rightarrow B$ such that $\text{Spec}(B) \rightarrow \text{Spec}(A)$ gives rise to the given map $T \rightarrow \pi_0(\text{Spec}(A))$ on applying π_0 . Moreover, the association $T \mapsto \text{Spec}(B)$ is a limit-preserving functor.*

— — — — picture — — — —

Remark 19 (No number). One may make the following more precise statement: for any affine scheme X , the functor $Y \mapsto \pi_0(Y)$ from affine X -schemes to profinite $\pi_0(X)$ -sets has a fully faithful right adjoint $S \mapsto S \times_{\pi_0(X)} X$, the fibre product in the category of topological spaces ringed using the pullback of the structure sheaf on X . Moreover, the natural map $S \times_{\pi_0(X)} X \rightarrow X$ is a pro-(Zariski localisation) and pro-finite.

Proof. Notice that $\pi_0(\text{Spec}(A)) = \varinjlim_{I \in D} I$ where the limit is over the set $D = \{\text{Spec}(A) = \sqcup_{i \in I} Z_i\}$ of decompositions of $\text{Spec}(A)$ into finitely many subsets $Z_i = \text{Spec}(A_i)$ which are both open and closed. Writing $T = \varprojlim_{\lambda \in \Lambda} T_\lambda$, the map $T \rightarrow \pi_0(\text{Spec}(A))$ comes from a map of pro-systems $\phi : (T_\lambda)_\Lambda \rightarrow (I)_{I \in D}$. For each $\lambda \in \Lambda$ define $B_\lambda = \prod_{t \in T_\lambda} \Gamma(T_{\phi_\lambda(t)}, \mathcal{O}_{\text{Spec}(A)})$. Then $\varinjlim_{\lambda \in \Lambda} B_\lambda$ satisfies the requirements. \square

3 Henselisation

To pass from w-strictly local covers of absolutely flat rings to arbitrary rings, we use henselizations:

Definition 20 (Def.2.2.10). *Given a map of rings $A \rightarrow B$, let $\text{Hens}_A(-) : \text{Ind}(B_{\text{et}}) \rightarrow \text{Ind}(A_{\text{et}})$ be the functor right adjoint to the base change functor*

$\text{Ind}(A_{\text{et}}) \rightarrow \text{Ind}(B_{\text{et}})$. Explicitly, for $B_0 \in \text{Ind}(B_{\text{et}})$, we have $\text{Hens}_A(B_0) = \varinjlim A'$, where the colimit is indexed by diagrams $A \rightarrow A' \rightarrow B_0$ of A -algebras with $A \rightarrow A'$ étale.

Lemma 21 (Lem.2.2.13). *Let A be a ring henselian along an ideal I . Then A is w-strictly local if and only if A/I is so.*

Corollary 22 (Cor.2.2.14). *Any ring A admits an ind-étale faithfully flat map $A \rightarrow A'$ with A' w-strictly local.*

Proof. Set $A' := \text{Hens}_{A^Z}(\overline{A^Z/I_{A^Z}})$, where $\overline{A^Z/I_{A^Z}}$ is a w-strictly local ind-étale faithfully flat A^Z/I_{A^Z} -algebra; then A' satisfies the required property by Lemma 2.2.13. \square

4 Weakly étale versus pro-étale

Definition 23 (Def.2.3.1). *A morphism $A \rightarrow B$ of commutative rings is called weakly étale if both $A \rightarrow B$ and the multiplication morphism $B \otimes_A B \rightarrow B$ are flat.*

p:PropWeaklyEtale

Proposition 24 (Prop.2.3.3). *Fix maps $f : A \rightarrow B$, $g : B \rightarrow C$, and $h : A \rightarrow D$ of rings.*

1. *If f is ind-étale, then f is weakly étale.*
2. *If f is weakly étale and finitely presented, then f is étale.*
3. *If f and g are weakly étale (resp. ind-étale), then $g \circ f$ is weakly étale (resp. ind-étale). If $g \circ f$ and f are weakly étale (resp. ind-étale), then g is weakly étale (resp. ind-étale).*
4. *If h is faithfully flat, then f is weakly étale if and only if $f \otimes_A D : D \rightarrow B \otimes_A D$ is weakly étale.*

Theorem 25 (Thm.2.3.4). *Let $f : A \rightarrow B$ be weakly étale. Then there exists a faithfully flat ind-étale morphism $g : B \rightarrow C$ such that $g \circ f : A \rightarrow C$ is ind-étale.*

Proof. Lemma 2.3.7 (omitted from the lecture) gives a diagram

$$\begin{array}{ccc} A & \longrightarrow & A' \\ \downarrow f & & \downarrow f' \\ B & \longrightarrow & B' \end{array}$$

with f' a w-local map of w-strictly local rings, and both horizontal maps being ind-étale and faithfully flat. The map f' is also weakly étale since all other maps in the square are so. Then f' is a ind-(Zariski localization) by Lemma 2.3.8 which says that any w-local weakly étale map of w-local rings from a w-strictly local ring is an ind-(Zariski localisation). Setting $C = B'$ then proves the claim. \square

5 Local contractibility

Definition 26 (Def.2.4.1). *A ring A is w -contractible if every faithfully flat ind-étale map $A \rightarrow B$ has a section.*

Lemma 27 (Lem.2.4.2). *A w -contractible ring A is w -local (and thus w -strictly local).*

Lemma 28 (Lem.2.4.3). *Let A be a ring henselian along an ideal I . Then A is w -contractible if and only if A/I is so.*

Definition 29 (Def.2.4.4). *A profinite set is extremally disconnected if the closure of every open is open.*

Remark 30. We are interested in extremally disconnected spaces because they are weakly local. In fact, a theorem of Gleason from 1958 says that they are exactly the projective objects in the category of all compact Hausdorff spaces: any continuous surjection $Y \rightarrow X$ from a compact Hausdorff space to an extremally disconnected profinite set has a section.

Example 31 (Exa.2.4.6). Every compact Hausdorff space X admits a continuous surjection from an extremally disconnected space. (Proof omitted).

Lemma 32 (Lem.2.4.8). *A w -strictly local ring A is w -contractible if and only if $\pi_0(\text{Spec}(A))$ is extremally disconnected.*

Proof. Suppose A is w -contractible and $T \rightarrow \pi_0(\text{Spec}(A))$ is a continuous surjection of profinite sets. Then $\text{Spec}(T) \times_{\pi_0(\text{Spec}(A))} T \rightarrow \text{Spec}(A)$ is a pro-(Zariski-localisation) and therefore has a section by w -contractibility. Composing with $\text{Spec}(A)^c \rightarrow \text{Spec}(A)$, we get a factorisation $\text{Spec}(A)^c \rightarrow T \rightarrow \pi_0(\text{Spec}(A))$, but $\text{Spec}(A)^c \rightarrow \pi_0(\text{Spec}(A))$ is an isomorphism (by w -locality of A) so we have found a section to $T \rightarrow \pi_0(\text{Spec}(A))$.

The converse reduces the dimension zero (i.e., absolutely flat) case. In this case since the residue fields of A are all separable closed, every ind-étale faithfully flat A -algebra B is induced by a continuous surjection of profinite sets $T \rightarrow \text{Spec}(A)$. Then we just apply that $\pi_0(\text{Spec}(A))$ is extremally disconnected. \square

lem:cwcontractiblecover

Lemma 33 (Lem.2.4.9). *For any ring A , there is an ind-étale faithfully flat A -algebra A' with A' w -contractible.*

Proof. By Lemmas 2.1.10, 2.2.3, and 2.2.4, the pro-finite set $\text{Spec}(A^Z/I_{A^Z})$ is homeomorphic to $\text{Spec}(A)$ with the constructible topology, where I_{A^Z} is the Jacobson radical. In particular, it is a compact Hausdorff space. Choose a continuous surjection $T \rightarrow \text{Spec}(A^Z/I_{A^Z})$ from an extremally disconnected profinite set as mentioned in Example 2.4.6. Using Lemma 2.2.8, choose an algebra $A^Z/I_{A^Z} \rightarrow B$ such that $T = \pi_0(B)$. Using Lemma 2.2.7, we find an ind-étale faithfully flat A^Z/I_{A^Z} -algebra A_0 with A_0 w -strictly local and $\text{Spec}(A_0)$ an extremally disconnected profinite set. Let $A' = \text{Hens}_{A^Z}(A_0)$. Then A_0 is w -contractible because $\pi_0(\text{Spec}(A_0))$ is extremally disconnected (Lem.2.4.8), and A' is w -contractible because A_0 is (Lemma 2.4.3). The map $A \rightarrow A'$ is faithfully flat and ind-étale since both $A \rightarrow A^Z$ and $A^Z \rightarrow A'$ are so individually. \square

A Some point-set topology

A.1 Profinite sets

Definition 34. A profinite set is a filtered inverse limit $T = \varprojlim T_i$ of finite sets T_i , equipped with the limit topology. So the open sets are (possibly infinite) unions of sets of the form $\pi_i^{-1}(t)$ where $\pi_i : T \rightarrow T_i$ is the canonical projection and $t \in T_i$.

Example 35.

1. Any finite set is a profinite set.
2. The set $\{0\} \cup \{\frac{1}{n} : n = 1, 2, 3, \dots\}$ is profinite.
3. The Cantor set is profinite.
4. Any product of profinite sets is profinite.

- - - - *** add picture *** - - - -

Proposition 36 ([Stacks Project, Tag 08ZY]). A topological space is a profinite set if and only if it is compact³ Hausdorff⁴ and totally disconnected⁵.

This proposition gives a canonical choice for the filtered limit.

Lemma 37. Let T be a profinite set. Then $T = \varprojlim_{\sqcup_{i \in I} T_i} I$ where the limit is over the finite partitions $T = \sqcup_{i \in I} T_i$ of T into a disjoint union of subsets T_i which are both open and closed.

Note that any closed open U is compact (any covering of U extends to a covering of T by adjoining $T \setminus U$, and T is compact). On the other hand, any compact open U is a finite union of basic opens⁶ and these are closed, so U is closed.

So, to summarise:

$$\left\{ \begin{array}{c} \text{basic} \\ \text{opens} \end{array} \right\} \subseteq \left\{ \begin{array}{c} \text{compact} \\ \text{opens} \end{array} \right\} = \left\{ \begin{array}{c} \text{closed} \\ \text{opens} \end{array} \right\}$$

³Compact means every open cover admits a finite subcover. This property is called *quasi-compact* when talking about schemes.

⁴Hausdorff means for every pair of distinct points $x \neq y$ there are open sets U, V with $x \in U, y \in V$ and $U \cap V = \emptyset$. When talking about schemes, we say quasi-compact for this same property.

⁵Totally disconnected means that every subset $V \subseteq T$ containing more than one point can be written as a disjoint union $V = V_1 \amalg V_2$ of nonempty sets $V_1, V_2 \subseteq T$ both of which are both open and closed.

⁶By *basic* we mean the preimage of some point $t \in T_i$ for some presentation $T = \varprojlim_{i \in I} T_i$.

A.2 Finite sober spaces

A topological space X is called *sober*⁷ if every irreducible closed subset has a unique generic point. I.e., if a closed subset Z is irreducible, then Z is the closure $Z = \overline{\{\eta\}}$ of some unique point η). The reader is probably already familiar with finite sober spaces in a different form:

Proposition 38. *The category of finite sober spaces is equivalent to the category of finite partially ordered sets.*

Proof. Given a partially ordered set P we define $X(P)$ to be the topological space whose points are the points of P , and whose opens are the “upwards closed” subsets. I.e., subsets U satisfying

$$x \in U, x \leq y \Rightarrow y \in U.$$

Exercise: check that the U really form a topology.

Conversely, given a topological space X we define $P(X)$ to be the partially ordered set whose points are the points of X , and relation


$$x \leq y \text{ iff } y \in \bigcap_{U \ni x} U$$

where the intersection is over all opens U containing x .

Exercise: Check that we have $x \leq y$ and $y \leq z \Rightarrow x \leq z$, that we have $x \leq x$, and that when X is sober we have $x \leq y$ and $y \leq x \Rightarrow x = y$. Note that an equivalent condition for $x \leq y$ is that $x \in \bigcap_{Z \ni y} Z$ where the intersection is over all closed subsets containing y (its straightforward to prove $y \notin \bigcap_{U \ni x} U \iff x \notin \bigcap_{Z \ni y} Z$).

Finally, one should check that $X(P(X)) = X$ and that $P(X(P)) = P$, and also that the operations $X \mapsto P(X)$ and $P \mapsto X(P)$ send continuous morphisms (resp. morphisms of partially ordered sets) to morphisms of partially ordered sets (resp. continuous morphisms). \square

So it’s quite easy to produce examples of finite sober spaces.



⁷From Johnstone’s Topos Theory book, p.230: “If we regard two distinct points having the same closure as an instance of double vision (and an irreducible closed set with no generic point as a species of pink elephant!), then the reason for the term ‘sober space’ will be apparent.”

A.3 Spectral spaces

Definition 39. A spectral space is a topological space of the form $\text{Spec}(R)$ for some ring R .

Spaces of the form $\text{Spec}(R)$ satisfy a number of nice properties.

1. Each $\text{Spec}(R)$ is sober.
2. Each $\text{Spec}(R)$ is quasi-compact, i.e., every open cover $\{U_i \rightarrow \text{Spec}(R)\}_{i \in I}$ has a finite subcover.⁸
3. The topology on $\text{Spec}(R)$ is generated by the opens of the form $D(f) = \text{Spec}(R[\frac{1}{f}])$; note that these are also quasi-compact.
4. In fact an open $U \subset \text{Spec}(R)$ is quasi-compact if and only if $\text{Spec}(R) \setminus U = V(I)$ with I a finitely generated ideal. Consequently, intersections of quasi-compact opens of $\text{Spec}(R)$ are also quasi-compact.

It is an old theorem of Hochster that the above four properties *characterise* spectral spaces: I.e., A topological space X is (i) sober, (ii) quasi-compact, (iii) has topology generated by quasi-compact opens, and (iv) has it's set of quasi-compact opens preserved by finite intersection if and only if there is some ring R with $\text{Spec}(R)$ homeomorphic to X . Moreover, a continuous morphism of spectral spaces is spectral if and only if the inverse image of any quasicompact open is quasicompact.

In particular, we can deduce from this that for every finite sober space X , there is some ring R with $X = \text{Spec}(R)$. In light of the above proposition, an equivalent statement is: every partially ordered set is the partially ordered set of primes of some ring.

On the other hand, there is another characterisation of spectral spaces that is useful for us.

Proposition 40. A topological space is spectral if and only if it is a filtered inverse limit of finite sober spaces.

B Constructible sets

Just as any profinite set X is the filtered inverse limit of it's partitions into closed-open sets (see above), any spectral space is the inverse limit of it's partitions into constructible sets.

Definition 41. Suppose that X is a spectral space. The family of constructible sets of X is the smallest family of subsets closed under finite intersection, finite union, complement, and containing the quasi-compact open subsets of X .

⁸This follows from a partition of unity type argument: If $V(I_i)$ are the closed complements of the U_i , then $\cap V(I_i) = \emptyset$, but $\cap V(I_i) = V(\sum_i I_i)$ and this is empty if and only if $1 \in \sum_i I_i$, so $1 = a_{i_1} + \dots + a_{i_n}$ for some $a_{i_j} \in I_j$. But then $V(I_{i_1} + \dots + I_{i_n})$ is also empty, so $\{U_{i_1}, \dots, U_{i_n}\}$ is also an open cover.

— — — — *** add picture *** — — — —

Constructible sets generate a new topology on X : the *constructible topology*. The opens of the constructible topology are (possibly infinite) unions of constructible sets (which, of course, may not be open in the original topology). Since the original topology on our spectral space X is generated by quasi-compact opens, we see that the constructible topology is finer than the original topology. In fact, equipped with the constructible topology, X becomes a profinite set!

Proposition 42 ([Stacks Project, Tag 0901]). *Let X be a spectral space. The constructible topology on X is compact, Hausdorff, and totally disconnected. In other words, it is profinite.*

The claim about being totally disconnected is not in the Stacks Project statement, but follows easily from the fact that the constructible topology is Hausdorff, and that it has a basis consisting of sets which are both open and closed.

On the other hand, since the compact opens of a profinite set are precisely the closed opens, for a profinite set the constructible topology is the same as the original topology.

So changing a spectral spaces topology to the constructible topology is a kind of “profinitefication”.

Construction 43. Suppose that $X = \text{Spec}(R)$ is a spectral space and $X = \sqcup_{P \in \mathcal{P}} X_P$ is a decomposition into constructible subsets. Sending $x \in X$ to the index of the X_P that contains it defines a map $\pi : X \rightarrow P$. Then we give P the induced topology. So a subset $U \subset P$ is open if and only if $\pi^{-1}U$ is open.

Since the constructible open sets are the quasicompact ones, one sees that the map $X \rightarrow P$ is spectral. That is, there is a ring homomorphism $S \rightarrow R$ such that $\text{Spec}(S) \rightarrow \text{Spec}(R)$ induces the map of topological spaces $X \rightarrow P$ (this R may be different from the R in the construction, even though it gives rise to the same topological space).

Lemma 44. *Suppose that X is a spectral space. Then X is the inverse limit of its finite constructible decompositions $X = \varprojlim_{X = \sqcup_{P \in \mathcal{P}} X_P} P$.*

Proof. If we equip each P with the discrete topology, then we get the morphism $\pi : X \rightarrow \varprojlim_{X = \sqcup_{P \in \mathcal{P}} X_P} P$ from the profinite version of this lemma, which we have already proven is injective and surjective. So it suffices to show that $U \subseteq X$ is open if and only if $\pi(U)$ is open. Suppose U is a quasicompact open. Then U induces a constructible partition $X = U \amalg (X \setminus U)$ and so $\pi(U)$ is open. Since all opens of X are unions of quasicompact opens, this shows that if U is open, $\pi(U)$ is open. Conversely, if $\pi(U)$ is open, then U must be open by continuity and bijectivity of π . \square