

Corrections
Geometric Function Theory in Several Complex Variables
(Version, 1990)

- p. xi, \downarrow 3: §1 \Rightarrow §2
 p. 14, \uparrow 3: $\tilde{\gamma}_i \Rightarrow \dot{\tilde{\gamma}}_i$
 p. 15, \downarrow 2, 5, 6: $\tilde{\gamma}_i \Rightarrow \dot{\tilde{\gamma}}_i$
 p. 17, \downarrow 16: $F \Rightarrow G$
 p. 19, \uparrow 6: 0 Then \Rightarrow 0. Then
 p. 30, \uparrow 1: $F \Rightarrow L$
 p. 31, \downarrow 15: $B(3/4) \Rightarrow \overline{B(3/4)}$
 p. 32, \downarrow 3: $\Delta^* \Rightarrow \Delta^*(1)$
 p. 32, \downarrow 6: ' (2 places) \Rightarrow delete
 p. 32, \downarrow 13: min \Rightarrow delete
 p. 34, \downarrow 5: $h(z) \Rightarrow |h(z)|$ (2 places)
 p. 41, \downarrow 11: $U_\nu \Rightarrow V_\nu$
 p. 44, \downarrow 11: $\psi_i: (\overset{\wedge}{z_i^0}, \dots, z_i^i, \dots, z_i^m) \in \mathbf{C}^m \rightarrow \rho(z_i^0, \dots, \overset{i-\text{th}}{1}, \dots, z_i^m) \in U_i \Rightarrow$
 $\psi_i: \rho(\overset{\wedge}{z_i^1}, \dots, \overset{i-\text{th}}{1}, \dots, z_i^m) \in U_i \rightarrow (\overset{\wedge}{z_i^1}, \dots, z_i^i, \dots, z_i^m) \in \mathbf{C}^{m-1}$
 p. 44, \downarrow 12: $\overset{\wedge}{\mathbf{C}^m} \Rightarrow E$
 p. 44, \downarrow 14: $\mathbf{C}^m \Rightarrow \mathbf{C}^{m-1}$
 p. 44, \downarrow 15: $\overset{\wedge}{\mathbf{C}^m} \Rightarrow E$
 p. 44, \downarrow 18: $z_i^0 \Rightarrow z_i^1$
 p. 60, \downarrow 13: $\mathbf{1}_M \Rightarrow \mathbf{1}_U$
 p. 64, \downarrow 20: $H \Rightarrow H_i$
 p. 65, \uparrow 10: $- > \Rightarrow \rightarrow$
 p. 68, \downarrow 5: $X \Rightarrow x$
 p. 70, \downarrow 11: $= m \Rightarrow = 2m$
 p. 71, \uparrow 8: $\mathbf{K}_{\mathbf{C}^m} \Rightarrow \mathbf{K}(\mathbf{C}^m)$
 p. 72, \uparrow 11: $)^{k-m-1} \Rightarrow)^{d-m-1}$
 p. 72, \uparrow 11: $k \Rightarrow d$
 p. 72, \uparrow 11: $\{ \Rightarrow [$
 p. 72, \uparrow 11: $\} \Rightarrow]$
 p. 73, \uparrow 3: $\Omega \geq 0$ for $\Rightarrow \Omega(x) \geq 0$ for
 p. 76, \downarrow 4: $tr_m) \Rightarrow tr_m)(z)$
 p. 76, \downarrow 15: $-\partial\bar{\partial} \Rightarrow -i\partial\bar{\partial}$
 p. 76, \uparrow 9: $\Omega^m \Rightarrow \Omega$
 p. 78, \downarrow 13: $a(y) < b(y) \Rightarrow a(y) \leq b(y)$
 p. 79, \downarrow 8: $\Lambda^m \Rightarrow \Lambda$
 p. 79, \downarrow 12: $j = 1 \Rightarrow j = 2$
 p. 81, \downarrow 2: $\int_M \Rightarrow \int_{B'}$
 p. 81, \uparrow 9: $\sum_{j=1}^{\infty} \Psi_M \Rightarrow \sum_{j=1}^{\infty} \int_{f_j(E_j)} \Psi_M$
 p. 83, \downarrow 16: (5.4.1) \Rightarrow (5.5.1)
 p. 83, \downarrow 21: (2.1.8) \Rightarrow (2.1.22)
 p. 83, \downarrow 22: (2.1.9) \Rightarrow (2.1.23)
 p. 84, \downarrow 10: Put equation number (2.6.3).
 p. 86, \downarrow 15: (iv) \Rightarrow (d)
 p. 86, \downarrow 16: (3.3.43) \Rightarrow (3.3.44)
 p. 87, \downarrow 14: $\Phi \Rightarrow \Psi$
 p. 88, \uparrow 9: $\|f^*\xi\| \Rightarrow \|f^*\xi\|_N$
 p. 88, \uparrow 6: $\|\xi\| \Rightarrow \|\xi\|_M$
 p. 91, \downarrow 7: manifolds \Rightarrow spaces
 p. 91, \downarrow 12: of Lang \Rightarrow (delete)
 p. 91, \downarrow 12: Kähler \Rightarrow a Kähler manifold
 p. 91, \downarrow 13: compact \Rightarrow (delete)
 p. 91, \downarrow 13: Moisezon \Rightarrow Kähler manifold
 p. 99, \uparrow 1: $\|\phi\|_0 T(\phi) \Rightarrow \|\phi\|_0 T(\phi_A) \pm T(\phi)$.
 p. 111, \downarrow 4: **real current.** \Rightarrow **real current**, and $p = q$.
 p. 113, \downarrow 7: (3.2.14) \Rightarrow (3.1.14)
 p. 114, \downarrow 4: positive distributions \Rightarrow distributions of order 0
 p. 114, \downarrow 13: $\sigma \Rightarrow \sigma_k$
 p. 117, \uparrow 1: $\int_{\overline{B(r_2)} - \overline{B(r_1)}} \Rightarrow \int_{B(r_2) - \overline{B(r_1)}}$

- p. 118, \uparrow 5: $< T \implies \leq T$
 p. 120, \downarrow 11: (two places) $\frac{1}{r^{2k}} \implies$ (1'st) $\frac{1}{r_2^{2k}}$; (2'nd) $\frac{1}{r_1^{2k}}$
 p. 120, \uparrow 8: $< \implies \leq$
 p. 121, \uparrow 10: $\epsilon^{i\theta} \implies \epsilon e^{i\theta}$
 p. 122, \downarrow 12: *increasing* \implies *decreasing*
 p. 122, \downarrow 13: $u_j(z) \leq u_{j+1} \implies u_j(z) \geq u_{j+1}(z)$
 p. 122, \downarrow 13: $u_j \implies u_j(z)$
 p. 122, \uparrow 8: above \implies above on K
 p. 128, \uparrow 4: $[u] \implies [u_\epsilon]$
 p. 129, \downarrow 10, 12: (two places) $\leq \implies =$
 p. 130, \uparrow 8: $\Delta(1) \implies \Delta(R)$
 p. 140, \downarrow 11: $b_U \implies b_V$
 p. 144, \downarrow 15: $I_{M,x} \implies I_{X,x}$
 p. 145, \downarrow 13: (ii) \implies (iii)
 p. 148, \downarrow 12: $= m(\implies -m($
 p. 149, \uparrow 13: $f \implies F$
 p. 150, \downarrow 3: $a \implies b$
 p. 150, \uparrow 8: $e^{-u_i} \implies e^{-u_{(n)i}}$
 p. 151, \downarrow 7: $f^{-1}(x) \implies f^{-1}(f(x))$
 p. 152, \uparrow 13: $M \implies X$
 p. 154, \uparrow 3: $U \cap U_2 = \emptyset \implies U_1 \cap U_2 = \emptyset$
 p. 160, \downarrow 4: $\tilde{\mathbf{C}}^{n+1^{n+1}} \implies \tilde{\mathbf{C}}^{n+1}$
 p. 166, \downarrow 6: $V \implies A$
 p. 188, \downarrow 7: $B(r) \implies \overline{B(r)}$
 p. 206, \downarrow 2: (5.4.17) \implies (5.5.17)
 p. 207, \uparrow 1: Put equation number (5.5.22).
 p. 214, \uparrow 6: (5.4.6) \implies (5.5.46)
 p. 242, \downarrow 6: $X \implies Y$
 p. 244, \uparrow 7: (6.2.5) \implies (6.1.5)
 p. 247, \downarrow 6: states \implies stated
 p. 247, \uparrow 3: variety. \implies variety A .
 p. 247, \uparrow 2: *any* \implies *Any*
 p. 254, \uparrow 11: $u(x) \implies |u(x)|$
 p. 254, \uparrow 7: $u_r \implies \hat{u}_r$
 p. 254, \uparrow 5: $u^m \implies y^m$
 p. 255, \downarrow 2 \sim 4: $n \implies m$
 p. 256, \downarrow 2: $x \implies z$
 p. 258, \uparrow 12: $dz^j \wedge d\bar{z}^k \implies T_{jk} dz^j \wedge d\bar{z}^k$
 p. 260, \uparrow 6: $q(q; a, z) \implies e(q; a, z)$
 p. 261, \downarrow 9 \sim 10: $q, \implies q;$
 p. 261, \downarrow 11: ((\implies (\implies)
 p. 263, \uparrow 3: $r \rightarrow \infty \implies \nu \rightarrow \infty$
 p. 264, \uparrow 2 \sim 1: By \sim (iii), \implies By (5.2.25), Corollary (5.2.30), condition (iii) and Lemma (3.2),
 p. 265, \downarrow 1 \sim 2: Change these 2 lines by the following 2 lines:

$$T(r_\nu, A^{\pm 1}) \leq T(r_\nu, F) + T(r_\nu, G) + O(1) = o(r_\nu^{q+1}).$$

Thus Theorem (5.3.13) implies that $\log M(r_\nu, A^{\pm 1}) = o(r_\nu^{q+1})$.

- p. 265, \downarrow 4: $|\operatorname{Re} \log A(z)| \implies \sup\{|\operatorname{Re} \log A(z)|; z \in B(r_\nu)\}$
 p. 269, \uparrow 12 \sim 11: “Compact \sim (1987). \implies “A finiteness criterion for compact varieties of surjective holomorphic mappings,” Kodai Math. J. **13**, pp. 373-376 (1990).
 p. 269, \uparrow 8: K. \implies Y.