

ANNIHILATORS OF GENERALIZED VERMA MODULES OF THE SCALAR TYPE FOR CLASSICAL LIE ALGEBRAS

TOSHIO OSHIMA

Graduate School of Mathematical Sciences, University of Tokyo
Komaba 3-8-1, Tokyo 153-8914, Japan
e-mail: oshima@ms.u-tokyo.ac.jp

ABSTRACT. We construct a generator system of the annihilator of a generalized Verma module of a classical reductive Lie algebra induced from a character of a parabolic subalgebra as an analogue of the minimal polynomial of a matrix. In a classical limit it gives a generator system of the defining ideal of any semisimple co-adjoint orbit of the Lie algebra. We also give some applications to integral geometry.

1. INTRODUCTION

In [O3] generalized Capelli operators are defined in the universal enveloping algebra of $GL(n, \mathbb{C})$ and it is shown that they characterize the differential equations satisfied by the functions in degenerate principal series representations of $GL(n, \mathbb{R})$. The operators are used to formulate boundary value problems for various boundaries of the symmetric space $GL(n, \mathbb{R})/O(n)$ and to construct generalized hypergeometric equations related to Radon transformations on Grassmannian manifolds. In [O4] using these operators we construct a generator system of the annihilator of the generalized Verma module for $\mathfrak{gl}(n, \mathbb{C})$ induced from any character of any parabolic subalgebra.

In this paper the generator system is constructed for the classical Lie algebra \mathfrak{g} . Here \mathfrak{g} equals $\mathfrak{gl}(n, \mathbb{C})$, $\mathfrak{o}(2n, \mathbb{C})$, $\mathfrak{o}(2n+1, \mathbb{C})$ or $\mathfrak{sp}(n, \mathbb{C})$. In the case of $\mathfrak{gl}(n, \mathbb{C})$ the generator system in [O4] is an analogue of minors and elementary divisors. The generator system here is an analogue of the minimal polynomial of a matrix and different from the one constructed in [O4]. For the generator system of the center of the universal enveloping algebra the former corresponds to Capelli identity in [C1] and [C2] and the latter to the trace of the power of a matrix with components in the Lie algebra which is presented by [Ge].

In §2 we define a matrix F with components in \mathfrak{g} or the universal enveloping algebra $U(\mathfrak{g})$ associated to a finite dimensional representation of a Lie algebra \mathfrak{g} and define a *minimal polynomial* of F with respect to a \mathfrak{g} -module (cf. Definition 2.4).

In §3 we calculate the Harish-Chandra homomorphism of certain polynomials of F . It is a little complicated but elementary. A complete answer is given in Theorem 4.19 when \mathfrak{g} is $\mathfrak{gl}(n, \mathbb{C})$. Owing to this calculation, in §4 we introduce some polynomials of F and study their action on the generalized Verma module.

Then we construct a two-sided ideal of $U(\mathfrak{g})$ generated by the components $q(F)_{ij}$ for the minimal polynomial $q(x)$ of F and prove Theorem 4.4, which is the main result in this paper. It says (cf. Remark 4.5 ii)) that the ideal describes the *gap* between the generalized Verma module and the usual Verma module (cf. (5.1) and (5.7)) if at least the infinitesimal character of the Verma module is regular (resp. strongly regular) in the case when \mathfrak{g} equals $\mathfrak{gl}(n, \mathbb{C})$, $\mathfrak{o}(2n+1, \mathbb{C})$ or $\mathfrak{sp}(n, \mathbb{C})$

(resp. $\mathfrak{o}(2n, \mathbb{C})$). The main motivation to write this paper is to construct a two-sided ideal with this property originated in the problem in [O1].

It follows from this theorem that the ideal equals the annihilator of the generalized Verma module of the scalar type for the classical Lie algebra if at least the infinitesimal character is (strongly) regular (cf. Corollary 4.6).

We will use the homogenized universal enveloping algebra $U^\epsilon(\mathfrak{g})$ introduced in [O4] so that we can compare the generator system of a co-adjoint orbit in the dual of \mathfrak{g} . As a classical limit we get the generator system of any semisimple co-adjoint orbit for a classical Lie algebra, which is described in Theorem 4.11 (cf. Remark 4.12).

In §5 we show some applications of our two-sided ideals to integral transformations of sections of a line bundle over a generalized flag manifold. For example, Theorem 5.1 is a typical application, which shows that the system of differential equations defined by our two-sided ideal characterizes the image of the Poisson transform of the functions on any boundary of the Riemannian symmetric space of the non-compact type.

In §6 we discuss the infinitesimal character which is excluded in the results in §4 and present some problems.

In the subsequent paper [OO] we will give a simple explicit formula of minimal polynomials of generalized Verma modules of the scalar type for any reductive Lie algebra and study the same problem as in this paper.

In order to explain our idea, suppose $G = GL(2n, \mathbb{C})$ and put $A = \begin{pmatrix} \lambda I_n & 0 \\ B & \mu I_n \end{pmatrix} \in \mathfrak{g} = \text{Lie}(G)$. Here $\lambda, \mu \in \mathbb{C}$ and $B \in M(n, \mathbb{C})$ is a generic element. Note that A is conjugate to $\lambda I_n \oplus \mu I_n$ if $\lambda \neq \mu$ and to $\begin{pmatrix} \lambda & 0 \\ 1 & \lambda \end{pmatrix} \oplus \cdots \oplus \begin{pmatrix} \lambda & 0 \\ 1 & \lambda \end{pmatrix}$ otherwise. We will identify \mathfrak{g} and its dual \mathfrak{g}^* by the symmetric bilinear form $\langle X, Y \rangle = \text{Trace } XY$. Let $I_\Theta(\subset S(\mathfrak{g}))$ be the defining ideal of the closure \bar{V}_Θ of the conjugacy class $V_\Theta = \sum_{g \in G} \text{Ad}(g)A$ with $\text{Ad}(g)X = gXg^{-1}$.

Note that $I_\Theta = I_\Theta^0$ by denoting

$$\begin{cases} I_\Theta^\epsilon = \bigcap_{g \in G} \text{Ad}(g)J_\Theta^\epsilon, \\ J_\Theta^\epsilon = \sum_{X_1, X_2, X_3 \in M(n, \mathbb{C})} U^\epsilon(\mathfrak{g}) \left(\begin{pmatrix} X_1 & 0 \\ X_3 & X_2 \end{pmatrix} - \lambda \text{Trace } X_1 - \mu \text{Trace } X_2 \right). \end{cases}$$

Here $U^\epsilon(\mathfrak{g})$ is the quotient of the tensor algebra of \mathfrak{g} by the two-sided ideal generated by elements of the form $X \otimes Y - Y \otimes X - \epsilon[X, Y]$. Then $U^0(\mathfrak{g})$ is the symmetric algebra $S(\mathfrak{g})$ of \mathfrak{g} and $U^1(\mathfrak{g})$ is the universal enveloping algebra of \mathfrak{g} . We call a generalization of I_Θ^ϵ a quantization of I_Θ^0 and the quantization I_Θ^1 is nothing but the annihilator of the generalized Verma module $U(\mathfrak{g})/J_\Theta^1$.

Since $\text{rank}(X - \lambda I_{2n}) \leq n$ and $\text{rank}(X - \mu I_{2n}) \leq n$ for $X \in \bar{V}_\Theta$, the $(n+1)$ -minors ($\in S(\mathfrak{g})$) of $((E_{ij}) - \lambda I_{2n})$ and $((E_{ij}) - \mu I_{2n})$ are in I_Θ . On the contrary, they generate I_Θ if $\lambda \neq \mu$. The quantizations of the minors are generalized Capelli operators studied by [O3].

If $\lambda = \mu$, the derivatives of $(n+1)$ -minors of $((E_{ij}) - xI_{2n})$ at $x = \lambda$ are also in I_Θ and in general the generators are described by using the elementary divisors. In [O4], we define their quantizations, namely, we explicitly construct the corresponding generators for any generalized Verma module of the scalar type for $\mathfrak{gl}(n, \mathbb{C})$ using generalized Capelli operators and quantized elementary divisors. Moreover in [O4] we determine the condition that the annihilator determines the gap between the generalized Verma module and the usual Verma module. In the

example here, the equality

$$(1.1) \quad J_{\Theta}^{\epsilon} = I_{\Theta}^{\epsilon} + \sum_{i>j} U^{\epsilon}(\mathfrak{g})E_{ij} + \sum_{i=1}^n U^{\epsilon}(\mathfrak{g})(E_{ii} - \lambda) + \sum_{i=n+1}^{2n} U^{\epsilon}(\mathfrak{g})(E_{ii} - \mu)$$

holds if and only if $\lambda - \mu \notin \{\epsilon, 2\epsilon, \dots, (n-1)\epsilon\}$. When $\epsilon = 1$, this condition is satisfied if $U(\mathfrak{g})/J_{\Theta}^{\epsilon}$ has a regular infinitesimal character, which is equivalent to the condition that $\lambda - \mu \notin \{1, 2, \dots, 2n-1\}$. If (1.1) holds, the quantized generators are considered to be the differential equations which characterize the representations of the group G related to the generalized Verma module. Hence they are important and the motivation of our study in this note is this fact.

Now since $(x - \lambda)(x - \mu)$ is the minimal polynomial of A , all the components of $((E_{ij} - \lambda I_{2n})((E_{ij} - \mu I_{2n}))$ are in J_{Θ}^0 . They generate I_{Θ}^0 together with $\sum_{i=1}^{2n} E_{ii} - n\lambda - n\mu$ if $\lambda \neq \mu$. We can quantize this minimal polynomial and the quantized minimal polynomial in this example equals $q^{\epsilon}(x) = (x - \lambda)(x - \mu - n\epsilon)$. We can show that the $4n^2$ components of the matrix $q^{\epsilon}((E_{ij})) \in M(2n, U^{\epsilon}(\mathfrak{g}))$ and the element $\sum_{i=1}^{2n} E_{ii} - n\lambda - n\mu$ generate I_{Θ}^{ϵ} if $\lambda - \mu \notin \{\epsilon, 2\epsilon, \dots, (n-1)\epsilon\}$. When we identify $U(\mathfrak{g})$ with the ring of left invariant holomorphic differential operators on $GL(n, \mathbb{C})$, we have $q^1((E_{ij})) = ({}^t X \partial - \lambda)({}^t X \partial - \mu - n)$ with the matrices $X = (x_{ij}) \in GL(n, \mathbb{C})$ and $\partial = (\frac{\partial}{\partial x_{ij}})$.

The main topic in this paper is to construct the elements in $U(\mathfrak{g})$ which kills the generalized Verma module of the scalar type for the classical Lie algebra by using the quantized minimal polynomial.

The author expresses his sincere gratitude to Mittag-Leffler Institute. The result in this paper for $\mathfrak{g} = \mathfrak{gl}(n, \mathbb{C})$ was obtained when the author was invited there from September until November in 1995 and it is reported in [O2].

2. MINIMAL POLYNOMIALS

For a module \mathfrak{A} and positive integers N and N' , we denote by $M(N, N', \mathfrak{A})$ the set of matrices of size $N \times N'$ with components in \mathfrak{A} . If $N = N'$, we simply denote it by $M(N, \mathfrak{A})$ and then $M(N, \mathfrak{A})$ is naturally an associative algebra if so is \mathfrak{A} .

We use the standard notation \mathfrak{gl}_n , \mathfrak{o}_n and \mathfrak{sp}_n for classical Lie algebras over \mathbb{C} . For a Lie algebra \mathfrak{g} we denote by $U(\mathfrak{g})$ and $S(\mathfrak{g})$ the universal enveloping algebra and the symmetric algebra of \mathfrak{g} , respectively. For a non-negative integer k let $S(\mathfrak{g})^{(k)}$ be the subspace of $S(\mathfrak{g})$ formed by elements of degree at most k . If we fix a Poincare-Birkhoff-Witt base of $U(\mathfrak{g})$, we can identify $U(\mathfrak{g})$ and $S(\mathfrak{g})$ as vector spaces and we denote by $U(\mathfrak{g})^{(k)}$ the subspace of $U(\mathfrak{g})$ corresponding to $S(\mathfrak{g})^{(k)}$.

The Lie algebra \mathfrak{gl}_N is identified with $M(N, \mathbb{C}) \simeq \text{End}(\mathbb{C}^N)$ by $[X, Y] = XY - YX$. Let $E_{ij} = \left(\delta_{\mu i} \delta_{\nu j} \right)_{\substack{1 \leq \mu \leq N \\ 1 \leq \nu \leq N}} \in M(N, \mathbb{C})$ be the standard matrix units. Note that the symmetric bilinear form

$$(2.1) \quad \langle X, Y \rangle = \text{Trace } XY \quad \text{for } X, Y \in \mathfrak{gl}_N$$

on \mathfrak{gl}_N is non-degenerate and satisfies

$$\begin{aligned} \langle E_{ij}, E_{\mu\nu} \rangle &= \delta_{i\nu} \delta_{j\mu}, \\ X &= \sum_{i,j} \langle X, E_{ji} \rangle E_{ij}, \end{aligned}$$

$$\langle \text{Ad}(g)X, \text{Ad}(g)Y \rangle = \langle X, Y \rangle \quad \text{for } X, Y \in \mathfrak{gl}_N \text{ and } g \in GL(N, \mathbb{C}).$$

Lemma 2.1. *Let \mathfrak{g} be a Lie algebra over \mathbb{C} and let (π, \mathbb{C}^N) be a representation of \mathfrak{g} . We denote by $U(\pi(\mathfrak{g}))$ the subalgebra of the universal enveloping algebra $U(\mathfrak{gl}_N)$*

of \mathfrak{gl}_N generated by $\pi(\mathfrak{g})$. Let p be a linear map of \mathfrak{gl}_N to $U(\pi(\mathfrak{g}))$ satisfying

$$(2.2) \quad p([X, Y]) = [X, p(Y)] \quad \text{for } X \in \pi(\mathfrak{g}) \text{ and } Y \in \mathfrak{gl}_N,$$

that is, $p \in \text{Hom}_{\pi(\mathfrak{g})}(\mathfrak{gl}_N, U(\pi(\mathfrak{g})))$.

Fix $f(x) \in \mathbb{C}[x]$ and put

$$(2.3) \quad \begin{cases} F = \left(p(E_{ij}) \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} \in M(N, U(\pi(\mathfrak{g}))), \\ \left(Q_{ij} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = f(F) \in M(N, U(\pi(\mathfrak{g}))). \end{cases}$$

Then

$$(2.4) \quad \left(p(\text{Ad}(g)E_{ij}) \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = {}^t g F {}^t g^{-1} \quad \text{for } g \in GL(n, \mathbb{C})$$

and

$$(2.5) \quad \begin{aligned} [X, Q_{ij}] &= \sum_{\nu=1}^N X_{\nu i} Q_{\nu j} - \sum_{\nu=1}^N X_{j \nu} Q_{i \nu} \\ &= \sum_{\nu=1}^N \langle X, Q_{i \nu} \rangle Q_{\nu j} - \sum_{\nu=1}^N Q_{i \nu} \langle X, Q_{\nu j} \rangle \quad \text{for } X = \left(X_{ij} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} \in \pi(\mathfrak{g}) \end{aligned}$$

with $X_{ij} \in \mathbb{C}$.

Proof. Put $g = (g_{ij})$ and $g^{-1} = (g'_{ij})$. Then

$$\left(\text{Ad}(g)E_{ij} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = \left(\sum_{\mu, \nu} g_{\mu i} g'_{j \nu} E_{\mu \nu} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = {}^t g F {}^t g^{-1}.$$

Fix $X \in \pi(\mathfrak{g})$. Since

$$[X, E_{ij}] = \left[\sum_{\mu, \nu} X_{\mu \nu} E_{\mu \nu}, E_{ij} \right] = \sum_{\mu=1}^N X_{\mu i} E_{\mu j} - \sum_{\nu=1}^N X_{j \nu} E_{i \nu},$$

we have (2.5) for $f(x) = x$ by (2.2).

Suppose (Q_{ij}^1) and $(Q_{ij}^2) \in M(N, U(\pi(\mathfrak{g})))$ satisfy (2.5). Put $Q_{ij}^3 = \sum_{k=1}^N Q_{ik}^1 Q_{kj}^2$ in $U(\pi(\mathfrak{g}))$. Then

$$\begin{aligned} [X, Q_{ij}^3] &= \sum_{k=1}^N [X, Q_{ik}^1] Q_{kj}^2 + \sum_{k=1}^N Q_{ik}^1 [X, Q_{kj}^2] \\ &= \sum_{k=1}^N \left(\sum_{\mu=1}^N X_{\mu i} Q_{\mu k}^1 Q_{kj}^2 - \sum_{\nu=1}^N X_{k \nu} Q_{i \nu}^1 Q_{kj}^2 \right) \\ &\quad + \sum_{k=1}^N \left(\sum_{\mu=1}^N Q_{ik}^1 X_{\mu k} Q_{\mu j}^2 - \sum_{\nu=1}^N Q_{ik}^1 X_{j \nu} Q_{k \nu}^2 \right) \\ &= \sum_{\mu=1}^N X_{\mu i} Q_{\mu j}^3 - \sum_{\nu=1}^N X_{j \nu} Q_{i \nu}^3 \end{aligned}$$

and therefore the elements (Q_{ij}^3) of $M(N, U(\pi(\mathfrak{g})))$ satisfying (2.5) form a subalgebra of $M(N, U(\pi(\mathfrak{g})))$. \square

Definition 2.2. If the symmetric bilinear form (2.1) is non-degenerate on $\pi(\mathfrak{g})$, the orthogonal projection of \mathfrak{gl}_N onto $\pi(\mathfrak{g})$ satisfies the assumption for p in Lemma 2.1, which we call the *canonical projection* of \mathfrak{gl}_N to $\pi(\mathfrak{g})$.

Remark 2.3. Suppose that \mathfrak{g} is reductive. Let G be a connected and simply connected Lie group with the Lie algebra \mathfrak{g} and let G_U be a maximal compact subgroup of G . Assume that the finite dimensional representation (π, V) in Lemma 2.1 can be lifted to the representation of G_U . Let $\mathfrak{g} = \bar{\mathfrak{n}} \oplus \mathfrak{a} \oplus \mathfrak{n}$ be a triangular decomposition of \mathfrak{g} such that $\exp \mathfrak{a} \cap G_U$ is a maximal torus of G_U . Let $\Sigma(\mathfrak{a})$ and $\Sigma(\mathfrak{a})^+$ be the sets of the roots for the pair $(\mathfrak{g}, \mathfrak{a})$ and $(\mathfrak{n}, \mathfrak{a})$, respectively, and let $\Psi(\mathfrak{a})$ denote the fundamental system of $\Sigma(\mathfrak{a})^+$. We fix a Hermitian inner product on V so that π is a unitary representation of G_U . Moreover let $\{v_1, \dots, v_N\}$ be an orthonormal basis of V such that v_j is a weight vector of a weight ϖ_j with respect to the Cartan subalgebra \mathfrak{a} . We may assume that $\varpi_i - \varpi_j \in \Sigma(\mathfrak{a})^+$ means $i > j$. Hence ϖ_1 is the lowest weight and ϖ_N is the highest weight of the representation π . Under this basis we identify $\pi(X) = (\pi(X)_{ij}) \in M(N, \mathbb{C}) \simeq \text{End}(\mathbb{C}^N) \simeq \mathfrak{gl}_N$ for $X \in \mathfrak{g}$ by $\pi(X)v_j = \sum_{i=1}^N \pi(X)_{ij}v_i$. Note that $\pi(\mathfrak{a}) \subset \mathfrak{a}_N$, $\pi(\mathfrak{n}) \subset \mathfrak{n}_N$ and $\pi(\bar{\mathfrak{n}}) \subset \bar{\mathfrak{n}}_N$ by denoting

$$(2.6) \quad \mathfrak{a}_N = \sum_{j=1}^N \mathbb{C}E_{jj}, \quad \mathfrak{n}_N = \sum_{1 \leq j < i \leq N} \mathbb{C}E_{ij} \quad \text{and} \quad \bar{\mathfrak{n}}_N = \sum_{1 \leq i < j \leq N} \mathbb{C}E_{ij}.$$

Since $\pi(X)$ is skew Hermitian for the element X in the Lie algebra \mathfrak{g}_U of G_U and $\mathbb{C}\pi(\mathfrak{g}_U) = \pi(\mathfrak{g})$, we have ${}^t\overline{\pi(\mathfrak{g})} = \pi(\mathfrak{g})$. Hence the symmetric bilinear form (2.1) is non-degenerate on $\pi(\mathfrak{g})$ and there exists the canonical projection of \mathfrak{gl}_N to $\pi(\mathfrak{g})$.

Definition 2.4 (Characteristic polynomials and minimal polynomials). Given a Lie algebra \mathfrak{g} , a faithful finite dimensional representation (π, \mathbb{C}^N) and a \mathfrak{g} -homomorphism p of $\text{End}(\mathbb{C}^N) \simeq \mathfrak{gl}_N$ to $U(\mathfrak{g})$. Here we identify \mathfrak{g} as a subalgebra of \mathfrak{gl}_N through π . Let $U(\mathfrak{g})$ and $Z(\mathfrak{g})$ be the universal enveloping algebra of \mathfrak{g} and the center of $U(\mathfrak{g})$, respectively. For $F = (p(E_{ij})) \in M(N, U(\mathfrak{g}))$, we say $q_F(x) \in U(\mathfrak{g})^G[x]$ is the *characteristic polynomial* of F if it is a non-zero polynomial of x satisfying

$$q_F(F) = 0$$

with the minimal degree.

Suppose moreover a \mathfrak{g} -module M is given. Then we call $q_{F,M}(x) \in \mathbb{C}[x]$ is the *minimal polynomial* of F with respect to M if it is the monic polynomial with the minimal degree which satisfies

$$q_{F,M}(F)M = 0.$$

If p is the canonical projection in Definition 2.2, we sometimes denote F_π , q_π and $q_{\pi,M}$ in place of F , q_F and $q_{F,M}$, respectively.

Remark 2.5. i) After the results in this paper was obtained, the author was informed that [Go2] studied the characteristic polynomial of F_π for the irreducible representation π of the reductive Lie algebra.

ii) If \mathfrak{g} is reductive, the characteristic polynomial is uniquely determined by (π, p) up to a constant multiple of the element of $Z(\mathfrak{g})$ since $Z(\mathfrak{g})$ is an integral domain.

iii) If \mathfrak{g} is reductive and M has an infinitesimal character χ , that is, χ is an algebra homomorphism of $Z(\mathfrak{g})$ to \mathbb{C} with $(D - \chi(D))M = 0$ for $D \in Z(\mathfrak{g})$, then $\chi(q_F(x)) \in \mathbb{C}[x]q_{F,M}(x)$.

iv) The characteristic polynomial and minimal polynomial of a matrix in the linear algebra can be regarded as a classical limit of our definition. See the proof of Proposition 4.16.

Theorem 2.6. *Let \mathfrak{g} be a reductive Lie algebra and let F be a matrix of $U(\mathfrak{g})$ defined from a representation of π under Definition 2.4.*

i) There exists the characteristic polynomial $q_F(x)$ whose degree is not larger than $\sum_{\varpi} m_{\pi}(\varpi)^2$. Here ϖ runs through the weights of π and $m_{\pi}(\varpi)$ denotes the multiplicity of the generalized weight ϖ in π .

ii) The minimal polynomial $q_{F,M}(x)$ exists if a \mathfrak{g} -module M has a finite length or an infinitesimal character. Its degree is not larger than that of the characteristic polynomial $q_F(x)$ if M has an infinitesimal character.

Proof. Let $\hat{Z}(\mathfrak{g})$ denote the quotient field of $Z(\mathfrak{g})$ and put $\hat{U}(\mathfrak{g}) = \hat{Z}(\mathfrak{g}) \otimes_{Z(\mathfrak{g})} U(\mathfrak{g})$. Owing to [Ko] it is known that $U(\mathfrak{g}) = \Lambda(H(\mathfrak{g})) \otimes Z(\mathfrak{g})$, where $H(\mathfrak{g})$ is the space of \mathfrak{g} -harmonic polynomials of $S(\mathfrak{g})$ and Λ is the map of the symmetrization of $S(\mathfrak{g})$ onto $U(\mathfrak{g})$. It is also known that $H(\mathfrak{g}) \simeq \sum_{\tau \in \hat{\mathfrak{g}}_f} m_{\tau}(0)\tau$ as a representation space of \mathfrak{g} by denoting $\hat{\mathfrak{g}}_f$ the equivalence classes of the finite dimensional irreducible representations of \mathfrak{g} .

Hence the dimension of the \mathfrak{g} -homomorphisms of $\pi \otimes \pi^*$ to $\hat{U}(\mathfrak{g})$ over the field $\hat{Z}(\mathfrak{g})$ is not larger than $\sum_{\tau \in \hat{\mathfrak{g}}_f} [\pi \otimes \pi^*, \tau] m_{\tau}(0)$. Here $[\pi \otimes \pi^*, \tau]$ is the multiplicity of τ appeared in $[\pi \otimes \pi^*]$ in the sense of the Grothendieck group. Moreover it is clear that $\sum_{\tau \in \hat{\mathfrak{g}}_f} [\pi \otimes \pi^*, \tau] m_{\tau}(0) = m_{\pi \otimes \pi^*}(0) = \sum_{\varpi} m_{\pi}(\varpi)^2$. On the other hand Lemma 2.1 says that the space $V_k = \sum_{i,j} \mathbb{C}F_{ij}^k$ is naturally a subrepresentation of the representation of \mathfrak{g} which is realized in $M(N, \mathbb{C})$ and belongs to $\pi \otimes \pi^*$ and that the map $T_k : E_{ij} \mapsto F_{ij}^k$ defines a \mathfrak{g} -homomorphism of $M(N, \mathbb{C})$ to $U(\mathfrak{g})$. Hence T_1, \dots, T_m are linearly dependent over $\hat{Z}(\mathfrak{g})$ if $m > \sum_{\varpi} m_{\pi}(\varpi)^2$. Thus we have proved the existence of the characteristic polynomial with the required degree.

For the existence of the minimal polynomial it is sufficient to prove the existence of a non-zero polynomial $f(x)$ with $f(F)M = 0$. Considering the irreducible subquotients of M in Definition 2.4, we may assume M has an infinitesimal character λ . Let $q_F(x)$ be the characteristic polynomial. We can choose $\mu \in \mathfrak{a}^*$ so that $\bar{\omega}(q_F(x))(\lambda + \mu t) \in \mathbb{C}[x, t]$ is not zero. We can find a non-negative integer k such that $f(x, t) = t^{-k} \bar{\omega}(q_F(x)) \in \mathbb{C}[x, t]$ and $f(x, 0)$ is not zero. Here $\bar{\omega}$ is the Harish-Chandra homomorphism defined by (4.10). Put $I_{\lambda} = \sum_{Z \in Z(\mathfrak{g})} U(\mathfrak{g})(Z - \bar{\omega}(Z)(\lambda))$. We define $h(t) \in M(N, H(\mathfrak{g}) \otimes \mathbb{C}[t])$ so that $f(F, t) - \Lambda(h(t)) \in M(N, I_{\lambda + \mu t})$. Since $d_F(F)(\lambda + \mu t) \in M(N, I_{\lambda + \mu t})$, $h(t) = 0$ for $t \neq 0$ and hence $h = 0$ and therefore $f(F, 0)(U(\mathfrak{g})/I_{\lambda}) = 0$. Hence $f(F, 0)M = 0$ because $\text{Ann}(M) \supset I_{\lambda}$. \square

Hereafter in this note we assume

$$(2.7) \quad \begin{cases} \pi \text{ is injective,} \\ p(\mathfrak{gl}_N) \subset \mathfrak{g}, \\ p(X) = CX \quad \text{for } X \in \mathfrak{g} \end{cases}$$

in Lemma 2.1 with a suitable non-zero constant C . Then we have the following.

Remark 2.7. i) Since π is faithful, \mathfrak{g} is identified with the Lie subalgebra $\pi(\mathfrak{g})$ of \mathfrak{gl}_N and $U(\pi(\mathfrak{g}))$ is identified with the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} . We note that the existence of p with (2.7) is equivalent to the existence of a \mathfrak{g} -invariant subspace of \mathfrak{gl}_N complementary to \mathfrak{g} .

ii) Fix $g \in GL(N, \mathbb{C})$. If we replace (π, \mathbb{C}^N) by (π^g, \mathbb{C}^N) with $\pi^g(X) = \text{Ad}(g)\pi(X)$ for $X \in \mathfrak{g}$ in Lemma 2.1, $(F_{ij}) \in M(N, \mathfrak{g})$ naturally changes into ${}^t g^{-1}(F_{ij}){}^t g$ and therefore the corresponding characteristic polynomial and minimal polynomial does not depend on the realization of the representation π . In fact, the map p^g of $U(\mathfrak{g})$ to $\pi(\mathfrak{g})$ is naturally defined by $\pi^g(X) = \text{Ad}(g)(p(\text{Ad}(g)^{-1}X))$ and hence $p^g(E) = \text{Ad}(g)(p(\text{Ad}(g)^{-1}E)) = \text{Ad}(g)(p({}^t g^{-1}E{}^t g)) = \pi^g({}^t g^{-1}(F_{ij}){}^t g)$.

iii) Suppose \mathfrak{g} is semisimple. Then the existence of p is clear because any finite dimensional representation of \mathfrak{g} satisfies the assumption in Remark 2.3.

iv) Let σ be an involutive automorphism of \mathfrak{gl}_N . Put

$$\mathfrak{g} = \{X \in \mathfrak{gl}_N; \sigma(X) = X\}.$$

Let π be the inclusion map of $\mathfrak{g} \subset \mathfrak{gl}_N$. Since $\mathfrak{q} = \{X \in \mathfrak{gl}_N; \sigma(X) = -X\}$ is \mathfrak{g} -stable, we may put $p(X) = \frac{X + \sigma(X)}{2}$ in Lemma 2.1, which is the canonical projection with respect to the bilinear form of \mathfrak{gl}_N .

v) For a positive integer k and complex numbers $\lambda_1, \dots, \lambda_k$, the vector space spanned by the N^2 components of the matrix $(p(E) - \lambda_1 I_N) \cdots (p(E) - \lambda_k I_N)$ is $\text{ad}(\mathfrak{g})$ -invariant. Moreover the trace of the matrix is a central element of $U(\mathfrak{g})$, which is clear from Lemma 2.1 and studied by [Ge] and [Gol] etc.

3. PROJECTION TO THE CARTAN SUBALGEBRA

Now we consider the natural realization of classical simple Lie algebras. Denoting

$$\tilde{I}_n = \left(\delta_{i,n+1-j} \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}} = \begin{pmatrix} & & 1 \\ & \cdots & \\ 1 & & \end{pmatrix} \text{ and } \tilde{J}_n = \begin{pmatrix} & \tilde{I}_n \\ -\tilde{I}_n & \end{pmatrix},$$

we naturally identify

$$(3.1) \quad \begin{aligned} \mathfrak{o}_n &= \{X \in \mathfrak{gl}_n; \sigma_{\mathfrak{o}_n}(X) = X\} \quad \text{with } \sigma_{\mathfrak{o}_n}(X) = -\tilde{I}_n^t X \tilde{I}_n, \\ \mathfrak{sp}_n &= \{X \in \mathfrak{gl}_{2n}; \sigma_{\mathfrak{sp}_n}(X) = X\} \quad \text{with } \sigma_{\mathfrak{sp}_n}(X) = -\tilde{J}_n^t X \tilde{J}_n. \end{aligned}$$

Definition 3.1. Let $\mathfrak{g} = \mathfrak{gl}_n$ or \mathfrak{o}_{2n} or \mathfrak{o}_{2n+1} or \mathfrak{sp}_n and put $N = n$ or $2n$ or $2n+1$ or $2n$, respectively, so that \mathfrak{g} is a subalgebra of \mathfrak{gl}_N . Put

$$(3.2) \quad \bar{i} = N + 1 - i$$

for any integer i and define

$$(3.3) \quad \epsilon_i = \begin{cases} 0 & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ 1 & \text{if } \mathfrak{g} = \mathfrak{o}_N, \\ 1 & \text{if } \mathfrak{g} = \mathfrak{sp}_n \text{ and } i \leq n, \\ -1 & \text{if } \mathfrak{g} = \mathfrak{sp}_n \text{ and } i > n. \end{cases}$$

Then the involutions $\sigma_{\mathfrak{g}}$ of \mathfrak{gl}_N defining \mathfrak{g} with $\mathfrak{g} = \mathfrak{o}_N$ and \mathfrak{sp}_n satisfy

$$\sigma_{\mathfrak{g}}(E_{ij}) = -\epsilon_i \epsilon_j E_{\bar{j}\bar{i}}.$$

We moreover define

$$(3.4) \quad F = \left(F_{ij} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} = \left(E_{ij} - \epsilon_i \epsilon_j E_{\bar{j}\bar{i}} \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}}.$$

This definition of F means $C = 2$ in (2.7) if $\mathfrak{g} = \mathfrak{o}_N$ or \mathfrak{sp}_n . We will denote F_i in place of F_{ii} for simplicity. Then $\mathfrak{g} = \sum_{i,j} \mathbb{C} F_{ij}$ and

$$(3.5) \quad [X, F_{ij}] = \sum_{\nu=1}^N (X_{\nu i} F_{\nu j} - X_{j \nu} F_{i \nu}) \quad \text{for } X = (X_{ij}) \in \mathfrak{g} \subset M(N, \mathbb{C})$$

by Lemma 2.1.

Use the notation (2.6) and define $\mathfrak{a} = \mathfrak{g} \cap \mathfrak{a}_N$, $\mathfrak{n} = \mathfrak{g} \cap \mathfrak{n}_N$ and $\bar{\mathfrak{n}} = \mathfrak{g} \cap \bar{\mathfrak{n}}_N$. Then

$$(3.6) \quad \mathfrak{g} = \bar{\mathfrak{n}} \oplus \mathfrak{a} \oplus \mathfrak{n}$$

is a triangular decomposition of \mathfrak{g} .

Definition 3.2. For a positive integer k and complex numbers $\lambda_1, \dots, \lambda_k$ put

$$F^k(\lambda_1, \dots, \lambda_k) = (F - \lambda_1 I_N) \cdots (F - \lambda_k I_N)$$

and define an element $\bar{F}^k(\lambda_1, \dots, \lambda_k)$ in $M(N, U(\mathfrak{a}))$ by

$$(3.7) \quad F^k(\lambda_1, \dots, \lambda_k) \equiv \bar{F}^k(\lambda_1, \dots, \lambda_k) \pmod{M(N, \bar{\mathfrak{n}}U(\mathfrak{g}) + U(\mathfrak{g})\mathfrak{n})}$$

In this section we will study the image $\bar{F}^k(\lambda_1, \dots, \lambda_k)$ of $F^k(\lambda_1, \dots, \lambda_k)$ under the Harish-Chandra homomorphism with respect to (3.6). First we note that if

$$(3.8) \quad F_{ij} \in \begin{cases} \bar{\mathfrak{n}} & \text{if } i < j, \\ \mathfrak{a} & \text{if } i = j, \\ \mathfrak{n} & \text{if } i > j, \end{cases}$$

we have

$$(3.9) \quad \begin{aligned} F_{ij}^k(\lambda_1, \dots, \lambda_k) &\equiv \sum_{\mu=1}^j F_{i\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_{\mu j} - \lambda_k \delta_{\mu j}) \pmod{U(\mathfrak{g})\mathfrak{n}} \\ &= F_{ij}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_j - \lambda_k) \\ &\quad + \sum_{\mu=1}^{j-1} \left(F_{\mu j} F_{i\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) - [F_{\mu j}, F_{i\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1})] \right) \\ &= F_{ij}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_j - \lambda_k) + \sum_{\mu=1}^{j-1} \left(F_{\mu j} F_{i\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \right. \\ &\quad \left. - \sum_{\nu=1}^{i-1} \langle F_{\mu j}, E_{i\nu} \rangle F_{\nu\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) + \sum_{\nu=\mu+1}^N \langle F_{\mu j}, E_{\nu\mu} \rangle F_{i\nu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \right) \end{aligned}$$

by Lemma 2.1.

The following is clear by the induction on k .

Remark 3.3. i) The highest homogeneous part of $\bar{F}^k(\lambda_1, \dots, \lambda_k)$ with the degree k is given by

$$\bar{F}^k(\lambda_1, \dots, \lambda_k) \equiv \left(\delta_{ij} F_{ii}^k \right)_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}} \pmod{M(N, U(\mathfrak{a})^{(k-1)}}.$$

ii) If $\mathfrak{g} = \mathfrak{gl}_n$ or \mathfrak{o}_{2n+1} or \mathfrak{sp}_n and π is the natural representation of \mathfrak{g} , it is clear that $\text{Trace } F_{\pi}^k$ for $k = 1, 2, \dots, n$ or $k = 2, 4, \dots, 2n$ or $2, 4, \dots, 2n$, respectively, generate $Z(\mathfrak{g})$ as an algebra. In particular for any $D \in Z(\mathfrak{g})$ there uniquely exists a polynomial $f(x)$ with $\text{Trace } f(F) = D$. In the case when $\mathfrak{g} = \mathfrak{o}_{2n}$ we use both the natural representation π and the half-spin representation π' of \mathfrak{g} and then $\text{Trace } F_{\pi}^k$ for $k = 2, 4, \dots, 2(n-1)$ and $\text{Trace } F_{\pi'}^n$ generate $Z(\mathfrak{g})$.

iii) The Killing form of \mathfrak{g} is a positive constant multiple of the restriction of the bilinear form (2.1) to \mathfrak{g} if \mathfrak{g} is simple.

Hereafter suppose that $\mathfrak{g} = \mathfrak{gl}_n$ or \mathfrak{o}_{2n} or \mathfrak{o}_{2n+1} or \mathfrak{sp}_n and that F is given by (3.4). Then (3.4) means

$$\langle F_{\mu j}, E_{i\nu} \rangle = \delta_{ij} \delta_{\mu\nu} - \epsilon_{\mu} \epsilon_j \delta_{\bar{\mu}i} \delta_{\bar{j}\nu} \quad \text{and} \quad \langle F_{\mu j}, E_{\mu\nu} \rangle = \delta_{j\nu} - \epsilon_{\mu} \epsilon_j \delta_{\bar{\mu}\nu} \delta_{\bar{j}\mu}$$

and therefore it follows from (3.9) that

$$(3.10) \quad \begin{aligned} F_{ij}^k(\lambda_1, \dots, \lambda_k) &\equiv F_{ij}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_j - \lambda_k + j - 1) \\ &\quad + \sum_{\mu=1}^{j-1} \left(F_{\mu j} F_{i\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \right. \\ &\quad \left. - \delta_{ij} F_{\mu\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) + \epsilon_{\mu} \epsilon_j \delta_{\bar{\mu}\bar{i}} F_{\bar{j}\bar{i}}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \right. \\ &\quad \left. - \epsilon_{\mu} \epsilon_j \delta_{\mu\bar{j}} F_{i\bar{j}}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \right) \pmod{U(\mathfrak{g})\mathfrak{n}}. \end{aligned}$$

Since $[U(\mathfrak{g}\mathfrak{n}, U(\mathfrak{a})) \subset U(\mathfrak{g}\mathfrak{n})$ and since $F_{ij} \in \mathfrak{n}$ and $F_{\bar{j}\bar{i}} \in \mathfrak{n}$ for $i > j$, the equation (3.10) shows $F_{ij}^k(\lambda_1, \dots, \lambda_k) \equiv 0 \pmod{U(\mathfrak{g}\mathfrak{n})}$ for $i > j$ by the induction on k . Similarly we have $F_{\mu\nu}^k(\lambda_1, \dots, \lambda_k) \in \bar{\mathfrak{n}}U(\mathfrak{g})$ if $i < j$. Hence by denoting

$$(3.11) \quad \begin{aligned} \omega_i &= \begin{cases} 0 & \text{if } i \leq n, \\ \epsilon_i & \text{if } i > n, \end{cases} \\ \omega'_j &= \begin{cases} 0 & \text{if } j \leq n \text{ or } \bar{j} \geq j, \\ \epsilon_j & \text{if } j > n \text{ and } \bar{j} < j, \end{cases} \end{aligned}$$

we have

$$(3.12) \quad \begin{aligned} F_{ii}^k(\lambda_1, \dots, \lambda_k) &\equiv F_{ii}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_i - \lambda_k + i - 1 - \omega_i) + \omega_i F_{ii}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \\ &\quad - \sum_{\mu=1}^{i-1} F_{\mu\mu}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \pmod{U(\mathfrak{g}\mathfrak{n})}, \end{aligned}$$

$$(3.13) \quad \begin{aligned} F_{i+1}^k(\lambda_1, \dots, \lambda_k) &\equiv F_{i+1}^{k-1}(\lambda_1, \dots, \lambda_{k-1})(F_{i+1} - \lambda_k + i - \omega'_{i+1}) + \omega_i F_{i-1}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \\ &\quad + F_{i+1} F_{ii}^{k-1}(\lambda_1, \dots, \lambda_{k-1}) \pmod{U(\mathfrak{g}\mathfrak{n})}. \end{aligned}$$

Now we give the main result in this section:

Proposition 3.4. *Suppose that $\mathfrak{g} = \mathfrak{gl}_n$ or \mathfrak{o}_{2n} or \mathfrak{o}_{2n+1} or \mathfrak{sp}_n and that F is given by (3.4). Let $\Theta = \{n_1 < n_2 < \dots < n_L = n\}$ be a sequence of positive integers. Put $n'_\nu = n_\nu - n_{\nu-1}$ for $\nu = 1, \dots, L$ with $n_0 = 0$ and fix a positive number k . Let μ_1, \dots, μ_k be complex numbers. Put $n_0 = 0$ and $n_\nu = n$ for $\nu > L$ and define*

$$\begin{aligned} \iota_\Theta(\nu) &= p \quad \text{if } n_{p-1} < \nu \leq n_p, \\ \tilde{J}(\mu)_i &= U(\mathfrak{g}\mathfrak{n}) + \sum_{\nu=1}^i U(\mathfrak{g})(F_\nu - \mu_{\iota_\Theta(\nu)} + n_{\iota_\Theta(\nu)-1}). \end{aligned}$$

If $\mathfrak{g} = \mathfrak{gl}_n$, we put $H(\Theta, \mu_1, \dots, \mu_L) = F^L(\mu_1, \dots, \mu_L)$.

If $\mathfrak{g} = \mathfrak{sp}_n$ or \mathfrak{o}_{2n} , we put

$$\begin{aligned} H(\Theta, \mu_1, \dots, \mu_L) &= F^{2L}(\mu_1, \dots, \mu_L, \\ &\quad -\mu_1 - n'_1 + 2n + \delta, \dots, -\mu_L - n'_L + 2n + \delta). \end{aligned}$$

If $\mathfrak{g} = \mathfrak{o}_{2n+1}$, we put

$$\begin{aligned} H(\Theta, \mu_1, \dots, \mu_L) &= F^{2L+1}(\mu_1, \dots, \mu_L, n, \\ &\quad -\mu_1 - n'_1 + 2n, \dots, -\mu_L - n'_{L-1} + 2n). \end{aligned}$$

Moreover we define

$$\begin{aligned} \tilde{H}(\Theta, \mu_1, \dots, \mu_{L-1}) &= F^{2L-1}(\mu_1, \dots, \mu_{L-1}, n_{L-1}, \\ &\quad -\mu_1 - n'_1 + 2n + \delta, \dots, -\mu_{L-1} - n'_{L-1} + 2n + \delta). \end{aligned}$$

Here

$$(3.14) \quad \delta = \begin{cases} 1 & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ 0 & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1} \text{ or } \mathfrak{gl}_n, \\ -1 & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}. \end{cases}$$

i) The off-diagonal elements of $F^k(\mu_1, \dots, \mu_k)$ satisfy

$$F_{ij}^k(\mu_1, \dots, \mu_k) \equiv 0 \pmod{U(\mathfrak{g}\mathfrak{n})} \quad \text{if } i > j,$$

$$F_{ij}^k(\mu_1, \dots, \mu_k) \equiv 0 \pmod{\bar{n}U(\mathfrak{g})} \quad \text{if } i < j.$$

ii) If $i \leq n$, then

$$F_{ii}^k(\mu_1, \dots, \mu_k) \equiv \begin{cases} 0 & \pmod{\tilde{J}(\mu)_i} \quad \text{if } i \leq n_k, \\ \prod_{\nu=1}^k (\mu_{k+1} - \mu_\nu - n'_\nu) & \pmod{\tilde{J}(\mu)_i} \quad \text{if } n_k < i \leq n_{k+1}. \end{cases}$$

iii) If $i < n$, then

$$F_{ii+1}^k(\mu_1, \dots, \mu_k) \equiv \left(\prod_{\nu=1}^{\ell-1} (\mu_\ell - \mu_\nu - n'_\nu - n_{\ell-1} + i) \prod_{\nu=\ell+1}^k (\mu_\ell - \mu_\nu - n_{\ell-1} + i) \right) F_{ii+1} \pmod{\tilde{J}(\mu)_i} \quad \text{if } n_{\ell-1} < i < n_\ell \text{ and } k \geq \ell.$$

iv) Suppose $\mathfrak{g} = \mathfrak{o}_{2n}$ or \mathfrak{o}_{2n+1} or \mathfrak{sp}_n . Then

$$H_{ii}(\Theta, \mu_1, \dots, \mu_L) \equiv 0 \pmod{\tilde{J}(\mu)_n} \quad \text{for } i = 1, \dots, N.$$

In particular, if $\mu_L = n_{L-1}$, then

$$\tilde{H}_{ii}(\Theta, \mu_1, \dots, \mu_{L-1}) \equiv 0 \pmod{\tilde{J}(\mu)_n} \quad \text{for } i = 1, \dots, N$$

and

$$\begin{aligned} \tilde{H}_{nn+1}(\Theta, \mu_1, \dots, \mu_{L-1}) \\ \equiv (-1)^{L-1} \left(\prod_{\nu=1}^{L-1} (\mu_\nu + n'_\nu - n)(\mu_\nu + n'_\nu - n - \delta) \right) F_{nn+1} \\ \pmod{U(\mathfrak{g})\tilde{J}(\mu)_n}. \end{aligned}$$

Proof. Put $F_{ij}^k(\mu) = F_{ij}^k(\mu_1, \dots, \mu_k)$ for simplicity. If $i < n$, it follows from (3.12) that

$$F_{i+1i+1}^k(\mu) - F_{ii}^k(\mu) \equiv F_{i+1i+1}^{k-1}(\mu)(F_{i+1} - \mu_k + i) - F_{ii}^{k-1}(\mu)(F_i - \mu_k + i) \pmod{U(\mathfrak{g})\mathfrak{n}}$$

and therefore by the induction on k we have

$$(3.15) \quad F_{ii}^k(\mu) \equiv F_{i+1i+1}^k(\mu) \pmod{U(\mathfrak{g})\mathfrak{n} + U(\mathfrak{g})(F_{i+1} - F_i)}.$$

Here we note that $F_{\nu+1} - F_\nu \in \tilde{J}(\mu)_{n_\ell}$ if $n_{\ell-1} < \nu < n_\ell$. Hence we have

$$F_{ii}^k(\mu) + \tilde{J}(\mu)_{n_\ell} = F_{n_\ell n_\ell}^k(\mu) + \tilde{J}(\mu)_{n_\ell} \quad \text{for } n_{\ell-1} < i \leq n_\ell \text{ and } 1 \leq \ell \leq L.$$

Put $s_\nu = n_\nu - n_{\nu-1}$ and introduce polynomials $f(k, \ell)$ of $(\mu_1, \dots, \mu_L, s_1, \dots, s_L)$ with $\ell \leq L$ so that

$$(3.16) \quad F_{n_\ell n_\ell}^k(\mu_1, \dots, \mu_k) \equiv f(k, \ell) \pmod{\tilde{J}(\mu)_{n_\ell}}.$$

Similarly for i with $n_{\ell-1} < i < n_\ell$, we put $t = \mu_\ell - n_{\ell-1} + i$ and define polynomials $g(k, \ell)$ of $(t, \mu_1, \dots, \mu_L, s_1, \dots, s_L)$ so that

$$(3.17) \quad F_{ii+1}^k(\mu_1, \dots, \mu_k) \equiv g(k, \ell) E_{ii+1} \pmod{\tilde{J}(\mu)_i}.$$

Then we have

$$(3.18) \quad \begin{aligned} f(k, \ell) &= \begin{cases} 1 & \text{if } k = 0, \\ f(k-1, \ell)(\mu_\ell - \mu_k) - \sum_{\nu=1}^{\ell-1} s_\nu f(k-1, \nu) & \text{if } k \geq 1, \end{cases} \\ g(k, \ell) &= \begin{cases} 1 & \text{if } k = 1, \\ g(k-1, \ell)(t - \mu_k) + f(k-1, \ell) & \text{if } k > 1. \end{cases} \end{aligned}$$

We will first prove $f(k, \ell) = 0$ if $k \geq \ell$ by the induction on ℓ . Putting $\ell = 1$ in (3.18), we have $f(k, 1) = f(k-1, 1)(\mu_1 - \mu_k)$ and $f(1, 1) = 0$ and therefore $f(k, 1) = 0$ for $k \geq 1$. Then if $k \geq \ell + 1$, we have $f(k, \ell + 1) = f(k-1, \ell + 1)(\mu_{\ell+1} - \mu_k) - \sum_{\nu=1}^{\ell} s_{\nu} f(k-1, \nu) = f(k-1, \ell + 1)(\mu_{\ell+1} - \mu_k)$ by the hypothesis of the induction. Hence we have $f(k, \ell + 1) = 0$ for $k \geq \ell + 1$ by the induction on k .

Putting $\mu_{\ell} = \mu_{\ell-1} + s_{\ell-1}$ in (3.18), we have $f(k, \ell) - f(k, \ell-1) = f(k-1, \ell) - f(k-1, \ell-1) = \dots = 0$ and therefore $f(\ell-1, \ell)|_{\mu_{\ell}=\mu_{\ell-1}+s_{\ell-1}} = 0$. Hence there exist polynomials $h(\ell)$ with $f(\ell-1, \ell) = h(\ell)(\mu_{\ell} - \mu_{\ell-1} - s_{\ell-1})$. Then (3.18) shows

$$h(\ell)(\mu_{\ell} - \mu_{\ell-1} - s_{\ell-1}) = f(\ell-2, \ell)(\mu_{\ell} - \mu_{\ell-1}) - s_{\ell-1}f(\ell-2, \ell-1).$$

It follows from (3.18) that $f(k, \ell)$ is a polynomial of degree at most 1 with respect to $s_{\ell-1}$ because $f(k, \nu)$ does not contain $s_{\ell-1}$ for $\nu < \ell$. Hence $h(\ell) = f(\ell-2, \ell)|_{s_{\ell-1}=0}$. Moreover by putting $s_{\ell-1} = 0$ in (3.18), it is clear that $f(\ell-2, \ell)|_{s_{\ell-1}=0}$ does not contain $\mu_{\ell-1}$. Hence $h(\ell) = f(\ell-2, \ell-1)|_{\mu_{\ell-1} \mapsto \mu_{\ell}}$ and we get

$$(3.19) \quad f(\ell-1, \ell) = \prod_{\nu=1}^{\ell-1} (\mu_{\ell} - \mu_{\nu} - s_{\nu})$$

by the induction on ℓ . Thus we have ii).

Now we put

$$(3.20) \quad f(\ell-1, \ell) = \sum_{\nu=0}^{\ell-1} c(\nu, \ell)(\mu_{\ell} - \mu_{\nu+1})(\mu_{\ell} - \mu_{\nu+2}) \cdots (\mu_{\ell} - \mu_{\ell-1})$$

with homogeneous polynomials $c(\nu, \ell)$ of $(\mu_1, \dots, \mu_{\ell-1}, s_1, \dots, s_{\ell-1})$ with degree ν . Here $c(\nu, \ell)$ does not contain μ_{ℓ} . Then by the induction on $k = \ell-1, \ell-2, \dots, 0$, (3.18) shows

$$(3.21) \quad \begin{aligned} f(k, \ell) &= \sum_{\nu=0}^k c(\nu, \ell)(\mu_{\ell} - \mu_{\nu+1})(\mu_{\ell} - \mu_{\nu+2}) \cdots (\mu_{\ell} - \mu_k), \\ - \sum_{\nu=1}^{\ell-1} s_{\nu} f(k-1, \nu) &= c(k, \ell) \end{aligned}$$

because $\sum_{\nu=1}^{\ell-1} s_{\nu} f(k-1, \nu)$ does not contain μ_{ℓ} . We will show

$$(3.22) \quad g(\ell, \ell) = \sum_{k=0}^{\ell-1} (t - \mu_{\ell})(t - \mu_{\ell-1}) \cdots (t - \mu_{k+2}) f(k, \ell)$$

$$(3.23) \quad = \prod_{\nu=1}^{\ell-1} (t - \mu_{\nu} - s_{\nu}).$$

Note that (3.22) is a direct consequence of (3.18). Denoting

$$g_k(\ell) = \sum_{\nu=0}^{\ell-1} c(\nu, \ell)(\mu_{\ell} - \mu_{\nu+1}) \cdots (\mu_{\ell} - \mu_{k-1})(\mu_{\ell} - \mu_k)(t - \mu_{k+1}) \cdots (t - \mu_{\ell-1})$$

for $k = 0, \dots, \ell-1$, we have

$$\begin{aligned} g_{k-1}(\ell) - g_k(\ell) &= \sum_{\nu=0}^{k-1} c(\nu, \ell)(\mu_{\ell} - \mu_{\nu+1}) \cdots (\mu_{\ell} - \mu_{k-1})(t - \mu_{\ell})(t - \mu_{k+1}) \cdots (t - \mu_{\ell-1}) \\ &= (t - \mu_{\ell})(t - \mu_{\ell-1}) \cdots (t - \mu_{k+1}) f(k-1, \ell) \end{aligned}$$

from (3.21) and therefore (3.22) shows

$$g(\ell, \ell) = g_{\ell-1}(\ell) + \sum_{k=1}^{\ell-1} (g_{k-1}(\ell) - g_k(\ell)) = g_0(\ell) = f(\ell-1, \ell)|_{\mu_{\ell} \mapsto t},$$

which implies (3.23). Since $f(k, \ell) = 0$ for $k \geq \ell$, (3.18) shows

$$(3.24) \quad g(k, \ell) = \prod_{\nu=1}^{\ell-1} (t - \mu_{\nu} - s_{\nu}) \prod_{\nu=\ell+1}^k (t - \mu_{\nu}) \quad \text{if } k \geq \ell,$$

from which iii) follows.

In general we have proved the following lemma.

Lemma 3.5. *The functions $f(k, \ell)$ and $g(k, \ell)$ of $\mu_1, \mu_2, \dots, s_1, s_2, \dots$ and t which are recursively defined by (3.18) satisfy (3.19), (3.24) and $f(k, \ell) = 0$ for $k \geq \ell \geq 1$.*

Now suppose $\mathfrak{g} = \mathfrak{sp}_n$ or \mathfrak{o}_{2n} . Then

$$(3.25) \quad F_{n+1n+1}^k(\mu) \equiv F_{n+1n+1}^{k-1}(\mu)(F_{n+1} - \mu_k) + \sum_{\nu=1}^n (F_{n+1n+1}^{k-1}(\mu) - F_{\nu\nu}^{n-1}(\mu)) \\ + \delta(F_{n+1n+1}^{k-1}(\mu) - F_{nn}^{k-1}(\mu)) \equiv 0 \quad \text{mod } U(\mathfrak{g}\mathfrak{n}).$$

Hence

$$(3.26) \quad F_{n+1n+1}^k(\mu) - F_{nn}^k(\mu) \\ \equiv F_{n+1n+1}^{k-1}(\mu)(F_{n+1} - \mu_k + n + \delta) - F_{nn}^{k-1}(\mu)(F_n - \mu_k + n + \delta) \\ \equiv 0 \quad \text{mod } U(\mathfrak{g}\mathfrak{n}) + U(\mathfrak{g})(F_{n+1} - F_n)$$

by the induction on k and

$$(3.27) \quad F_{n+1n+1}^k(\mu) \equiv 0 \quad \text{mod } \sum_{\nu=1}^n U(\mathfrak{g})F_{\nu\nu}^{k-1}(\mu) + U(\mathfrak{g}\mathfrak{n}) + U(\mathfrak{g})(F_{n+1} - \mu_k + n + \delta).$$

Since $F_{n+1} = -F_n$, we have from (3.25)

$$F_{n+1n+1}^L(\mu_1, \dots, \mu_{L-1}, n_{L-1}) \equiv 0 \quad \text{mod } \tilde{J}(\mu)_{n_{L-1}} + \sum_{\nu=n_{L-1}+1}^n U(\mathfrak{g})F_{\nu}$$

in the case $\mu_L = n_{L-1}$ and from (3.27) with $-(\mu_L - n_{L-1}) - \mu_{L+1} + n + \delta = 0$

$$F_{n+1n+1}^{L+1}(\mu_1, \dots, \mu_L, -\mu_L + n_{L-1} + n + \delta) \equiv 0 \quad \text{mod } \tilde{J}(\mu)_n.$$

Suppose $i < n$. Then

$$F_{ii}^k(\mu) \equiv F_{ii}^{k-1}(\mu)(F_{\bar{i}} - \mu_k) + \sum_{\nu=1}^{\bar{i}-1} (F_{ii}^{k-1}(\mu) - F_{\nu\nu}^{k-1}(\mu)) + \delta(F_{ii}^{k-1}(\mu) - F_{ii}^{k-1}(\mu)) \\ \text{mod } U(\mathfrak{g}\mathfrak{n})$$

and therefore

$$(3.28) \quad F_{i+1\bar{i}+1}^k(\mu) - F_{ii}^k(\mu) \equiv F_{i+1\bar{i}+1}^{k-1}(\mu)(F_{\bar{i}+1} - \mu_k + \bar{i} + \delta) - F_{ii}^{k-1}(\mu)(F_{\bar{i}} - \mu_k + \bar{i} + \delta) \\ + \delta(F_{i-1\bar{i}-1}^{k-1}(\mu) - F_{ii}^{k-1}(\mu)) \quad \text{mod } U(\mathfrak{g}) \\ \equiv 0 \quad \text{mod } U(\mathfrak{g}\mathfrak{n}) + U(\mathfrak{g})(F_{\bar{i}} - F_{i-1}) + U(\mathfrak{g})(F_{\bar{i}-1} - F_{\bar{i}}), \\ F_{\bar{n}_p \bar{n}_p}^k(\mu) \equiv 0 \quad \text{mod } \sum_{\nu=1}^{\bar{n}_p-1} U(\mathfrak{g})F_{\nu\nu}^{k-1}(\mu) + U(\mathfrak{g}\mathfrak{n}) + U(\mathfrak{g})(F_{\bar{n}_p} - \mu_k + \bar{n}_p - 1 + \delta).$$

Note that $F_{\bar{n}_p} - \mu_k + \bar{n}_p - 1 + \delta = (F_{\bar{n}_p} + \mu_p - n_{p-1}) - \mu_k - \mu_p + n_{p-1} - n_p + 2n + \delta$. Since $F_{\bar{\nu}} = -F_{\nu}$, we have

$$(3.29) \quad F_{\bar{i}+1\bar{i}+1}^k(\mu) \equiv F_{\bar{i}\bar{i}}^k(\mu) \pmod{\tilde{J}(\mu)_n} \quad \text{for } \bar{n}_p \leq \bar{i} < \bar{n}_{p-1}$$

and hence by the induction on $p = L, L-1, \dots, 1$, we have

$$(3.30) \quad F_{\bar{n}_p\bar{n}_p}^{2L+1-p}(\mu_1, \dots, \mu_L, -\mu_L + n_{L-1} - n_L + 2n + \delta, \\ \dots, -\mu_p + n_{p-1} - n_p + 2n + \delta) \equiv 0 \pmod{\tilde{J}(\mu)_n}$$

and if $\mu_L = n_{L-1}$, then

$$(3.31) \quad F_{\bar{n}_p\bar{n}_p}^{2L-p}(\mu_1, \dots, \mu_{L-1}, n_{L-1}, \mu_{L-1} + n_{L-2} - n_{L-1} + 2n + \delta, \\ \dots, -\mu_p + n_{p-1} - n_p + 2n + \delta) \equiv 0 \pmod{\tilde{J}(\mu)_n}.$$

Suppose $\mu_L = n_{L-1}$ and $\mathfrak{g} = \mathfrak{sp}_n$. Then from (3.13) we have

$$(3.32) \quad F_{nn+1}^k(\mu) \equiv F_{nn+1}^{k-1}(\mu)(F_{n+1n+1} - \mu_k + n + \delta) + F_{nn+1}F_{nn}^{k-1}(\mu) \pmod{U(\mathfrak{g})\mathfrak{n}} \\ \equiv F_{nn+1}^L(\mu) \prod_{\nu=L+1}^k (-\mu_\nu + n + \delta) \pmod{\tilde{J}(\mu)_n} \quad \text{if } k \geq L.$$

It follows from Lemma 3.5 with $t = n+1$ that

$$\bar{H}_{nn+1}(\Theta, \mu) \equiv F_{nn+1} \prod_{\nu=1}^{L-1} (-\mu_\nu + n_{\nu-1} - n_\nu + n + \delta) \prod_{\nu=1}^{L-1} (\mu_\nu - n_{\nu-1} + n_\nu - n) \\ \pmod{\tilde{J}(\mu)_n}.$$

Thus we have proved iv).

Lastly suppose $\mathfrak{g} = \mathfrak{o}_{2n+1}$. Note that $F_{n+1} = 0$ and $F_{n+2} = -F_n$. Then

$$F_{n+1n+1}^k(\mu) \equiv F_{n+1n+1}^{k-1}(F_{n+1} - \mu_k) + \sum_{\nu=1}^n (F_{n+1n+1}^{k-1}(\mu) - F_{\nu\nu}^{k-1}(\mu)) \pmod{U(\mathfrak{g})\mathfrak{n}} \\ \equiv 0 \pmod{\sum_{\nu=1}^n U(\mathfrak{g})F_{\nu\nu}^{k-1}(\mu) + U(\mathfrak{g})(-\mu_k + n)},$$

$$F_{n+2n+2}^k(\mu) \equiv F_{n+2n+2}^{k-1}(F_{n+2} - \mu_k) + \sum_{\nu=1}^{n+1} (F_{n+2n+2}^{k-1}(\mu) - F_{\nu\nu}^{k-1}(\mu)) \\ - (F_{n+2n+2}^{k-1}(\mu) - F_{nn}^{k-1}(\mu)) \pmod{U(\mathfrak{g})\mathfrak{n}} \\ \equiv 0 \pmod{\sum_{\nu=1}^{n+1} U(\mathfrak{g})F_{\nu\nu}^{k-1}(\mu) + U(\mathfrak{g})(-F_n - \mu_k + n)}$$

and

$$F_{n+1n+1}^{L+1}(\mu_1, \dots, \mu_L, n) \equiv 0 \pmod{\tilde{J}(\mu)_n}, \\ F_{n+2n+2}^{L+2}(\mu_1, \dots, \mu_L, n, -\mu_L + n_{L-1} + n) \equiv 0 \pmod{\tilde{J}(\mu)_n},$$

Since

$$F_{n+1n+1}^k(\mu) - F_{nn}^k(\mu) \equiv F_{n+1n+1}^{k-1}(F_{n+1} - \mu_k + n) - F_{nn}^{k-1}(F_n - \mu_k + n) \\ \pmod{U(\mathfrak{g})\mathfrak{n}} \\ F_{n+2n+2}^k(\mu) - F_{n+1n+1}^k(\mu) \equiv F_{n+2n+2}^{k-1}(F_{n+2} - \mu_k + n) - F_{n+1n+1}^{k-1}(F_{n+1} - \mu_k + n) \\ - (F_{n+1n+1}^{k-1}(\mu) - F_{nn}^{k-1}(\mu)) \pmod{U(\mathfrak{g})\mathfrak{n}},$$

we have

$$F_{n+2n+2}^k(\mu) \equiv F_{n+1n+1}^k(\mu) \equiv F_{nn}^k(\mu) \pmod{U(\mathfrak{g})\mathfrak{n} + U(\mathfrak{g})F_n}$$

and

$$F_{n+1n+1}^L(\mu) \equiv F_{n+2n+2}^L(\mu) \equiv 0 \pmod{\tilde{J}(\mu)_n} \quad \text{if } \mu_L = n_{L-1}.$$

Note that (3.28) is valid if $\bar{i} < n$. But since $F_{\bar{n}_p} - \mu_k + \bar{n}_p - 1 + \delta = -\mu_k - (F_{n_p} - \mu_p + n_{p-1}) - \mu_p + n_{p-1} - n_p + 2n$, we have

$$F_{\bar{n}_p \bar{n}_p}^{2L+2-p}(\mu_1, \dots, \mu_L, n, -\mu_L + n_{L-1} - n_L + 2n, \dots, -\mu_p + n_{p-1} - n_p + 2n) \equiv 0 \pmod{\tilde{J}(\mu)_n}$$

for $p = L, L-1, \dots, 1$. Similarly we have (3.31) with $\delta = 0$ if $\mu_L = n_{L-1}$. Moreover (3.32) is valid with $\delta = 0$ and we have iv) as in the case of $\mathfrak{g} = \mathfrak{sp}_n$. \square

4. GENERALIZED VERMA MODULES

In this section we define by (4.14) a two-sided ideal of $U^\epsilon(\mathfrak{g})$ associated to every generalized Verma module of the scalar type for the classical Lie algebra \mathfrak{g} . We have Theorem 4.4 which shows that the ideal describes the gap between the generalized Verma module and the usual Verma module and then Corollary 4.6 says that the ideal equals the annihilator of the generalized Verma module. In the classical limit, an explicit generator system of every semisimple co-adjoint orbit of \mathfrak{g} is given in Theorem 4.11.

The ideal is constructed from a polynomial of a matrix with elements in \mathfrak{g} and the polynomial is proved to be a minimal polynomial in a certain sense (cf. Proposition 4.16 and 4.18). Lastly in this section Theorem 4.19 gives a description of the image of the Harish-Chandra homomorphism of any polynomial function of the matrix $(E_{ij}) \in M(n, \mathfrak{g})$.

Retain the notation in the previous section. Let $\Theta = \{(0 <)n_1 < n_2 < \dots < n_L (= n)\}$ be the sequence of strictly increasing positive integers ending at n . Put

$$H_\Theta = \sum_{k=1}^L \sum_{i=1}^{n_k} F_i \quad \text{and} \quad H_{\bar{\Theta}} = \sum_{k=1}^{L-1} \sum_{i=1}^{n_k} F_i.$$

Recall that $F_i = F_{ii}$, $F = (F_{ij}) \in M(N, \mathfrak{g})$, $\mathfrak{n} = \sum_{i>j} \mathbb{C}F_{ij}$, $\mathfrak{a} = \sum_i \mathbb{C}F_i$, $\bar{\mathfrak{n}} = \sum_{i<j} \mathbb{C}F_{ij}$ and $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{a} \oplus \bar{\mathfrak{n}}$. Note that $F_{ij} = E_{ij}$ in the case $\mathfrak{g} = \mathfrak{gl}_n$ and $F_{ij} = E_{ij} + \sigma_{\mathfrak{g}}(E_{ij})$ in the case $\mathfrak{g} = \mathfrak{o}_{2n+1}$, \mathfrak{sp}_n or \mathfrak{o}_{2n} . Here $\sigma_{\mathfrak{g}}$ is the involution of \mathfrak{gl}_N to define \mathfrak{g} in (3.1) so that \mathfrak{g} is the subalgebra of \mathfrak{gl}_N fixed by $\sigma_{\mathfrak{g}}$. Let G be the analytic subgroup of $GL(N, \mathbb{C})$ with the Lie algebra \mathfrak{g} . Namely $G = GL(n, \mathbb{C})$, $O(2n+1, \mathbb{C})$, $Sp(n, \mathbb{C})$ or $O(2n, \mathbb{C})$.

Define

$$(4.1) \quad \begin{cases} \mathfrak{m}_\Theta = \{X \in \mathfrak{g}; \text{ad}(H_\Theta)X = 0\}, \\ \mathfrak{n}_\Theta = \{X \in \mathfrak{n}; \langle X, \mathfrak{m}_\Theta \rangle = 0\}, \quad \bar{\mathfrak{n}}_\Theta = \{X \in \bar{\mathfrak{n}}; \langle X, \mathfrak{m}_\Theta \rangle = 0\}, \\ \mathfrak{p}_\Theta = \mathfrak{m}_\Theta + \mathfrak{n}_\Theta. \end{cases}$$

We similarly define $m_{\bar{\Theta}}$, $\mathfrak{n}_{\bar{\Theta}}$, $\bar{\mathfrak{n}}_{\bar{\Theta}}$ and $\mathfrak{p}_{\bar{\Theta}}$ replacing Θ by $\bar{\Theta}$ in the above definition. Then $\mathfrak{n} = \mathfrak{n}_{\{1,2,\dots,n\}}$, $\bar{\mathfrak{n}} = \bar{\mathfrak{n}}_{\{1,2,\dots,n\}}$, $\mathfrak{a} = \mathfrak{a}_{\{1,2,\dots,n\}}$ and \mathfrak{p}_Θ and $\mathfrak{p}_{\bar{\Theta}}$ are parabolic subalgebras of \mathfrak{g} containing the Borel subalgebra $\mathfrak{b} = \mathfrak{n} + \mathfrak{a}$.

Let $\{e_1, \dots, e_n\}$ be the dual bases of $\{F_1, \dots, F_n\}$. Then the fundamental system $\Psi(\mathfrak{a})$ for the pair $(\mathfrak{n}, \mathfrak{a})$ is

$$(4.2) \quad \Psi(\mathfrak{a}) = \begin{cases} \{e_2 - e_1, e_3 - e_2, \dots, e_n - e_{n-1}\} & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ \{e_2 - e_1, e_3 - e_2, \dots, e_n - e_{n-1}, -e_n\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}, \\ \{e_2 - e_1, e_3 - e_2, \dots, e_n - e_{n-1}, -2e_n\} & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ \{e_2 - e_1, e_3 - e_2, \dots, e_n - e_{n-1}, -e_n - e_{n-1}\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}. \end{cases}$$

We put $\alpha_j = e_{j+1} - e_j$ for $j = 1, \dots, n-1$ and $\alpha_n = -e_n$ or $-2e_n$ or $-e_n - e_{n-1}$ if $\mathfrak{g} = \mathfrak{o}_{2n+1}$ or \mathfrak{sp}_n or \mathfrak{o}_{2n} , respectively. Then the fundamental system for $(\mathfrak{m}_\Theta \cap \mathfrak{n}, \mathfrak{a})$ is $\Psi(\mathfrak{a}) \setminus \{\alpha_{n_1}, \dots, \alpha_{n_{L-1}}\}$ and that for $(\mathfrak{m}_{\bar{\Theta}} \cap \mathfrak{n}, \mathfrak{a})$ is

$$\begin{cases} \Psi(\mathfrak{a}) \setminus \{\alpha_{n_1}, \dots, \alpha_{n_{L-1}}, \alpha_n\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1} \text{ or } \mathfrak{sp}_n, \\ \Psi(\mathfrak{a}) \setminus \{\alpha_{n_1}, \dots, \alpha_{n_{L-1}}\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n} \text{ and } n_{L-1} \neq n-1, \\ \Psi(\mathfrak{a}) \setminus \{\alpha_{n_1}, \dots, \alpha_{n_{L-1}}, \alpha_n\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n} \text{ and } n_{L-1} = n-1. \end{cases}$$

Then the Dynkin diagram of \mathfrak{g} is as follows:

$$(4.3) \quad \begin{array}{ccc} \begin{array}{c} \alpha_1 \quad \alpha_2 \quad \dots \quad \alpha_{n-2} \quad \alpha_{n-1} \\ \circ \text{---} \circ \text{---} \dots \text{---} \circ \text{---} \circ \\ \mathfrak{gl}_n \end{array} & \begin{array}{c} \alpha_1 \quad \alpha_2 \quad \dots \quad \alpha_{n-1} \quad \alpha_n \\ \circ \text{---} \circ \text{---} \dots \text{---} \circ \text{---} \circ \Rightarrow \\ \mathfrak{o}_{2n+1} \end{array} \\ \\ \begin{array}{c} \alpha_1 \quad \alpha_2 \quad \dots \quad \alpha_{n-1} \quad \alpha_n \\ \circ \text{---} \circ \text{---} \dots \text{---} \circ \Leftarrow \circ \\ \mathfrak{sp}_n \end{array} & \begin{array}{c} \alpha_1 \quad \alpha_2 \quad \dots \quad \alpha_{n-2} \quad \alpha_{n-1} \\ \circ \text{---} \circ \text{---} \dots \text{---} \circ \text{---} \circ \\ \alpha_n \\ \circ \\ \mathfrak{o}_{2n} \end{array} \end{array}$$

Fix $\lambda = (\lambda_1, \dots, \lambda_L) \in \mathbb{C}^L$ and define a character λ_Θ of \mathfrak{p}_Θ

$$(4.4) \quad \lambda_\Theta(X + \sum_{i=1}^n C_i F_i) = \sum_{i=1}^n C_i \lambda_{\iota_\Theta(i)} \quad \text{for } X \in \mathfrak{n}_\Theta + [\mathfrak{m}_\Theta, \mathfrak{m}_\Theta].$$

We similarly define a character $\lambda_{\bar{\Theta}}$ of $\mathfrak{p}_{\bar{\Theta}}$ if $\lambda_L = 0$.

We introduce the *homogenized universal enveloping algebra*

$$(4.5) \quad U^\epsilon(\mathfrak{g}) = \left(\sum_{k=0}^{\infty} \otimes^k \mathfrak{g} \right) / \langle X \otimes Y - Y \otimes X - \epsilon[X, Y]; X, Y \in \mathfrak{g} \rangle.$$

of \mathfrak{g} as in [O4]. Here ϵ is a central element of $U^\epsilon(\mathfrak{g})$. Let $U^\epsilon(\mathfrak{g})^{(m)}$ be the image of $\sum_{k=0}^m \otimes^k \mathfrak{g}$ in $U^\epsilon(\mathfrak{g})$ and let $Z^\epsilon(\mathfrak{g})$ be the subalgebra of G -invariants of $U^\epsilon(\mathfrak{g})$. Fix generators $\Delta_1, \dots, \Delta_n$ of $Z^\epsilon(\mathfrak{g})$ so that

$$(4.6) \quad \begin{cases} \Delta_j \in U^\epsilon(\mathfrak{g})^{(j)} & (1 \leq j \leq n) & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ \Delta_j \in U^\epsilon(\mathfrak{g})^{(2j)} & (1 \leq j \leq n) & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1} \text{ or } \mathfrak{sp}_n, \\ \Delta_j \in U^\epsilon(\mathfrak{g})^{(2j)} & (1 \leq j < n), \quad \Delta_n \in U^\epsilon(\mathfrak{g})^{(n)} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}. \end{cases}$$

If $\mathfrak{g} = \mathfrak{o}_{2n}$, we assume that Δ_n changes into $-\Delta_n$ by the outer automorphism of \mathfrak{o}_{2n} which maps $(F_1, \dots, F_{n-1}, F_n)$ to $(F_1, \dots, F_{n-1}, -F_n)$. Moreover put

$$(4.7) \quad \begin{cases} J_{\bar{\Theta}}^\epsilon(\lambda) = \sum_{X \in \mathfrak{p}_{\bar{\Theta}}} U^\epsilon(\mathfrak{g})(X - \lambda_{\bar{\Theta}}(X)), & M_{\bar{\Theta}}^\epsilon(\lambda) = U^\epsilon(\mathfrak{g})/J_{\bar{\Theta}}^\epsilon(\lambda), \\ J_{\Theta}^\epsilon(\lambda) = \sum_{X \in \mathfrak{p}_{\Theta}} U^\epsilon(\mathfrak{g})(X - \lambda_{\Theta}(X)), & M_{\Theta}^\epsilon(\lambda) = U^\epsilon(\mathfrak{g})/J_{\Theta}^\epsilon(\lambda), \\ J^\epsilon(\lambda_\Theta) = \sum_{X \in \mathfrak{b}} U^\epsilon(\mathfrak{g})(X - \lambda_\Theta(X)), & M^\epsilon(\lambda_\Theta) = U^\epsilon(\mathfrak{g})/J^\epsilon(\lambda_\Theta). \end{cases}$$

For a $U^\epsilon(\mathfrak{g})$ -module M the annihilator of M is denoted by $\text{Ann}(M)$ and put $\text{Ann}_G(M) = \bigcap_{g \in G} \text{Ad}(g) \text{Ann}(M)$. Note that $\text{Ann}_G(M) = \text{Ann}(M)$ if $\epsilon \neq 0$.

When $\epsilon = 1$, $U^\epsilon(\mathfrak{g})$ is the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} and we will sometimes omit the superfix ϵ for $J_{\bar{\Theta}}^\epsilon(\lambda)$ and $M_{\bar{\Theta}}^\epsilon(\lambda)$ etc. Then $M_{\Theta}(\lambda)$ and $M_{\bar{\Theta}}(\lambda)$ are generalized Verma modules which are quotients of the Verma module $M(\lambda_{\Theta})$.

Remark 4.1. i) Suppose $\mathfrak{g} = \mathfrak{o}_{2n}$. Then we have not considered the parabolic subalgebra \mathfrak{p} such that the fundamental system for $(\mathfrak{m}_{\mathfrak{p}}, \mathfrak{a})$ contains α_{n-1} and does not contain α_n . But this is reduced to the case when it contains α_n and does not contain α_{n-1} by the outer automorphism of \mathfrak{o}_{2n} .

ii) Considering the above remark, the parabolic subalgebra \mathfrak{p} containing the Borel subalgebra \mathfrak{b} corresponds to \mathfrak{p}_{Θ} or $\mathfrak{p}_{\bar{\Theta}}$ and therefore we will sometimes use the notation $\mathfrak{m}_{\mathfrak{p}}, \mathfrak{n}_{\mathfrak{p}}, \bar{\mathfrak{n}}_{\mathfrak{p}}, \lambda_{\mathfrak{p}}, J_{\mathfrak{p}}^\epsilon(\lambda), M_{\mathfrak{p}}^\epsilon(\lambda)$ and $M^\epsilon(\lambda_{\mathfrak{p}})$ for $\mathfrak{m}_{\Theta'}, \mathfrak{n}_{\Theta'}, \bar{\mathfrak{n}}_{\Theta'}, \lambda_{\Theta}, L_{\Theta'}^\epsilon(\lambda), M_{\bar{\Theta}'}^\epsilon(\lambda)$ and $M^\epsilon(\lambda_{\Theta})$, respectively, by this correspondence. Note that $\Theta' = \bar{\Theta}$ means $\lambda_n = 0$.

Let $\rho \in \mathfrak{a}^*$ with $\rho(H) = \frac{1}{2} \text{Trace}(\text{ad}(H))|_{\mathfrak{n}}$ for $H \in \mathfrak{a}$. Then with δ in (3.14)

$$(4.8) \quad \rho = \begin{cases} \sum_{\nu=1}^n (\nu - \frac{n+1}{2}) e_{\nu} & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ \sum_{\nu=1}^n (\nu - n - \frac{1}{2}) e_{\nu} = \sum_{\nu=1}^n (\nu - n - \frac{\delta+1}{2}) e_{\nu} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}, \\ \sum_{\nu=1}^n (\nu - n - 1) e_{\nu} = \sum_{\nu=1}^n (\nu - n - \frac{\delta+1}{2}) e_{\nu} & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ \sum_{\nu=1}^n (\nu - n) e_{\nu} = \sum_{\nu=1}^n (\nu - n - \frac{\delta+1}{2}) e_{\nu} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}. \end{cases}$$

We define $\bar{\lambda} = (\bar{\lambda}_1, \dots, \bar{\lambda}_n) \in \mathbb{C}^n$ by

$$(4.9) \quad \lambda_{\Theta}|_{\mathfrak{a}} + \epsilon\rho = \bar{\lambda}_1 e_1 + \bar{\lambda}_2 e_2 + \dots + \bar{\lambda}_n e_n.$$

For $P \in U^\epsilon(\mathfrak{g})$ let $\omega(P)$ and $\bar{\omega}(P)$ denotes the elements of $S(\mathfrak{a}) \simeq U^\epsilon(\mathfrak{a})$ with

$$(4.10) \quad \begin{aligned} P - \omega(P) &\in \bar{\mathfrak{n}}U^\epsilon(\mathfrak{g}) + U^\epsilon(\mathfrak{g})\mathfrak{n}, \\ \bar{\omega}(P)(\mu + \epsilon\rho) &= \omega(P)(\mu) \quad \text{for } \forall \mu \in \mathfrak{a}^*. \end{aligned}$$

Then $\bar{\omega}$ induces the Harish-Chandra isomorphism

$$(4.11) \quad \bar{\omega} : Z^\epsilon(\mathfrak{g}) \xrightarrow{\sim} S(\mathfrak{a})^W.$$

Here W is the Weyl group for the pair $(\mathfrak{g}, \mathfrak{a})$ and $S(\mathfrak{a})^W$ denotes the totality of W -invariants in $S(\mathfrak{a})$.

Definition 4.2. Retain the above notation and define polynomials

$$(4.12) \quad \begin{cases} q_{\bar{\Theta}}^\epsilon(\mathfrak{gl}_n; x, \lambda) = \prod_{j=1}^L (x - \lambda_j - n_{j-1}\epsilon), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{o}_{2n+1}; x, \lambda) = (x - n\epsilon) \prod_{j=1}^L (x - \lambda_j - n_{j-1}\epsilon)(x + \lambda_j + (n_j - 2n)\epsilon), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{sp}_n; x, \lambda) = \prod_{j=1}^L (x - \lambda_j - n_{j-1}\epsilon)(x + \lambda_j + (n_j - 2n - 1)\epsilon), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{o}_{2n}; x, \lambda) = \prod_{j=1}^L (x - \lambda_j - n_{j-1}\epsilon)(x + \lambda_j + (n_j - 2n + 1)\epsilon) \end{cases}$$

and if $\mathfrak{g} = \mathfrak{sp}_n$ or \mathfrak{o}_{2n+1} or \mathfrak{o}_{2n} ,

$$(4.13) \quad q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; x, \lambda) = (x - n_{L-1}\epsilon) \prod_{j=1}^{L-1} (x - \lambda_j - n_{j-1}\epsilon)(x + \lambda_j + (n_j - 2n - \delta)\epsilon)$$

with the δ given by (3.3). Furthermore define two-sided ideals of $U^\epsilon(\mathfrak{g})$

$$(4.14) \quad \begin{cases} I_{\bar{\Theta}}^\epsilon(\lambda) = \sum_{i=1}^N \sum_{j=1}^N U^\epsilon(\mathfrak{g}) q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; F, \lambda)_{ij} + \sum_{j \in \bar{J}} U^\epsilon(\mathfrak{g}) (\Delta_j - \omega(\Delta_j)(\lambda_{\bar{\Theta}})), \\ I_{\Theta}^\epsilon(\lambda) = \sum_{i=1}^N \sum_{j=1}^N U^\epsilon(\mathfrak{g}) q_{\Theta}^\epsilon(\mathfrak{g}; F, \lambda)_{ij} + \sum_{j \in \bar{J}} U^\epsilon(\mathfrak{g}) (\Delta_j - \omega(\Delta_j)(\lambda_{\Theta})) \end{cases}$$

with

$$(4.15) \quad \begin{cases} J = \{1, 2, \dots, L-1\} & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ J = \{1, 2, \dots, L\}, \bar{J} = \{1, 2, \dots, L-1\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}, \\ J = \bar{J} = \{1, 2, \dots, L-1\} & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ J = \bar{J} = \{1, 2, \dots, L-1\} \cup \{n\} & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}. \end{cases}$$

Remark 4.3. i) Let $p(x)$ and $q(x)$ be monic polynomials with $q(x) \in \mathbb{C}[x]p(x)$. Then

$$\begin{cases} \sum_{i=1}^N p(F)_{ii} \in Z^\epsilon(\mathfrak{g}), \\ \sum_{i=1}^N p(F)_{ii} - \sum_{i=1}^N F_i^{\deg p} \in \bar{\mathfrak{n}}U^\epsilon(\mathfrak{g}) + U^\epsilon(\mathfrak{g})\mathfrak{n} + U^\epsilon(\mathfrak{g})^{(\deg p-1)}, \\ q(F)_{ij} \in \sum_{\substack{1 \leq \mu \leq N \\ 1 \leq \nu \leq N}} U^\epsilon(\mathfrak{g}) p(F)_{\mu\nu}. \end{cases}$$

Hence it is clear

$$(4.16) \quad I_{\Theta'}^\epsilon(\lambda) \supset \sum_{D \in Z^\epsilon(\mathfrak{g})} U^\epsilon(\mathfrak{g})(D - \omega(D)(\lambda_{\Theta'})) \quad \text{for } \Theta' = \Theta \text{ and } \bar{\Theta}.$$

Note that it is known that the right hand side of the above equals $\text{Ann}_G(M^\epsilon(\lambda_{\Theta}))$.

ii) $I_{\bar{\Theta}}^\epsilon(\lambda)$ and $I_{\Theta}^\epsilon(\lambda)$ are homogeneous ideals with respect to $(\mathfrak{g}, \lambda, \epsilon)$.

Now we give the main theorem in this paper:

Theorem 4.4. i) Let $\mathfrak{g} = \mathfrak{gl}_n, \mathfrak{o}_{2n+1}, \mathfrak{sp}_n$ or \mathfrak{o}_{2n} . Then

$$(4.17) \quad \begin{cases} I_{\bar{\Theta}}^\epsilon(\lambda) \subset \text{Ann}(M_{\bar{\Theta}}^\epsilon(\lambda)), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; F, \lambda)_{ii+1} \equiv r_i^\epsilon(\mathfrak{g}; \Theta, \lambda) F_{ii+1} \pmod{J^\epsilon(\lambda_{\bar{\Theta}})} & \text{if } n_{k-1} < i < n_k, \\ J_{\bar{\Theta}}^\epsilon(\lambda) = I_{\bar{\Theta}}^\epsilon(\lambda) + J^\epsilon(\lambda_{\bar{\Theta}}) & \text{if } r^\epsilon(\mathfrak{g}; \Theta, \lambda) \neq 0. \end{cases}$$

Here

$$\begin{aligned} r_i^\epsilon(\mathfrak{gl}_n; \Theta, \lambda) &= \prod_{\nu=1}^{k-1} (\lambda_k - \lambda_\nu - (n_\nu - i)\epsilon) \prod_{\nu=k+1}^L (\lambda_k - \lambda_\nu - (n_{\nu-1} - i)\epsilon) \\ &= \prod_{\nu=1}^{k-1} (\bar{\lambda}_i - \bar{\lambda}_{n_\nu}) \prod_{\nu=k+1}^L (\bar{\lambda}_{i+1} - \bar{\lambda}_{n_{\nu-1}+1}), \end{aligned}$$

$$\begin{aligned} r_i^\epsilon(\mathfrak{sp}_n; \Theta, \lambda) &= r_i^\epsilon(\mathfrak{o}_{2n}; \Theta, \lambda) \\ &= r_i^\epsilon(\mathfrak{gl}_n; \Theta, \lambda) \prod_{\nu=1}^L (\lambda_k + \lambda_\nu + (n_\nu - 2n - \delta + i)\epsilon) \\ &= \prod_{\nu=1}^{k-1} (\bar{\lambda}_i - \bar{\lambda}_{n_\nu}) \prod_{\nu=k+1}^L (\bar{\lambda}_{i+1} - \bar{\lambda}_{n_{\nu-1}+1}) \prod_{\nu=1}^L (\bar{\lambda}_{i+1} + \bar{\lambda}_{n_\nu}), \end{aligned}$$

$$r_i^\epsilon(\mathfrak{o}_{2n+1}; \Theta, \lambda) = r_i^\epsilon(\mathfrak{gl}_n; \Theta, \lambda) (\lambda_k - (n - i)\epsilon) \prod_{\nu=1}^L (\lambda_k + \lambda_\nu + (n_\nu - 2n + i)\epsilon)$$

$$= \frac{1}{2}(\bar{\lambda}_i + \bar{\lambda}_{i+1}) \prod_{\nu=1}^{k-1} (\bar{\lambda}_i - \bar{\lambda}_{n_\nu}) \prod_{\nu=k+1}^L (\bar{\lambda}_{i+1} - \bar{\lambda}_{n_{\nu-1}+1}) \prod_{\nu=1}^L (\bar{\lambda}_{i+1} + \bar{\lambda}_{n_\nu})$$

if $n_{k-1} < i < n_k$ and then

$$(4.18) \quad r^\epsilon(\mathfrak{g}; \Theta, \lambda) = \prod_{k=1}^L \prod_{n_{k-1} < i < n_k} r_i^\epsilon(\mathfrak{g}; \Theta, \lambda).$$

ii) Suppose $\lambda_L = 0$. If $\mathfrak{g} = \mathfrak{sp}_n$ or \mathfrak{o}_{2n+1} or \mathfrak{o}_{2n} , then

$$(4.19) \quad \begin{cases} I_{\bar{\Theta}}^\epsilon(\lambda) \subset \text{Ann}(M_{\bar{\Theta}}^\epsilon(\lambda)), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; F, \lambda)_{ii+1} \equiv r_i^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) F_{ii+1} \pmod{J^\epsilon(\lambda_\Theta)} & \text{if } \iota_\Theta(i) = \iota_\Theta(i+1), \\ q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; F, \lambda)_{nn+1} \equiv \bar{r}^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) F_{nn+1} \pmod{J^\epsilon(\lambda_\Theta)} & \text{if } \mathfrak{g} \neq \mathfrak{o}_{2n}, \\ J_{\bar{\Theta}}^\epsilon(\lambda) = I_{\bar{\Theta}}^\epsilon(\lambda) + J^\epsilon(\lambda_\Theta) & \text{if } r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) \neq 0 \end{cases}$$

with denoting

$$\begin{aligned} r_i^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) &= \prod_{\nu=1}^{k-1} (\lambda_k - \lambda_\nu - (n_\nu - i)\epsilon) \prod_{\nu=k+1}^L (\lambda_k - \lambda_\nu - (n_{\nu-1} - i)\epsilon) \\ &\quad \cdot \prod_{\nu=1}^{L-1} (\lambda_k + \lambda_\nu + (n_\nu - 2n - \delta + i)\epsilon) \\ &= \prod_{\nu=1}^{k-1} (\bar{\lambda}_i - \bar{\lambda}_{n_\nu}) \prod_{\nu=k+1}^L (\bar{\lambda}_{i+1} - \bar{\lambda}_{n_{\nu-1}+1}) \prod_{\nu=1}^{L-1} (\bar{\lambda}_{i+1} + \bar{\lambda}_{n_\nu}) \\ &\quad \text{if } n_{k-1} < i < n_k, \\ \bar{r}_o^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) &= (-1)^{L-1} \prod_{\nu=1}^{L-1} (\lambda_\nu + (n_\nu - n)\epsilon)(\lambda_\nu + (n_\nu - n - \delta)\epsilon) \\ &= \begin{cases} (-1)^{L-1} \prod_{\nu=1}^{L-1} \bar{\lambda}_{n_\nu} (\bar{\lambda}_{n_\nu} - \bar{\lambda}_n) & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ (-1)^{L-1} \prod_{\nu=1}^{L-1} (\bar{\lambda}_{n_\nu} - \bar{\lambda}_n)^2 & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}, \end{cases} \end{aligned}$$

and

$$(4.20) \quad \begin{cases} r^\epsilon(\mathfrak{o}_{2n}; \bar{\Theta}, \lambda) = \prod_{k=1}^L \prod_{n_{k-1} < i < n_k} r_i^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda), \\ r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) = r^\epsilon(\mathfrak{o}_{2n}; \bar{\Theta}, \lambda) \bar{r}_o^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) & \text{if } \mathfrak{g} = \mathfrak{sp}_n \text{ or } \mathfrak{o}_{2n+1}. \end{cases}$$

Proof. Define the parameters μ_ν in Proposition 3.4 by

$$\begin{aligned} \mu_\nu &= \lambda_\nu + n_{\nu-1} \quad \text{for } 1 \leq \nu \leq L, \\ \mu_{\nu+L} &= -\lambda_\nu - n_\nu + 2n + \delta = -\mu_\nu - n'_\nu + 2n + \delta \quad \text{for } 1 \leq \nu \leq L, \\ \mu_{2L+1} &= -n. \end{aligned}$$

Then in the proposition $H(\Theta, \mu_1, \dots, \mu_L) = \prod_{\nu=1}^{L'} (F - \mu_\nu)$ with $L' = L, 2L, 2L$ or $2L + 1$ if $\mathfrak{g} = \mathfrak{gl}_n, \mathfrak{sp}_n, \mathfrak{o}_{2n}$ or \mathfrak{o}_{2n+1} , respectively. Moreover $\tilde{H}(\Theta, \mu_1, \dots, \mu_{L-1}) = \prod_{\nu=1}^{2L-1} (F - \mu_\nu)$ with $\lambda_L = 0$. Note that if $\ell \leq L$,

$$\begin{aligned} \mu_\ell - \mu_\nu - n'_\nu - n_{\ell-1} + i &= \lambda_\ell + i - \lambda_\nu - n_\nu, \\ \mu_\ell - \mu_\nu - n_{\ell-1} + i &= \lambda_\ell + i - \lambda_\nu - n_{\nu-1}, \\ \mu_\ell - \mu_{\nu+L} - n_{\ell-1} + i &= \lambda_\ell + i + \lambda_\nu + n'_\nu - 2n - \delta, \\ \mu_\ell - \mu_{2L+1} - n_{\ell-1} + i &= \lambda_\ell + i + n. \end{aligned}$$

For $\Theta' = \Theta$ or $\bar{\Theta}$ Proposition 3.4 and the isomorphism $U(\mathfrak{g}) \simeq U^\epsilon(\mathfrak{g})$ show that

$$q_{\Theta'}^\epsilon(\mathfrak{g}; F, \lambda)_{ij} \equiv 0 \pmod{\bar{n}U^\epsilon(\mathfrak{g}) + J^\epsilon(\lambda_{\Theta'})},$$

which assures $I_{\Theta'}^\epsilon(\lambda) \subset \text{Ann}(M_{\Theta'}^\epsilon(\lambda))$ (cf. [O4, Lemma 2.11 and Remark 2.12]) because $M_{\Theta'}^\epsilon(\lambda)$ is irreducible \mathfrak{g} -module for generic (λ, ϵ) and $\sum \mathbb{C}q_{\Theta'}^\epsilon(\mathfrak{g}; F, \lambda)_{ij}$ is \mathfrak{g} -invariant.

Suppose $\mathfrak{g} = \mathfrak{o}_{2n}$. Since $I_{\bar{\Theta}}^\epsilon + J^\epsilon(\lambda_{\Theta})$ is stable under the outer automorphism which maps α_{n-1} to α_n , $I_{\bar{\Theta}}^\epsilon + J^\epsilon(\lambda_{\Theta}) \ni F_{ii+1}$ if and only if $I_{\bar{\Theta}}^\epsilon + J^\epsilon(\lambda_{\Theta}) \ni F_{ii+2}$.

Hence other statements of the theorem are direct consequences of Proposition 3.4. Note that the functions in the theorem should be homogeneous with respect to $(x, \mathfrak{g}, \lambda, \epsilon)$. \square

Any zero of $r^\epsilon(\mathfrak{g}; \Theta', \lambda)$ in the above theorem corresponds to the hypersurface defined by a root in $\Sigma(\mathfrak{g})$ except for the term $\bar{\lambda}_{i+1} + \bar{\lambda}_{n_\nu}$ with $i = n_\nu - 1$ and $\nu = k$ in the case when $\mathfrak{g} = \mathfrak{o}_{2n}$. Hence we have the following remark.

Remark 4.5. i) Suppose $\mathfrak{g} = \mathfrak{gl}_n$. Considering the weights with respect to \mathfrak{a} , we have

$$\left(I_{\bar{\Theta}}^\epsilon(\lambda) + J^\epsilon(\lambda_{\Theta}) \right) \cap \mathbb{C}E_{ii+1} = \left(q_{\bar{\Theta}}^\epsilon(\mathfrak{g}; F, \lambda)_{ii+1} + J^\epsilon(\lambda_{\Theta}) \right) \cap \mathbb{C}E_{ii+1}$$

as in the argument in [O4, §3]. Hence $J_{\bar{\Theta}}^\epsilon(\lambda) = I_{\bar{\Theta}}^\epsilon(\lambda) + J^\epsilon(\lambda_{\Theta})$ if and only if $r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) \neq 0$.

ii) If the infinitesimal character of $M^\epsilon(\lambda_{\Theta})$ is regular (resp. strongly regular) in the case when $\mathfrak{g} = \mathfrak{gl}_n, \mathfrak{o}_{2n+1}$ or \mathfrak{sp}_n (resp. \mathfrak{o}_{2n}), then $r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) \neq 0$ and $r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) \neq 0$.

Here we defined that the infinitesimal character μ is *strongly regular* if it is not fixed by any non-trivial element of the group of automorphisms of the root system. Note that the group is generated by the reflections with respect to the simple roots and the automorphisms of the Dynkin diagram.

iii) Suppose $\mathfrak{g} = \mathfrak{o}_{2n}$. Then $r^\epsilon(\mathfrak{g}; \bar{\Theta}, \lambda) \neq 0$ if

$$(4.21) \quad \begin{aligned} &\langle \lambda_{\Theta}|_{\mathfrak{a}} + \epsilon\rho, \alpha \rangle \neq 0 \quad \text{for } \alpha \in \Sigma(\mathfrak{a}), \\ &\langle \lambda_{\Theta}|_{\mathfrak{a}} + \epsilon\rho, 2\alpha_j + \cdots + 2\alpha_{n-2} + \alpha_{n-1} + \alpha_n \rangle \neq 0 \quad \text{for } j = 2, \dots, n-1 \end{aligned}$$

under the notation in (4.3).

It is proved by [BG] and [Jo] that for $\mu \in \mathfrak{a}^*$ the map

$$(4.22) \quad \begin{aligned} &\{I; I \text{ are the two-sided ideals of } U(\mathfrak{g}) \text{ with } I \supset \text{Ann}(M(\mu))\} \\ &\ni I \mapsto I + J(\mu) \in \{J; J \text{ are the left ideals of } U(\mathfrak{g}) \text{ with } J \supset J(\mu)\} \end{aligned}$$

is injective if $\mu + \rho$ is dominant

$$(4.23) \quad 2 \frac{\langle \mu + \rho, \alpha \rangle}{\langle \alpha, \alpha \rangle} \notin \{-1, -2, \dots\} \quad \text{for any root } \alpha \text{ for the pair } (\mathfrak{n}, \mathfrak{a}).$$

Under the notation (4.9) with $\mu = \lambda_{\Theta}|_{\mathfrak{a}}$ and $\epsilon = 1$ the condition (4.23) is equal to

$$(4.24) \quad \begin{cases} \bar{\lambda}_i - \bar{\lambda}_j \notin \{1, 2, \dots\} & (1 \leq i < j \leq n) & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ \bar{\lambda}_i \pm \bar{\lambda}_j \notin \{1, 2, \dots\} & (1 \leq i < j \leq n) & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}, \\ \bar{\lambda}_i \pm \bar{\lambda}_j, \bar{\lambda}_k \notin \{1, 2, \dots\} & (1 \leq i < j \leq n, 1 \leq k \leq n) & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ \bar{\lambda}_i \pm \bar{\lambda}_j, 2\bar{\lambda}_k \notin \{1, 2, \dots\} & (1 \leq i < j \leq n, 1 \leq k \leq n) & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}. \end{cases}$$

Note that

$$(4.25) \quad \bar{\lambda}_i - \bar{\lambda}_{i+1} = -1 \quad \text{if } n_{k-1} < i < n_k \text{ and } 1 \leq k \leq L.$$

Corollary 4.6. i) If $\lambda_{\Theta}|_{\mathfrak{a}} + \rho$ is dominant and $r^1(\mathfrak{g}; \bar{\Theta}', \lambda) \neq 0$, then

$$(4.26) \quad \text{Ann}(M_{\bar{\Theta}'}(\lambda)) = I_{\bar{\Theta}'}^1(\lambda)$$

for $\Theta' = \Theta$ or $\bar{\Theta}$.

ii) Suppose the infinitesimal character $M_{\Theta'}(\lambda)$ is regular (resp. strongly regular) in the case when $\mathfrak{g} = \mathfrak{o}_{2n}$ with $\Theta' = \bar{\Theta}$ or $\mathfrak{g} = \mathfrak{gl}_n$ or $\mathfrak{g} = \mathfrak{o}_{2n+1}$ or $\mathfrak{g} = \mathfrak{sp}_n$ (resp. $\mathfrak{g} = \mathfrak{o}_{2n}$ with $\Theta' = \Theta$). Then (4.26) holds.

Proof. i) Since $J_{\Theta'}(\lambda) = I_{\Theta'}(\lambda) + J(\lambda_{\Theta'}) \subset \text{Ann}(M_{\Theta'}(\lambda)) + J(\lambda_{\Theta'}) \subset J_{\Theta'}(\lambda)$ by Theorem 4.4, we have (4.26) by the injectivity of (4.22).

ii) Let $\Psi(\mathfrak{m}_{\Theta}, \mathfrak{a})$ denote the fundamental system for $(\mathfrak{m}_{\Theta}, \mathfrak{a})$. Here $\Psi(\mathfrak{m}_{\Theta}, \mathfrak{a}) \subset \Psi(\mathfrak{a})$. Fix $w \in W$ satisfying $w\Psi(\mathfrak{m}_{\Theta}, \mathfrak{a}) \subset \Psi(\mathfrak{a})$. Define $w.\Theta'$ and $w.\lambda$ so that $\mathfrak{m}_{w.\Theta'} = \text{Ad}(w)\mathfrak{m}_{\Theta'}$ and $\overline{w.\lambda} = w\bar{\lambda} \in \mathbb{C}^n$. Then [Ja, Corollary 15.27] says

$$(4.27) \quad \text{Ann}(M_{\Theta'}(\lambda)) = \text{Ann}(M_{w.\Theta'}(w.\lambda)).$$

For example, $\text{Ann}(M_{\{k,n\}}(\lambda_1, \lambda_2)) = \text{Ann}(M_{\{n-k,n\}}(\lambda_2 + k, \lambda_1 - n + k))$. Here we note that $q_{\Theta'}(\mathfrak{g}; x, \lambda)$ does not change under this transformation.

Case $\mathfrak{g} = \mathfrak{gl}_n$. By a permutation of the L blocks $\{\{n_{k-1} + 1, \dots, n_k\}; k = 1, \dots, L\}$, we may assume $\Re\bar{\lambda}_{n_0+1} \leq \Re\bar{\lambda}_{n_1+1} \leq \dots \leq \Re\bar{\lambda}_{n_{L-1}+1}$. Here $\Re c$ denotes the real part of c for $c \in \mathbb{C}$. Since $M_{\Theta}(\lambda)$ has a regular infinitesimal character, (4.24) and (4.25) assure that $\lambda_{\Theta}|_{\mathfrak{a}} + \rho$ is dominant and $r^1(\mathfrak{g}; \Theta, \lambda) \neq 0$. Hence we have ii) from i).

Case $\mathfrak{g} = \mathfrak{sp}_n$ or \mathfrak{o}_{2n+1} . First suppose $\Theta' = \Theta$. We may assume $\Re\bar{\lambda}_{n_0+1} \leq \Re\bar{\lambda}_{n_1+1} \leq \dots \leq \Re\bar{\lambda}_{n_{L-1}+1} \leq 0$. Since $M_{\Theta}(\lambda)$ has a regular infinitesimal character, $\lambda_{\Theta}|_{\mathfrak{a}} + \rho$ is dominant and we have ii). Here we note that if $\bar{\lambda}_{n_{k+1}} < 0$ and $\bar{\lambda}_{n_{k+1}} > 0$, then $\bar{\lambda}_{n_{k+1}} \notin 2\mathbb{Z}$ because $\bar{\lambda}_i + \bar{\lambda}_{i+1} \neq 0$ if $n_k < i < n_{k+1}$.

Next suppose $\Theta' = \bar{\Theta}$. Then we may assume $\Re\bar{\lambda}_{n_0+1} \leq \Re\bar{\lambda}_{n_1+1} \leq \dots \leq \Re\bar{\lambda}_{n_{L-2}+1} \leq 0$ and $\bar{\lambda}_{n_L} = -\frac{\delta+1}{2}$ (cf. (3.14)). Hence we similarly have ii).

Case $\mathfrak{g} = \mathfrak{o}_{2n}$. The map $e_n \mapsto -e_n$ corresponds to an outer automorphism of \mathfrak{o}_{2n} which does not change \mathfrak{b} . Combining the corresponding automorphism of the universal enveloping algebra with the above argument, we may have the same assumption on $\bar{\lambda}$ as in the previous case. Thus we similarly have ii). \square

Remark 4.7. Suppose $\mathfrak{g} = \mathfrak{gl}_n$.

i) In [O4] another generator system of $\text{Ann}_G(M_{\bar{\Theta}}^{\epsilon}(\lambda))$ is given for every $(\Theta, \epsilon, \lambda)$. It is interesting to express them by the generators constructed in this note, which is done by [Sa] when \mathfrak{p}_{Θ} is a maximal parabolic subalgebra. In the case of the minimal parabolic subalgebra, that is, in the case of the central elements of $U(\mathfrak{g})$, it is studied by [I1], [I2] and [Um]. In general, it may be considered as a generalization of Newton's formula for symmetric polynomials.

ii) Put $\Theta = \{k, n\}$ with $1 \leq k < n$ and fix $\lambda = (\lambda_1, \lambda_2) \in \mathbb{C}^2$. Then

$$\lambda \text{ is regular} \Leftrightarrow \lambda_1 - \lambda_2 \notin \{1, 2, \dots, n-1\},$$

$$\lambda \text{ is dominant} \Leftrightarrow \lambda_1 - \lambda_2 \notin \{2, 3, 4, \dots\},$$

$$J_{\Theta}(\lambda) = I_{\Theta}(\lambda) + J(\lambda_{\Theta}) \Leftrightarrow \lambda_1 - \lambda_2 \notin \{1, 2, \dots, \max\{k, n-k\} - 1\},$$

$$J_{\Theta}(\lambda) = \text{Ann}(M_{\Theta}(\lambda)) + J(\lambda_{\Theta}) \Leftrightarrow \lambda_1 - \lambda_2 \notin \{1, 2, \dots, \min\{k, n-k, \frac{n}{2} - 1\}\}.$$

Here the last equivalence follows from [O4, Theorem 3.1] and the one before last is clear from Remark 4.5 i).

Remark 4.8. Considering the m -th exterior product of the natural representation of \mathfrak{gl}_n , we may put $p(E) = (E_{IJ})_{\#I=\#J=m} \in M\left(\binom{n}{m}, U(\mathfrak{g})\right)$ in Lemma 2.1, where $I = \{i_1, \dots, i_m\}$, $J = \{j_1, \dots, j_m\}$ with $1 \leq i_1 < \dots < i_m \leq n$ and $1 \leq j_1 < \dots < j_m \leq n$ and $E_{IJ} = \det(E_{i_{\mu}j_{\nu}} + (\mu - m)\epsilon\delta_{i_{\mu}j_{\nu}})_{1 \leq \mu \leq m, 1 \leq \nu \leq m}$. Here

$\det(A_{ij}) = \sum_{\sigma \in \mathfrak{S}_n} A_{\sigma(1)1} \cdots A_{\sigma(n)n}$. The study of $f(p(E))$ for polynomials $f(x)$ may be interesting because it may be a *quantization* of the ideals of the *rank varieties* (cf. [ES]) defined by the condition $\text{rank } f(A) = m$ for $A \in M(n, \mathbb{C})$.

Remark 4.9. For $\mathfrak{g} = \mathfrak{o}_n$ or \mathfrak{sp}_n we may expect an explicit generator system for $\text{Ann}_G(M_{\mathfrak{p}}^{\epsilon}(\lambda))$ which are of the same type given by [O4] for \mathfrak{gl}_n . It should be a *quantization* of determinants and Pfaffians (and elementary divisors for the singular case). The quantization of Pfaffians for \mathfrak{o}_n is studied by [I2], [IU] and [Od] etc. It is shown by [Od] that it gives $\text{Ann}(M_{\mathfrak{p}}(\lambda))$ for the expected \mathfrak{p} .

Remark 4.10. We have considered $\sum_{i,j} \mathbb{C}f(p(E))_{ij}$ for the construction of a two-sided ideal of $U(\mathfrak{g})$ with a required property. We may pick up a \mathfrak{g} -invariant subspace V of $\sum_{i,j} \mathbb{C}f(p(E))_{ij}$ to get a refined result. Moreover for a certain problem (cf. [O1]) related to a symmetric pair $(\mathfrak{g}, \mathfrak{k})$ it is useful to study \mathfrak{k} -invariant subspaces of $\sum_{i,j} \mathbb{C}f(p(E))_{ij}$ which should have required zeros under the map of Harish-Chandra homomorphism for the pair. This will be discussed in another paper [OSh].

In the case when $\epsilon = 0$ we have the following.

Theorem 4.11. *Let $\lambda \in \mathfrak{a}$ and suppose that the centralizer of λ in \mathfrak{g} equals $\mathfrak{m}_{\Theta'}$ with $\Theta' = \Theta$ or $\bar{\Theta}$. If $\mathfrak{g} = \mathfrak{o}_{2n}$, we moreover assume $\lambda_i \neq 0$ for $i = 1, \dots, n$. Then*

$$I_{\Theta'}^0(\lambda) = \{f \in S(\mathfrak{g}); f|_{\text{Ad}(G)\lambda} = 0\}.$$

Proof. It is clear from Theorem 4.4 that the element of $I_{\Theta'}^0(\lambda)$ vanishes on λ and therefore $I_{\Theta'}^0(\lambda)$ vanishes on $\text{Ad}(G)\lambda$ because $I_{\Theta'}^0(\lambda)$ is G -stable.

We will prove that the dimension of the space $\sum_{i=1}^N \sum_{j=1}^N \mathbb{C}dq_{\Theta'}^0(\mathfrak{g}; F, \lambda)_{ij}|_{\lambda}$ is not smaller than $\dim \mathfrak{m}_{\Theta'}$. This is shown by the direct calculation and it is almost the same in any case and therefore we give it in the case when $\mathfrak{g} = \mathfrak{sp}_n$ and $\Theta' = \bar{\Theta}$.

Put $\Theta = \{n_1, \dots, n_L\}$ and $\lambda = (\lambda_1, \dots, \lambda_L)$. Note that $\lambda_L = 0$ and $q_{\bar{\Theta}}^0(\mathfrak{sp}_n; x, \lambda) = x \prod_{1 \leq \nu < L} (x - \lambda_{\nu})(x + \lambda_{\nu})$. If $n_{k-1} \leq i < n_k$ and $n_{k-1} \leq j < n_k$ and $k < L$, we have

$$dq_{\bar{\Theta}}^0(\mathfrak{sp}_n; F, \lambda)_{ij}|_{\lambda_{\Theta}} = 2\lambda_k^2 \prod_{1 \leq \nu < L, \nu \neq k} (\lambda_k - \lambda_{\nu})(\lambda_k + \lambda_{\nu}) dF_{ij}.$$

If $n_{L-1} \leq i < 2n - n_{L-1}$ and $n_{L-1} \leq j < 2n - n_{L-1}$, then

$$dq_{\bar{\Theta}}^0(\mathfrak{sp}_n; F, \lambda)_{ij}|_{\lambda_{\Theta}} = \prod_{1 \leq \nu < L} (-\lambda_{\nu})(\lambda_{\nu}) dF_{ij}.$$

The assumption of the proposition implies $\lambda_k \neq 0$, $\lambda_{\nu} \neq 0$ and $\lambda_k^2 \neq \lambda_{\nu}^2$ in the above and therefore we get the required result.

Put $V = \{X \in \mathfrak{g}; f(X) = 0 (\forall f \in I_{\Theta'}(\lambda))\}$. Since $[\lambda, \mathfrak{g}] = \mathfrak{n}_{\Theta} + \bar{\mathfrak{n}}_{\Theta}$, the tangent space of $\text{Ad}(G)\lambda$ at λ is isomorphic $\mathfrak{n}_{\Theta} + \bar{\mathfrak{n}}_{\Theta}$. Since $\text{Ad}(G)\lambda \subset V$, it follows from the above calculation of the dimension that $\text{Ad}(G)\lambda$ and V are equal in a neighborhood of λ . In particular, V is non-singular at λ .

Let $X \in \mathfrak{g}$ with $f(X) = 0$ for all $f \in I_{\Theta'}^0(\lambda)$. We will show $X \in \text{Ad}(G)\lambda$, which completes the proof of the theorem. Let $X = X_s + X_n$ be the Jordan decomposition of X . Here X_s is semisimple and X_n is nilpotent. By the action of the element of $\text{Ad}(G)$, we may assume $X_s \in \mathfrak{a}$ and $X_n \in \mathfrak{n}$. Then it is clear that $f(X_s + tX_n) = 0$ for all $f \in I_{\Theta'}^0(\lambda)$ and $t \in \mathbb{C}$. Moreover it is also clear that X_s is a transformation of λ_{Θ} under a suitable element of the Weyl group of the root system for the pair $(\mathfrak{g}, \mathfrak{a})$ and therefore we may assume $X_s = \lambda$. Since the tangent space of V and λ is isomorphic to $\mathfrak{n}_{\Theta} + \bar{\mathfrak{n}}_{\Theta}$, we have $X_n \in \mathfrak{n}_{\Theta}$. Hence $X_n = 0$ because $[X_s, X_n] = 0$. and therefore $X \in \text{Ad}(G)\lambda$. \square

Remark 4.12. Theorem 4.11 shows that we have constructed a generator system of the defining ideal of the adjoint orbit of any semisimple element of any classical Lie algebra. In fact, for any $\lambda \in \mathfrak{a}$ in the orbit the centralizer of λ in \mathfrak{g} is \mathfrak{m}_{Θ} or $\mathfrak{m}_{\bar{\Theta}}$ or \mathfrak{g} with a suitable Θ .

On the other hand [O4] constructed a generator system of the ideal corresponding to the closure of an arbitrary conjugacy class of \mathfrak{gl}_n , which is of a different type from the one given here.

We will generalize the Cayley-Hamilton theorem in the linear algebra. Put

$$\bar{d}_{\mathfrak{g}}(x) = \begin{cases} \prod_{i=1}^n (x - F_i - \frac{n-1}{2}\epsilon) & \text{if } \mathfrak{g} = \mathfrak{gl}_n, \\ \prod_{i=1}^n (x - F_i - n\epsilon)(x + F_i - n\epsilon) & \text{if } \mathfrak{g} = \mathfrak{sp}_n, \\ \prod_{i=1}^n (x - F_i - (n-1)\epsilon)(x + F_i - (n-1)\epsilon) & \text{if } \mathfrak{g} = \mathfrak{o}_{2n}, \\ (x - n\epsilon) \prod_{i=1}^n (x - F_i - (n - \frac{1}{2})\epsilon)(x + F_i - (n - \frac{1}{2})\epsilon) & \text{if } \mathfrak{g} = \mathfrak{o}_{2n+1}. \end{cases}$$

Here we note that if $\Theta = \{1, 2, \dots, n\}$, then $n_j = j$, $\lambda_i + n_{i-1}\epsilon = \bar{\lambda}_i + (n + \frac{\delta-1}{2})\epsilon$ and $\lambda_i + (n_i - 2n - \delta)\epsilon = \bar{\lambda}_i - (n + \frac{\delta-1}{2})\epsilon$. Since $\bar{d}_{\mathfrak{g}}(x) \in S(\mathfrak{a})^W[x]$, there exists $d_{\mathfrak{g}}(x) \in Z^\epsilon(\mathfrak{g})[x]$ with

$$(4.28) \quad \bar{\omega}(d_{\mathfrak{g}}(x)) = \bar{d}_{\mathfrak{g}}(x),$$

which is equivalent to $d_{\mathfrak{g}}(x) \equiv \bar{d}_{\mathfrak{g}}(x)(\mu) \pmod{J^\epsilon(\mu - \epsilon\rho)}$. Then Theorem 4.4 assures $d_{\mathfrak{g}}(F) \equiv \bar{d}_{\mathfrak{g}}(F)(\mu) \equiv 0 \pmod{J^\epsilon(\mu - \epsilon\rho)}$. Hence $\bar{\omega}(d_{\mathfrak{g}}(F))(\mu) = 0$ for any $\mu \in \mathfrak{a}^*$ and therefore $\bar{\omega}(d_{\mathfrak{g}}(F)) = 0$, which assures $d_{\mathfrak{g}}(F) = 0$ because $\sum_{i,j} \mathbb{C}d_{\mathfrak{g}}(F)_{ij}$ is \mathfrak{g} -invariant (cf. [O4, Lemma 2.12]). Thus we have the following corollary.

Corollary 4.13 (The Cayley-Hamilton theorem for the natural representation of the classical Lie algebra \mathfrak{g}).

$$d_{\mathfrak{g}}(F) = 0.$$

Remark 4.14. This result for \mathfrak{gl}_n and \mathfrak{o}_n is given by [Um] and [I2], respectively. A much general result is given by [Go2] (cf. [OO]).

Remark 4.15. Suppose $\mathfrak{g} = \mathfrak{gl}_n$. Then it follows from [O3] that

$$d_{\mathfrak{g}}(x) = \det \left(x - F_{ij} - (i-n)\epsilon\delta_{ij} \right)_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}}.$$

In [O4] we define another generator system of $\text{Ann}_G(M_{\Theta}^\epsilon(\lambda))$ for any $(\Theta, \lambda, \epsilon)$ by using “elementary divisors” in place of the “minimal polynomial” $q_{\Theta}^\epsilon(\mathfrak{gl}_n; x, \lambda)$.

Proposition 4.16. *Suppose $\mathfrak{g} = \mathfrak{gl}_n$ and let π be its natural representation. Then the characteristic polynomial of $F = (E_{ij})$ in $U^\epsilon(\mathfrak{g})$ equals $d_{\mathfrak{g}}^\epsilon(x)$ and the minimal polynomial of $(F, M_{\Theta}^\epsilon(\lambda))$ equals $q_{\Theta}^\epsilon(\mathfrak{g}; x, \lambda)$.*

Proof. Suppose $\epsilon = 0$ and identify the dual space \mathfrak{g}^* of \mathfrak{g} with \mathfrak{g} by the bilinear form (2.1). Put $J_{\Theta}^0(\lambda)^\perp = \{X \in \mathfrak{g}^*; \langle X, Y \rangle = 0 \ (\forall Y \in J_{\Theta}^0(\lambda))\}$. Then the condition $q(F)M_{\Theta}^0(\lambda) = 0$ for a polynomial $q(x)$ is equivalent to $q(F)(J_{\Theta}^0(\lambda)^\perp) = 0$, which also equivalent to $q(A_{\Theta, \lambda}) = 0$ for a generic element $A_{\Theta, \lambda}$ of $J_{\Theta}^0(\lambda)^\perp$ because the closure of $\bigcup_{g \in GL(n, \mathbb{C})} gA_{\Theta, \lambda}g^{-1}$ equals $\bigcup_{g \in GL(n, \mathbb{C})} g(J_{\Theta}^0(\lambda)^\perp)g^{-1}$ (cf. [O4, §2]). In fact

$$A_{\Theta, \lambda} = \begin{pmatrix} \lambda_1 I_{n'_1} & & & & 0 \\ A_{21} & \lambda_2 I_{n'_2} & & & \\ A_{31} & A_{32} & \lambda_3 I_{n'_3} & & \\ \vdots & \vdots & \vdots & \ddots & \\ A_{L1} & A_{L2} & A_{L3} & \cdots & \lambda_L I_{n'_L} \end{pmatrix}$$

with generic $A_{ij} \in M(n'_i, n'_j, \mathbb{C})$. Hence our minimal polynomial is the same as that of $A_{\Theta, \lambda}$ in the linear algebra and the claim in the lemma for the minimal polynomial is clear.

We may assume $\epsilon = 1$. Let $p(x)$ be the minimal polynomial of $(F, M_\Theta(\lambda))$ with a fixed λ . Define a homogeneous and monic polynomial $p(x, \epsilon)$ of (x, ϵ) with $p(x) = p(x, 1)$. Then $p(x, \epsilon)M_\Theta^\epsilon(\epsilon\lambda) = 0$ for $\epsilon \in \mathbb{C}$. It follows from the result in the case $\epsilon = 0$ that the degree of $p(x)$ should not be smaller than that of $q_\Theta(\mathfrak{g}; x, \lambda)$. Hence $q_\Theta(\mathfrak{g}; x, \lambda)$ is the minimal polynomial for $M_\Theta^\epsilon(\lambda)$.

Since the degree of the minimal polynomial $q_\Theta(\mathfrak{g}; x, \lambda)$ for $\Theta = \{1, 2, \dots, n\}$ is n , the degree of the characteristic polynomial is not smaller than n . Hence $d_{\mathfrak{g}}(x)$ is the characteristic polynomial. \square

Definition 4.17. The non-zero element $q(x, \lambda, \epsilon) \in \mathbb{C}[x, \lambda, \epsilon] \simeq S(\mathfrak{a}_{\Theta'})[x, \epsilon]$ is called the *global minimal polynomial* of $(F, M_{\mathfrak{p}}(\lambda))$ if $q(x, \lambda, \epsilon)$ satisfies $q(F, \lambda, \epsilon)M_{\mathfrak{p}}(\lambda) = 0$ for any (λ, ϵ) in the parameter space $\mathfrak{a}_{\Theta'}^* \times \mathbb{C}$ and any other non-zero polynomial whose degree with respect to x is smaller than that of $q(x, \lambda, \epsilon)$ does not satisfy this.

Proposition 4.18. *The polynomials $q_{\Theta'}^\epsilon(\mathfrak{g}; x, \lambda)$ in Definition 4.2 are the global minimal polynomials of $(F, M_{\Theta'}(\lambda))$ for $\Theta' = \Theta$ and $\bar{\Theta}$.*

Proof. Let $q(x, \lambda, \epsilon)$ be a global minimal polynomial of $(F, M_{\Theta'}(\lambda))$. We may assume $q(x, \lambda, 0)$ is not zero by dividing by ϵ^k with an integer k if necessary. Put $\epsilon = 0$ and consider the generic λ . Since $q(\langle F, \lambda \rangle, \lambda, 0) = 0$, the minimality of the degree is clear. Here $\langle F, \lambda \rangle = (\lambda(F_i)\delta_{ij})_{\substack{1 \leq i \leq N \\ 1 \leq j \leq N}}$ is a diagonal matrix in $M(N, \mathbb{C})$. \square

Let $f(x)$ be any polynomial in $\mathbb{C}[x]$. We will give a characterization of the image $\bar{\omega}(f(F))$ under the Harish-Chandra homomorphism $\bar{\omega}$ defined by (4.10) in the case of the natural representation of $\mathfrak{g} = \mathfrak{gl}_n$.

Theorem 4.19. *Put $f(x) = \prod_{i=1}^m (x - \lambda_i)$. For sets A and B we denote by $\text{Map}(A, B)$ the totality of the maps of A to B . For $\tau \in \text{Map}(\{1, \dots, n\}, \{1, \dots, m\})$ we put*

$$\begin{cases} I_\tau(\lambda) = \sum_{j=1}^n \mathbb{C}[\mathfrak{a}, \lambda, \epsilon] \left(E_{jj} - \lambda_{\tau(j)} - (m_\tau(j) - \frac{n-1}{2})\epsilon \right), \\ m_\tau(j) = \#\{\nu \in \{1, \dots, j-1\}; \tau(\nu) = \tau(j)\} \quad \text{for } j = 1, \dots, n. \end{cases}$$

Then for $F = (E_{ij})_{\substack{1 \leq i \leq n \\ 1 \leq j \leq n}} \in M(n, \mathfrak{gl}_n)$ we have

$$(4.29) \quad \sum_{j=1}^n \mathbb{C}[\mathfrak{a}, \lambda, \epsilon] \bar{\omega}(f(F))_{jj} = \bigcap_{\tau \in \text{Map}(\{1, \dots, n\}, \{1, \dots, m\})} I_\tau(\lambda).$$

Moreover $\bar{\omega}(f(F)_{ij}) = 0$ if $i \neq j$ and the polynomial $\bar{\omega}(f(F))_{jj}$ is the unique homogeneous element in $\mathbb{C}[\mathfrak{a}, \lambda, \epsilon]$ belonging to the right hand side of (4.29) such that $\bar{\omega}(f(F)_{jj})|_{\lambda=\epsilon=0} = E_{jj}^m$.

Proof. First note that $\bar{\omega}(f(F)_{ij})|_{\epsilon=0} = \delta_{ij} \prod_{\nu=1}^m (E_{jj} - \lambda_\nu)$ and $\bar{\omega}(f(F)_{jj})$ are of homogeneous of degree m with respect to $(\mathfrak{a}, \lambda, \epsilon)$.

We have already proved that $\bar{\omega}(f(F)_{ij}) = 0$ if $i \neq j$ and $\bar{\omega}(f(F)_{jj}) \equiv F_{jj}^m \pmod{U^\epsilon(\mathfrak{a})^{(m-1)}[\lambda]}$. Hence for any fixed λ and ϵ , the system of the equations

$$(4.30) \quad \bar{\omega}(f(F)_{11}) = \dots = \bar{\omega}(f(F)_{nn}) = 0 \text{ for } (E_{11}, \dots, E_{nn}) \in \mathbb{C}^n$$

is in the complete intersection and has m^n roots counting their multiplicities. Hence to prove (4.29) it is sufficient to show that

$$(4.31) \quad \left(\lambda_{\tau(1)} + (m_\tau(1) - \frac{n-1}{2})\epsilon, \dots, \lambda_{\tau(n)} + (m_\tau(n) - \frac{n-1}{2})\epsilon \right)$$

is a root for any $\tau \in \text{Map}(\{1, \dots, n\}, \{1, \dots, m\})$ and a generic $(\lambda, \epsilon) \in \mathbb{C}^{m+1}$.

Fix any $\sigma \in \text{Map}(\{1, \dots, n\}, \{1, \dots, m\})$ which satisfies $\sigma(i) \leq \sigma(j)$ if $1 \leq i < j \leq n$. Define $\Theta = \{n_1, \dots, n_L\}$ so that $0 = n_0 < n_1 < \dots < n_L = n$ and that the condition $\sigma(i) = \sigma(n_k)$ is equals to $n_{k-1} < i \leq n_k$ for $k = 1, \dots, L$. Put $\mu = (\lambda_{\sigma(n_1)} + n_0\epsilon, \dots, \lambda_{\sigma(n_L)} + n_{L-1}\epsilon)$. Then Theorem 4.4 shows that $f(F)_{ij}$ is in $\text{Ann}(M/J_{\Theta}^{\epsilon}(\mu))$. Here we note that $\mu_{\Theta} + \epsilon\rho = \sum_{j=1}^n (\lambda_{\sigma(j)} + (m_{\sigma(j)} + \frac{n-1}{2})\epsilon)e_j$. On the other hand, [O4, Theorem 2.12] determines the common zeros of $\bar{\omega}(\text{Ann}(M/J_{\Theta}^{\epsilon}(\mu)))$ for $\epsilon \neq 0$. Namely they are (4.31) with τ in

$$\{\tau \in \text{Map}(\{1, \dots, n\}, \{1, \dots, m\}); \#\{\nu; \tau(\nu) = \sigma(j)\} = n_j - n_{j-1} \text{ for } j = 1, \dots, L\}.$$

For any τ we have the above σ so that τ is in this set and we have the theorem. \square

5. INTEGRAL TRANSFORMS ON GENERALIZED FLAG MANIFOLDS

Let \mathfrak{g} be a complex reductive Lie algebra and \mathfrak{p} be a parabolic subalgebra containing a Borel subalgebra \mathfrak{b} . For a holomorphic character λ of \mathfrak{p} we define left ideals

$$(5.1) \quad \begin{cases} J_{\mathfrak{p}}(\lambda) = \sum_{X \in \mathfrak{p}} (X - \lambda(X)), \\ J_{\mathfrak{b}}(\lambda) = \sum_{X \in \mathfrak{b}} (X - \lambda(X)) \end{cases}$$

of the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} . Let $I_{\mathfrak{p}}(\lambda)$ be the two-sided ideal of $U(\mathfrak{g})$ which satisfies

$$(5.2) \quad I_{\mathfrak{p}}(\lambda) \subset J_{\mathfrak{p}}(\lambda).$$

Let G be a connected real semisimple Lie group and let P be a parabolic subgroup of G such that the complexifications of $\text{Lie}(G)$ and $\text{Lie}(P)$ equal \mathfrak{g} and \mathfrak{p} , respectively. Let L_{λ} be a line bundle over G/P such that the local section of L_{λ} is killed by $J_{\mathfrak{p}}(\lambda)$. Then the image of any \mathfrak{g} -equivariant map of the space of sections of L_{λ} over an open subset of G/P is killed by $I_{\mathfrak{p}}(\lambda)$. Here the element of $I_{\mathfrak{p}}(\lambda)$ is identified with a left invariant differential operator but it may be identified with a right invariant differential operator through the anti-automorphism of $U(\mathfrak{g})$ ($X \mapsto -X$, $XY \mapsto (-Y)(-X)$ for $X, Y \in \mathfrak{g}$) because $I_{\mathfrak{p}}(\lambda)$ is a two-sided ideal. If the \mathfrak{g} -equivariant map is an integral transform to the space of functions on a homogeneous space X of G or sections of a vector bundle over X , it is a natural question how the system of differential equations induced from $I_{\mathfrak{p}}(\lambda)$ characterizes the image.

The same problem may be considered when L_{λ} is the holomorphic line bundle over the complexification of G/P .

5.1. Penrose transformations. Let $G_{\mathbb{C}}$ be a reductive complex Lie group with the Lie algebra \mathfrak{g} . Let G be a real form of $G_{\mathbb{C}}$ and let $P_{\mathbb{C}}$ be a parabolic subalgebra of $G_{\mathbb{C}}$ with the Lie algebra \mathfrak{p} and let V be a G -orbit in $G_{\mathbb{C}}/P_{\mathbb{C}}$. Suppose \mathcal{O}_{λ} is a holomorphic line bundle over $G_{\mathbb{C}}/P_{\mathbb{C}}$ which is killed by $J_{\mathfrak{p}}(\lambda)$. Here the element of $J_{\mathfrak{p}}(\lambda)$ is identified with a right invariant holomorphic differential operators on $G_{\mathbb{C}}$. Then the image of any G -equivariant map

$$(5.3) \quad T : H_V^*(G_{\mathbb{C}}/P_{\mathbb{C}}, \mathcal{O}_{\lambda}) \rightarrow E$$

is killed by $I_{\mathfrak{p}}(\lambda)$. This is obvious because $I_{\mathfrak{p}}(\lambda)$ is a two-sided ideal. Here E is usually a space of sections of a certain line (or vector) bundle over a homogeneous space of G . In this case $I_{\mathfrak{p}}(\lambda)$ is identified with a system of holomorphic differential equations and we may identify the element of $I_{\mathfrak{p}}(\lambda)$ as a left invariant differential operator on G through the anti-automorphism of the universal enveloping algebra or a right invariant differential operator on G .

5.2. Poisson transformations. Let G be a connected semisimple Lie group with finite center, let K be a maximal compact subgroup of G and let P be a parabolic subalgebra of G with the Langlands decomposition $P = MAN$ and let P_o be a minimal parabolic subgroup with the Langlands decomposition $P_o = M_o A_o N_o$ satisfying $M_o \subset M$, $A_o \supset A$, $N_o \supset N$ and $P_o \subset P$. Let λ be an element of the complexification \mathfrak{a}^* of the dual of the Lie algebra of A and put

$$\mathcal{B}(G/P, L_\lambda) = \{f \in \mathcal{B}(G); f(xman) = a^\lambda f(x) \quad (\forall m \in M, \forall a \in A, \forall n \in N)\}$$

which is the space of hyperfunction sections of spherical degenerate principal series. Let \mathfrak{p} be a complexification of the Lie algebra of P . The Poisson transformation of the space $\mathcal{B}(G/P, L_\lambda)$ is defined by

$$(5.4) \quad \mathcal{P}^\lambda : \mathcal{B}(G/P, L_\lambda) \rightarrow \mathcal{B}(G/K), \quad f \mapsto (\mathcal{P}^\lambda f)(x) = \int_K f(xk) dk$$

with the normalized Haar measure dk on K . Let $\mathbb{D}(G/K)$ be the ring of invariant differential operator of G and let χ_λ be the algebra homomorphism of $\mathbb{D}(G/K)$ to \mathbb{C} so that the image of \mathcal{P}_λ is in the solution space $\mathcal{A}(G/K, \mathcal{M}_\lambda)$ of the system

$$(5.5) \quad \mathcal{M}_\lambda : Du = \chi_\lambda(D)u \quad (\forall D \in \mathbb{D}(G/K))$$

for $u \in \mathcal{A}(G/K)$. Here $\mathcal{A}(G/K)$ denotes the space of real analytic functions on G/K .

Note that $\mathcal{B}(G/P, L_\lambda)$ is the subspace of the space of hyperfunction sections of spherical principal series

$$\mathcal{B}(G/P_o, L_\lambda) = \{f \in \mathcal{B}(G); f(xman) = a^\lambda f(x) \quad (\forall m \in M_o, \forall a \in A_o, \forall n \in N_o)\}.$$

Here λ is extended to the complexification \mathfrak{a}_o^* of the dual of the Lie algebra \mathfrak{a}_o of A_o so that it takes the value 0 on $\text{Lie}(M) \cap \text{Lie}(A_o)$.

Theorem 5.1. *Suppose that the Poisson transform*

$$(5.6) \quad \mathcal{P}_o^\lambda : \mathcal{B}(G/P_o, L_\lambda) \rightarrow \mathcal{A}(G/K, \mathcal{M}_\lambda), \quad f \mapsto (\mathcal{P}_o^\lambda f)(x) = \int_K f(xk) dk$$

for the boundary G/P_o of G/K is bijective. Assume the condition

$$(5.7) \quad J_{\mathfrak{p}}(\lambda) = I_{\mathfrak{p}}(\lambda) + J_{\mathfrak{b}}(\lambda)$$

for a two-sided ideal $I_{\mathfrak{p}}(\lambda)$ of $U(\mathfrak{g})$. Then the Poisson transform \mathcal{P}^λ for the boundary G/P is a G -isomorphism onto the simultaneous solution space of the system \mathcal{M}_λ and the system defined by $I_{\mathfrak{p}}(\lambda)$.

Proof. Since $\mathcal{B}(G/P, L_\lambda)$ is a subspace of $\mathcal{B}(G/P_o, L_\lambda)$ and \mathcal{P}^λ is a G -equivariant map, the image of \mathcal{P}_o^λ satisfies the systems \mathcal{M}_λ and $I_{\mathfrak{p}}(\lambda)$.

Suppose the function $u \in \mathcal{A}(G/K, \mathcal{M}_\lambda)$ satisfies $I_{\mathfrak{p}}(\lambda)$. Since the function $(\mathcal{P}_o^\lambda)^{-1}u \in \mathcal{B}(G/P_o, L_\lambda)$ also satisfies $I_{\mathfrak{p}}(\lambda)$, the condition (5.7) assures $(\mathcal{P}_o^\lambda)^{-1}u \in \mathcal{B}(G/P, L_\lambda)$ because we may assume $\mathbb{C} \otimes_{\mathbb{R}} \text{Lie}(P_o) \supset \mathfrak{b}$. \square

Remark 5.2. i) The above theorem with its proof is based on the idea given by [O3] which explains it in the case when $G = GL(n, \mathbb{R})$.

ii) The bijectivity of \mathcal{P}_o^λ is equivalent to the condition $e(\lambda + \rho) \neq 0$ by [K-]. This condition is introduced by [He] for the injectivity of \mathcal{P}_o^λ . Here

$$(5.8) \quad e(\lambda) = \prod_{\alpha \in \Sigma_o^+} \left\{ \Gamma \left(\frac{\langle \lambda, \alpha \rangle}{2\langle \alpha, \alpha \rangle} + \frac{m_\alpha}{4} + \frac{1}{2} \right) \Gamma \left(\frac{\langle \lambda, \alpha \rangle}{2\langle \alpha, \alpha \rangle} + \frac{m_\alpha}{4} + \frac{m_{2\alpha}}{2} \right) \right\},$$

Σ^+ is the set of the positive roots for the pair $(\mathfrak{g}, \mathfrak{a}_o)$ so that $\text{Lie}(N)$ corresponds to Σ^+ . Moreover $\Sigma_o^+ = \{\alpha \in \Sigma^+; \frac{1}{2}\alpha \notin \Sigma^+\}$, m_α is the multiplicity of the root $\alpha \in \Sigma^+$ and $\rho = \frac{1}{2} \sum_{\alpha \in \Sigma^+} m_\alpha \alpha$.

iii) Suppose G is simple and of the classical type and suppose the condition $e(\lambda + \rho) \neq 0$. Let $I_{\mathfrak{p}}(\lambda)$ be the system given by (4.14). Then if moreover the infinitesimal character of $\mathcal{B}(G/P, L_\lambda)$ is regular (or strongly regular in the case when $\text{Lie}(G)$ equals \mathfrak{o}_{2n} or its real form), \mathcal{P}^λ is G -isomorphic to the solution space of the system of differential equations $I_{\mathfrak{p}}(\lambda)$ on G/K since Theorem 4.4 assures (5.7). This is because the natural map of $Z(\mathfrak{g})$ to $\mathbb{D}(G/K)$ is surjective and therefore it follows from Remark 4.3 i) that \mathcal{M}_λ is contained in $I_{\mathfrak{p}}(\lambda)$. Here the function on G/K is identified with the right K -invariant function on G . Note that all the assumptions are valid when $\lambda = 0$.

iv) Let \mathfrak{p}_o be the complexification of the Lie algebra of P_o and put $J_{\mathfrak{p}_o}(\lambda) = \sum_{X \in \mathfrak{p}_o} (X - \lambda(X))$. Then we may replace the assumption (5.7) by

$$(5.9) \quad J_{\mathfrak{p}}(\lambda) = I_{\mathfrak{p}}(\lambda) + J_{\mathfrak{p}_o}(\lambda)$$

in Theorem 5.1, which is clear from its proof. Suppose that $I_{\mathfrak{p}}(\lambda)$ is the system given by (4.14). Then under the notation in Theorem 4.4, (5.9) is valid if $r_i^\epsilon(\mathfrak{g}; \Theta', \lambda) \neq 0$ for $i \notin \Theta$ and moreover if $\bar{r}^\epsilon(\mathfrak{g}; \Theta', \lambda) \neq 0$ for $\mathfrak{g} = \mathfrak{o}_{2n+1}$ and \mathfrak{sp}_n in the case $\Theta' = \bar{\Theta}$.

v) Owing to [K-] the abstract existence of the system of differential equations characterizing the image of \mathcal{P}^λ is clear (cf. [OSh]) but a certain existence theorem of the system in the case $\lambda = 0$ is given by [Jn]. A more precise study for this problem including the relation to the *Hua equations* will be discussed in [OSh].

5.3. Intertwining operators. Retain the notation in §5.2. Let $\tilde{\lambda}$ and $\tilde{\mu}$ be characters of P and P_0 , respectively. Put

$$(5.10) \quad \mathcal{B}(G/Q, L_\tau) = \{f \in \mathcal{B}(G); f(xq) = \tau(q)^{-1}f(x) \quad (\forall q \in Q)\}$$

for $(Q, \tau) = (P, \tilde{\lambda})$, $(P_0, \tilde{\lambda})$ and $(P_0, \tilde{\mu})$. Let \mathfrak{j} be a Cartan subalgebra of $\text{Lie}(G)$ containing \mathfrak{a} and let λ be an element of the complexification of the dual of \mathfrak{j} . Assume (5.7) and $\tilde{\lambda}(e^H) = e^{-\lambda(H)}$ for $H \in \mathfrak{j}$. If there exists a G -equivariant bijective map

$$\mathcal{T}_{\tilde{\mu}}^{\tilde{\lambda}} : \mathcal{B}(G/P_0, L_{\tilde{\lambda}}) \xrightarrow{\sim} \mathcal{B}(G/P_0, L_{\tilde{\mu}}),$$

$\mathcal{T}_{\tilde{\mu}}^{\tilde{\lambda}}(\mathcal{B}(G/P, L_{\tilde{\lambda}}))$ is identical with the space of solutions of the system of differential equations on $\mathcal{B}(G/P_0, L_{\tilde{\mu}})$ defined by $I_{\mathfrak{p}}(\lambda)$.

5.4. Radon transformations. Let G be a connected real semisimple Lie group and let P_1 and P_2 be maximal parabolic subgroups of G . If there exists a G -equivariant map $\mathcal{R} : \mathcal{B}(G/P_1, L_{\lambda_1}) \rightarrow \mathcal{B}(G/P_2, L_{\lambda_2})$ for certain characters λ_j of P_j under the notation (5.10), the image of \mathcal{R} satisfies the system $I_{\mathfrak{p}}(\lambda)$. Here \mathfrak{p} and λ correspond to P_1 and λ_1 , respectively.

5.5. Hypergeometric functions. Some special cases of Radon or Penrose transformations and their relations to Aomoto-Gelfand hypergeometric functions are discussed in [O3], [Se] and [Ta].

6. CLOSURE OF IDEALS

Now we will consider the non-regular λ which are excluded in Theorem 4.4. We begin with a general consideration.

Definition 6.1. Let M be a C^∞ -manifold and let U be an open subset of \mathbb{C}^ℓ . We denote by $\mathcal{D}'(M)$ the space of distributions on M . Suppose that meromorphic functions $f_1(\lambda), \dots, f_n(\lambda)$ of U with values in $\mathcal{D}'(M)$ are given. Moreover suppose there exists a non-zero holomorphic function $r(\lambda)$ on U such that f_1, \dots, f_n are holomorphic on $U_r = \{\lambda \in U; r(\lambda) \neq 0\}$ and $\dim V_\lambda = m$ for any $\lambda \in U_r$. For $\lambda \in U$ we define

$$\bar{V}_\mu = \{f(0); f \text{ is a holomorphic function on } \{t \in \mathbb{C}; |t| < 1\} \text{ valued in } \mathcal{D}'(M)\}$$

and there exists a holomorphic curve $c : \{t \in \mathbb{C}; |t| < 1\} \rightarrow U$ such that

$$c(t) \in U_r \text{ and } f(t) \in V_{c(t)} \text{ for } 0 < |t| \ll 1 \text{ and } c(0) = \mu\}.$$

We call \bar{V}_μ the *closure* of the holomorphic family of the spaces V_λ ($\lambda \in U_r$) at μ . It follows from [OS, Proposition 2.21] that $\dim \bar{V}_\mu \geq m$. We define that a point $\mu \in U \setminus U_r$ is a *removable* (resp. *un-removable*) *singular point* if $\dim V_\mu = m$ (resp. $\dim V_\mu > m$). Note that $\bar{V}_\lambda = V_\lambda$ if $\lambda \in U_r$, which follows from the last statement in Lemma 6.3 by replacing μ and U_r by λ ($\in U_r$) and $U_r \setminus \{\lambda\}$.

Example 6.2. The origin $\lambda = (\lambda_1, \lambda_2) = 0$ is a removable singular point of $V_\lambda = \mathbb{C}(x + \lambda_1) + \mathbb{C}(\lambda_2 x + \lambda_1 y + \lambda_1^2)$ and an un-removable singular point of $V_\lambda = \mathbb{C}(\lambda_1 x + \lambda_2 y)$.

Lemma 6.3. i) *If μ is a removable singular point of the spaces V_λ , then there exist a neighborhood U_μ of μ and holomorphic functions $h_1(\lambda), \dots, h_m(\lambda)$ on U_μ valued in $\mathcal{D}'(M)$ such that they are linearly independent for any $\lambda \in U_\mu$ and they span V_λ for any $\lambda \in U_\mu \cap U_r$. On the other hand, the existence of $h_j(\lambda)$ ($j = 1, \dots, m$) with these property implies that μ is a removable singular point.*

ii) *If U is convex and there is no un-removable singular point in U , we may choose $U_\mu = U$ in i).*

Proof. i) Suppose $\dim \bar{V}_\mu = m$. We may assume $f_1(\lambda), \dots, f_m(\lambda)$ are linearly independent for a generic point λ in U_r . Fix a curve c to U with $c(0) = \mu$ and $c(t) \in U_r$ for $0 < |t| \ll 1$. Then [OS, Proposition 2.21] assures the existence of holomorphic functions $v_i(t)$ ($1 \leq i \leq m$) on $\{t \in \mathbb{C}; |t| < 1\}$ valued in $\mathcal{D}'(M)$ and a holomorphic curve $c : \{t \in \mathbb{C}; |t| < 1\} \rightarrow U$ such that $c(0) = \mu$, $c(t) \in U_r$ and $v_i(t) \in V_{c(t)}$ for $0 < |t| \ll 1$ and $v_1(t), \dots, v_m(t)$ are linearly independent for any t . Then the set $\{v_1(0), \dots, v_m(0)\}$ is a basis of V_μ . Fix test functions ϕ_1, \dots, ϕ_m so that $\langle v_i(0), \phi_j \rangle = \delta_{ij}$ and put $c_{ij}(\lambda) = \langle f_i(\lambda), \phi_j \rangle$. If $0 < |t| \ll 1$, then $v_i(t), \dots, v_m(t)$ span $V_{c(t)} = \sum_{i=1}^m \mathbb{C}f_i(c(t))$ and therefore $f_i(c(t)) = \sum_{j=1}^m c_{ij}(c(t))v_j(t)$, which means $\det(c_{ij}(\lambda))$ is not identically zero. Let $(d_{ij}(\lambda))$ be the inverse of $(c_{ij}(\lambda))$ and define $h_i(\lambda) = \sum_{j=1}^m d_{ij}(\lambda)f_j(\lambda)$ so that $\langle h_i(c(t)), \phi_j \rangle = \delta_{ij}$ for $0 < |t| \ll 1$.

Suppose $h_k(\lambda)$ has a pole at $\lambda = \mu$. Then there exists a test function ϕ such that $\langle h_k(\lambda), \phi \rangle$ has a pole at μ . Then it follows from Weierstrass' preparation theorem that there exists a curve $c(t)$ as above and moreover $\langle h_k(c(t)), \phi \rangle$ has a pole at the origin. Choose a positive integer ℓ so that the function $\tilde{h}(t) = t^\ell h_k(c(t))$ is holomorphically extends to $t = 0$ and $\tilde{h}(0) \neq 0$. Since $\langle \tilde{h}(t), \phi_j \rangle = t^\ell \delta_{kj}$, $\langle \tilde{h}(0), \phi_j \rangle = 0$ for $j = 1, \dots, m$, which contradicts to the facts $\langle v_i(0), \phi_j \rangle = \delta_{ij}$ because $0 \neq \tilde{h}(0) \in V_\mu = \sum_{i=1}^m \mathbb{C}v_i(0)$ by definition.

Thus we have proved that $h_i(\lambda)$ are holomorphic functions on λ in a neighborhood of U_μ of μ . Since $\langle h_i(\lambda), \phi_j \rangle = \delta_{ij}$, they are the required functions. In fact, $f_i(\lambda) = \sum_{j=1}^m \langle f_i(\lambda), \phi_j \rangle h_j(\lambda)$ for generic λ and therefore $V_\lambda \subset \sum_{j=1}^m \mathbb{C}h_j(\lambda)$ for $\lambda \in U_\mu \cap U_r$.

Now suppose the existence of h_1, \dots, h_m and consider the function f to define \bar{V}_μ in Definition 6.1. Then under the above notation, $f(t) = \sum_{j=1}^m \langle f(t), \phi_j \rangle h_j(c(t))$ for $0 < |t| \ll 1$ and therefore $f(0) = \sum_{j=1}^m \langle f(0), \phi_j \rangle h_j(c(0))$, which means $\dim V_\mu = m$.

ii) The claim in i) reduces the global existence of h_i to the second problem of Cousin and it is solved for the convex open domain by Oka's principle. \square

Remark 6.4. i) Replacing “meromorphic” and “holomorphic” by “rational” and “regular”, respectively, we have also Lemma 6.3 in the algebraic sense.

ii) When M is a finite set in Lemma 6.3, $\mathcal{D}'(M)$ is a finite dimensional vector space V over \mathbb{C} and $f_i(\lambda)$ are the elements of V with a meromorphic parameter λ .

Definition 6.5. Fix a base $\{X_1, \dots, X_m\}$ of \mathfrak{g} . Let

$$(6.1) \quad q_\nu(\lambda, \epsilon) = \sum_{\alpha_1 \geq 0, \dots, \alpha_m \geq 0} q_{\nu, \alpha}(\lambda, \epsilon) X_1^{\alpha_1} \cdots X_m^{\alpha_m}$$

be elements of $U^\epsilon(\mathfrak{g})$ for $(\lambda, \epsilon) \in \mathbb{C}^{r+1}$ and $\nu = 1, \dots, k$. Here $q_{\nu, \alpha}$ are polynomial functions of (λ, ϵ) and $q_{\nu, \alpha} = 0$ if $\alpha_1 + \dots + \alpha_m$ is sufficiently large. Let $I(\lambda, \epsilon)$ is the left ideal of $U^\epsilon(\mathfrak{g})$ generated by q_ν for $\nu = 1, \dots, k$. Put $d_j = \max_{\lambda, \epsilon} \dim I(\lambda, \epsilon) \cap U^\epsilon(\mathfrak{g})^{(j)}$ for $j = 1, 2, \dots$. Then we can find

$$p_{j, \mu}(\lambda, \epsilon) = \sum_{\alpha_1 \geq 0, \dots, \alpha_m \geq 0} p_{j, \mu, \alpha}(\lambda, \epsilon) X_1^{\alpha_1} \cdots X_m^{\alpha_m}$$

such that $p_{j, \mu}(\lambda, \epsilon) \in I(\lambda, \epsilon) \cap U^\epsilon(\mathfrak{g})^{(j)}$ for any (λ, ϵ) , $p_{j, \mu, \alpha}(\lambda, \epsilon)$ are polynomial functions and $p_{j, 1}(\lambda, \epsilon), \dots, p_{j, d_j}(\lambda, \epsilon)$ are linearly independent for generic (λ, ϵ) . Then we denote by $\bar{I}(\lambda, \epsilon)^{(j)}$ the closure of the holomorphic family $\sum_{\mu=1}^{d_j} \mathbb{C} p_{j, \mu}$ at (λ, ϵ) and put $\bar{I}(\lambda, \epsilon) = \bigcup_{j=1}^{\infty} \bar{I}(\lambda, \epsilon)^{(j)}$. We call $\bar{I}(\lambda, \epsilon)$ the closure of the ideal $I(\lambda, \epsilon)$ with respect to the parameter (λ, ϵ) . We call a point $(\lambda, \epsilon) \in \mathbb{C}^{r+1}$ is an un-removable singular point if (λ, ϵ) is an un-removable singular point of $\sum_{\mu=1}^{d_j} \mathbb{C} p_{j, \mu}$ for a certain j . Note that $\bar{I}(\lambda, \epsilon)$ does not depend on the choice of $\{X_1, \dots, X_m\}$ or $p_{j, \mu}$.

Let $\bar{I}_{\Theta'}^\epsilon(\lambda)$ be the closure of the two-sided ideal $I_{\Theta'}^\epsilon(\lambda)$ given by (4.14) for $\Theta' = \Theta$ or $\bar{\Theta}$. Then we give some problems.

Problem 1. Does there exists no un-removable singular point in the parameter (λ, ϵ) of the holomorphic family $I_{\Theta'}^\epsilon(\lambda)$?

Problem 2. Does the equality $\bar{I}_{\Theta'}(\lambda) = \text{Ann}(M_{\Theta'}(\lambda))$ hold?

Remark 6.6. i) It is clear that $\bar{I}_{\Theta'}^1(\lambda) \subset \text{Ann}(M_{\Theta'}(\lambda))$.

ii) The non-existence of the un-removable singularity in Problem 1 is equivalent to the following conditions. We can choose $\{q_\nu(\lambda, \epsilon); \nu = 1, \dots, k\}$ of the form (6.1) such that $\bar{I}_{\Theta'}^\epsilon(\lambda) = \sum_{\nu=1}^k U^\epsilon(\mathfrak{g}) q_\nu(\lambda, \epsilon)$ for any fixed (λ, ϵ) . It is also equivalent to the fact that the graded ring

$$\text{gr}(\bar{I}_{\Theta'}^\epsilon(\lambda)) = \bigoplus_{j=1}^{\infty} \left(\bar{I}_{\Theta'}^\epsilon(\lambda) \cap U^\epsilon(\mathfrak{g})^{(j)} / \bar{I}_{\Theta'}^\epsilon(\lambda) \cap U^\epsilon(\mathfrak{g})^{(j-1)} \right)$$

does not depend on (λ, ϵ) because they are also equivalent to the fact that the dimension of the vector space $\bar{I}_{\Theta'}^\epsilon(\lambda) \cap U^\epsilon(\mathfrak{g})^{(j)}$ does not depend on (λ, ϵ) and the space is spanned by homogeneous elements with respect to $(\mathfrak{g}, \lambda, \epsilon)$.

iii) Problem 1 and Problem 2 are affirmative if $\mathfrak{g} = \mathfrak{gl}_n$ because there exist $q_\nu(\lambda, \epsilon)$ ($\nu = 1, \dots, k$) of the form (6.1) such that $\text{Ann}_G(M_{\Theta'}^\epsilon(\lambda))$ is generated by $q_\nu(\lambda, \epsilon)$ ($\nu = 1, \dots, k$) for any (λ, ϵ) (cf. [O4]). In this case $\text{gr}(I_{\Theta'}^\epsilon(\lambda))$ is a prime ideal of $S(\mathfrak{g})$ but this is not true in general.

iv) If $\text{gr}(I_{\Theta'}^0(\lambda))$ is a prime ideal for generic λ , then Problem 1 and Problem 2 are affirmative, which is proved by the same argument as in [O4]. Note that $I_{\Theta'}^0(\lambda)$ is the defining ideal of $\text{Ad}(G)\lambda$ for generic $\lambda \in \mathfrak{a}_{\Theta'}$ by Theorem 4.11.

v) Problem 2 is affirmative if the infinitesimal character is (strongly) regular.

REFERENCES

- [BG] J. N. Bernstein and S. I. Gelfand, *Tensor products of finite and infinite dimensional representations of semisimple Lie algebras*, Comp. Math. **41**(1980), 245–285.
- [C1] A. Capelli, *Über die Zurückführung der Cayley'schen Operation Ω auf gewöhnliche Polar Operationen*, Math. Ann. **29** (1887), 331–338.
- [C2] ———, *Sur les opérations dans la théorie des formes algébriques*, Math. Ann. **37**(1890), 1–37.

- [ES] D. Eisenbud and D. Saltman, *Rank variety of matrices*, Commutative algebra, Math. Sci. Res. Inst. Publ. **15**, 173–212, Springer-Verlag, 1989.
- [Ge] I. M. Gelfand, *Center of the infinitesimal groups*, Mat. Sb. Nov. Ser. **26**(68)(1950), 103–112; English transl. in “Collected Papers”, Vol. II, pp.22–30.
- [Go1] M. D. Gould, *A trace formula for semi-simple Lie algebras*, Ann. Inst. Henri Poincaré, Sect. A **32**(1980), 203–219.
- [Go2] ———, *Characteristic identities for semi-simple Lie algebras*, J. Austral Math. Ser. B **26**(1985), 257–283.
- [He] S. Helgason, *A duality for symmetric spaces with applications to group representations II*, Advances in Math. **22**(1976), 187–219.
- [I1] M. Itoh, *Explicit Newton’s formula for \mathfrak{gl}_n* , J. Alg. **208**(1998), 687–697.
- [I2] ———, *Capelli elements for the orthogonal Lie algebras*, J. Lie Theory **10**(2000), 463–489.
- [IU] M. Itoh and T. Umeda, *On the central elements in the universal enveloping algebra of the orthogonal Lie algebras*, Compositio Math. **127**(2001), 333–359.
- [K–] M. Kashiwara, A. Kowata, K. Minemura, K. Okamoto, T. Oshima and M. Tanaka, *Eigenfunctions of invariant differential operators on a symmetric space*, Ann. of Math. **107**(1978), 1–39.
- [Ko] B. Kostant, *Lie group representations on polynomial rings*, Amer. J. Math. **85**(1963), 327–404.
- [Ja] J. C. Jantzen, *Einhüllende Algebren halbeinfacher Lie-Algebren*, Springer-Verlag, 1983, pp. 298.
- [Jn] K. D. Johnson, *Generalized Hua operators and parabolic subgroups*, Ann. of Math. **120**(1984), 477–495.
- [Jo] A. Joseph, *Dixmier’s problem for Verma and principal series submodules*, J. London Math. Soc. **20**(1979), 193–204.
- [KW] K. Kinoshita and M. Wakayama, *Explicit Capelli identities for skew symmetric matrices*, preprint, 1998.
- [Od] H. Oda, *Annihilator operators of the degenerate principal series for simple Lie groups of type (B) and (D)*, Doctor thesis presented to the University of Tokyo, 2000.
- [OO] H. Oda and T. Oshima, *Minimal polynomials and annihilators of generalized Verma modules of the scalar type*, preprint, math.RT/0411006.
- [O1] T. Oshima, *Boundary value problems for various boundaries of symmetric spaces*, RIMS Kôkyûroku, Kyoto Univ. **281**(1976), 211–226, in Japanese.
- [O2] ———, *Capelli identities, degenerate series and hypergeometric functions*, Proceedings of a symposium on Representation Theory at Okinawa, 1995, 1–19.
- [O3] ———, *Generalized Capelli identities and boundary value problems for $GL(n)$* , Structure of Solutions of Differential Equations, World Scientific, 1996, 307–335.
- [O4] ———, *A quantization of conjugacy classes of matrices*, UTMS 2000-38, preprint, 2000, to appear in Adv. in Math.
- [OS] T. Oshima and J. Sekiguchi, *Eigenspaces of invariant differential operators on an affine symmetric spaces*, Invent. Math. **57**(1980), 1–81.
- [OSh] T. Oshima and N. Shimeno, *Boundary value problems on Riemannian symmetric spaces of the noncompact type*, in preparation.
- [Sa] H. Sakaguchi, *$U(\mathfrak{g})$ -modules associated to Grassmannian manifolds*, Master thesis presented to the University of Tokyo, 1999.
- [Se] H. Sekiguchi, *The Penrose transform for certain non-compact homogeneous manifolds of $U(n, n)$* , J. Math. Sci. Univ. Tokyo **3**(1996), 655–697.
- [Ta] T. Tanisaki, *Hypergeometric systems and Radon transforms for Hermitian symmetric spaces*, Adv. Studies in Pure Math. **26**(2000), 235–263.
- [Um] T. Umeda, *Newton’s formula for \mathfrak{gl}_n* , Proc. Amer. Math. Soc. **126** (1998), 3169–3175.