

The eleventh Takagi Lectures

November 17, 2012 (Sat) 15:10–16:10

November 18, 2012 (Sun) 11:30–12:30

Graduate School of Mathematical Sciences

The University of Tokyo

Non-Commutative Geometry and the Local Langlands Conjecture

Paul Frank Baum

(The Pennsylvania State University)

Abstract

Let G be a reductive p -adic group. Examples are $GL(n, F)$, $SL(n, F)$, etc where n can be any positive integer and F can be any finite extension of the field Q_p of p -adic numbers. The smooth dual of G is the set of (equivalence classes of) smooth representations of G . The representations are on vector spaces over the complex numbers. In a canonical way, the smooth dual of G is the disjoint union of countably many subsets known as the Bernstein components.

Results from non-commutative geometry—e.g. BC (Baum–Connes) conjecture, periodic cyclic homology of the Hecke algebra of G —indicate that a very simple geometric structure might be present in the smooth dual of G . The ABP (Aubert–Baum–Plymen) conjecture makes this precise by asserting that each Bernstein component in the smooth dual of G is a complex affine variety. These varieties are explicitly identified as certain extended quotients. For split G , (granted a mild restriction on the residual characteristic) the ABP conjecture has recently been proved for any Bernstein component in the principal series. A corollary is that the local Langlands conjecture is valid throughout the principal series. The above is joint work with Anne-Marie Aubert, Roger Plymen, and Maarten Solleveld.

Topics in these lectures:

- #1. Review of the LL (Local Langlands) conjecture.
- #2. Statement of the ABP conjecture.
- #3. Outline of the proof that for any split reductive p -adic group G both ABP and LL are valid throughout the principal series of G . Class field theory, founded by Professor Teiji Takagi, is a basic point in all three topics.