

DIFFERENTIAL SYMMETRY BREAKING OPERATORS. I. GENERAL THEORY AND F-METHOD.

TOSHIYUKI KOBAYASHI, MICHAEL PEVZNER

ABSTRACT. We prove a one-to-one correspondence between differential symmetry breaking operators for equivariant vector bundles over two homogeneous spaces and certain homomorphisms for representations of two Lie algebras, in connection with branching problems of the restriction of representations.

We develop a new method (F-method) based on the *algebraic Fourier transform for generalized Verma modules*, which characterizes differential symmetry breaking operators by means of certain systems of partial differential equations.

In contrast to the setting of real flag varieties, continuous symmetry breaking operators of Hermitian symmetric spaces are proved to be differential operators in the holomorphic setting. In this case symmetry breaking operators are characterized by differential equations of second order via the F-method.

Key words and phrases: *branching laws, F-method, symmetric pair, invariant theory, Verma modules, Hermitian symmetric spaces.*

CONTENTS

1. Introduction	1
2. Differential intertwining operators	4
3. Algebraic Fourier transform for generalized Verma modules	17
4. F-method	27
5. Localness and extension theorem for symmetry breaking operators	35
References	43

1. INTRODUCTION

Let $\mathcal{W} \rightarrow Y$ and $\mathcal{V} \rightarrow X$ be two vector bundles with a smooth map $p : Y \rightarrow X$. Then we can define “differential operators” $D : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ between the spaces of smooth sections (Definition 2.1).

Date: August 13, 2015.

2010 Mathematics Subject Classification. Primary 22E47; Secondary 22E46, 11F55, 53C10.

Suppose that $G' \subset G$ is a pair of Lie groups acting equivariantly on $\mathcal{W} \rightarrow Y$ and $\mathcal{V} \rightarrow X$, respectively, and that p is G' -equivariant. The object of the present work is the study of G' -intertwining differential operators (*differential symmetry breaking operators*). If \mathcal{W} is isomorphic to the pull-back $p^*\mathcal{V}$, then the restriction map $f \mapsto f|_Y$ is obviously a G' -intertwining operator (and a differential operator of order zero). In the general setting where there is no morphism from $p^*\mathcal{V}$ to \mathcal{W} , non-zero G' -intertwining differential operators may or may not exist.

Suppose that G acts transitively on X and G' acts transitively on Y . We write X (respectively Y) as a homogeneous space $= G/H$ (respectively $Y = G'/H'$). The first main result is a duality theorem that gives a one-to-one correspondence between G' -intertwining differential operators and (\mathfrak{g}', H') -homomorphisms for induced representations of Lie algebras (see Corollary 2.10 for the precise notation):

Theorem A. *Suppose $H' \subset H$. Then there is a natural bijection:*

$$(1.1) \quad D_{X \rightarrow Y} : \mathrm{Hom}_{(\mathfrak{g}', H')}(\mathrm{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \mathrm{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} \mathrm{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y).$$

This generalizes a well-known result in the case where G and G' are the same reductive group and where X and Y are the same flag variety ([Kos74, HJ82]).

By a *branching problem* we mean a problem of understanding how a given representation of a group G behaves when restricted to a subgroup G' . For a unitary representation π of G , branching problems concern a decomposition of π into the direct integral of irreducible unitary representations of G' (branching law).

More generally, for non-unitary representations π and τ of G and G' , respectively, we may consider the space $\mathrm{Hom}_{G'}(\pi|_{G'}, \tau)$ of continuous G' -homomorphisms. The right-hand side of (1.1) concerns branching problems with respect to the restriction from G to G' , whereas the left-hand side of (1.1) concerns branching laws of “generalized Verma modules”.

If $\mathrm{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$ in (1.1) is one-dimensional, we may regard its generator as canonical up to a scalar and be tempted to find an explicit description for such a natural differential symmetry breaking operator. It should be noted that seeking explicit formulæ of intertwining operators is much more involved than finding abstract branching laws, as we may observe with the celebrated Rankin–Cohen brackets which appear as symmetry breaking operators in the decomposition of the tensor product of two holomorphic discrete series representations of $SL(2, \mathbb{R})$ (see [DP07, KP15-2] for a detailed discussion).

The condition $\dim \mathrm{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) \leq 1$ is often fulfilled when \mathfrak{h} is a parabolic subalgebra of \mathfrak{g} with abelian nilradical, (see [K14, Theorem 2.7]). Moreover, finding all bundles \mathcal{W}_Y for which such nontrivial intertwining operators exist is a part of the

initial problem, which reduces to abstract branching problems (see [KP15-2, Fact 4.3]).

We propose a new method to find explicit expressions for differential symmetry breaking operators appearing in this geometric setting. We call it the *F-method*, where F stands for the Fourier transform. More precisely, we consider an “algebraic Fourier transform” of generalized Verma modules, and characterize symmetry breaking operators by means of certain systems of partial differential equations. If \mathfrak{h} is a parabolic subalgebra with abelian nilradical, then the system is of second order although the resulting differential symmetry breaking operators may be of any higher order. The characterization is performed by applying an algebraic Fourier transform (see Definition 3.1). A detailed recipe of the *F-method* is described in Section 4.4 relying on Theorem 4.1 and Proposition 3.11.

In general, symmetry breaking operators between two principal series representations of real reductive Lie groups $G' \subset G$ are given by integro-differential operators in geometric models. Among them, equivariant differential operators are very special (e.g. [KnSt71] for $G' = G$ and [KS14] for $G' \subsetneq G$). However, in the case where X is a Hermitian symmetric space, Y a subsymmetric space, $G' \subset G$ are the groups of biholomorphic transformations of $Y \rightarrow X$, respectively, we prove the following *localness and extension theorem*:

Theorem B. *Any continuous G' -homomorphism from $\mathcal{O}(X, \mathcal{V})$ to $\mathcal{O}(Y, \mathcal{W})$ is given by a holomorphic differential operator, which extends to the whole flag variety.*

See Theorem 5.3 for the precise statement. Theorem B includes the case of the tensor product of two holomorphic discrete series representations corresponding to the setting where $G \simeq G' \times G'$ and $X \simeq Y \times Y$ as a special case.

In the second part of the work [KP15-2] we apply the F-method to Hermitian symmetric spaces to find explicit formulæ of differential symmetry breaking operators in the six parabolic geometries arising from symmetric pairs of split rank one.

The authors are grateful to the referee for enlightening remarks and for suggesting to divide the original manuscript (old title “*Rankin-Cohen operators for symmetric pairs*”) into two parts and to write more detailed proofs and explanations in the first part, not only for specialists but also for a broader audience. We would like to extend a special thanks to Dr. T. Kubo for providing valuable and constructive suggestions in respect to its legibility.

Notation: $\mathbb{N} = \{0, 1, 2, \dots\}$.

2. DIFFERENTIAL INTERTWINING OPERATORS

In this section we discuss equivariant differential operators between sections of homogeneous vector bundles in a more general setting than the usual. Namely, we consider vector bundles admitting a morphism between their base spaces. In this generality, we establish a natural bijection between such differential operators (*differential symmetry breaking operators*) and certain homomorphisms arising from the branching problems for infinite-dimensional representations of Lie algebras, see Theorem 2.9 (*duality theorem*).

2.1. Differential operators between two manifolds. We understand the notion of differential operators between two vector bundles in the usual sense when the bundles are defined over the same base space. We extend this terminology in a more general setting, where there exists a morphism between base spaces. Let $\mathcal{V} \rightarrow X$ be a vector bundle over a smooth manifold X . We write $C^\infty(X, \mathcal{V})$ for the space of smooth sections, which is endowed with the Fréchet topology of uniform convergence of sections and their derivatives of finite order on compact sets. Let $\mathcal{W} \rightarrow Y$ be another vector bundle, and $p: Y \rightarrow X$ a smooth map between the base spaces.

Definition 2.1. We say that a continuous linear map $T: C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ is a *differential operator* if T satisfies

$$(2.1) \quad p(\text{Supp } Tf) \subset \text{Supp } f \quad \text{for any } f \in C^\infty(X, \mathcal{V}).$$

We write $\text{Diff}(\mathcal{V}_X, \mathcal{W}_Y)$ for the vector space of differential operators from $C^\infty(X, \mathcal{V})$ to $C^\infty(Y, \mathcal{W})$.

The condition (2.1) shows that T is a local operator in the sense that for any open subset U of X , T induces a continuous linear map:

$$T_U: C^\infty(U, \mathcal{V}|_U) \longrightarrow C^\infty(p^{-1}(U), \mathcal{W}|_{p^{-1}(U)}).$$

Remark 2.2. If $X = Y$ and p is the identity map, then the condition (2.1) is equivalent to T being a differential operator in the usual sense owing to Peetre's celebrated theorem [Pee59]. Our proof of Lemma 2.3 in this special case gives an account of this classical theorem by using the theory of distributions due to L. Schwartz [S66].

Let $\Omega_X := |\wedge^{\text{top}} T^*(X)|$ be the bundle of densities. For a vector bundle $\mathcal{V} \rightarrow X$, we set $\mathcal{V}^\vee := \coprod_{x \in X} \mathcal{V}_x^\vee$ where $\mathcal{V}_x^\vee := \text{Hom}_{\mathbb{C}}(\mathcal{V}_x, \mathbb{C})$, and denote by \mathcal{V}^* the dualizing bundle $\mathcal{V}^\vee \otimes \Omega_X$. In what follows $\mathcal{D}'(X, \mathcal{V}^*)$ (respectively, $\mathcal{E}'(X, \mathcal{V}^*)$) denotes the space of \mathcal{V}^* -valued distributions (respectively, those with compact support). We shall regard distributions as generalized functions à la Gelfand rather than continuous linear forms on $C_c^\infty(X)$ or $C^\infty(X)$. In particular, we sometimes write as

$$(2.2) \quad \mathcal{E}'(X, \Omega_X) \rightarrow \mathbb{C}, \quad \omega \mapsto \int_X \omega,$$

to denote the natural pairing $\langle \omega, \mathbf{1}_X \rangle$ of ω with the constant function $\mathbf{1}_X$ on X . Composing (2.2) with the contraction on the fiber, we get a natural bilinear map

$$(2.3) \quad C^\infty(X, \mathcal{V}) \times \mathcal{E}'(X, \mathcal{V}^*) \rightarrow \mathbb{C}, \quad (f, \omega) \mapsto \langle f, \omega \rangle = \int_X f \omega.$$

Let $\mathcal{V}^* \boxtimes \mathcal{W}$ denote the tensor product bundle over $X \times Y$ of the two vector bundles $\mathcal{V}^* \rightarrow X$ and $\mathcal{W} \rightarrow Y$. Then for any continuous linear map $T : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ there exists a unique distribution $K_T \in \mathcal{D}'(X \times Y, \mathcal{V}^* \boxtimes \mathcal{W})$ such that the projection on the second factor $\text{pr}_2 : X \times Y \rightarrow Y$ is proper on the support of K_T and such that

$$(Tf)(y) = \langle K_T(\cdot, y), f(\cdot) \rangle \text{ for any } f \in C^\infty(X, \mathcal{V}),$$

by the Schwartz kernel theorem.

Given a map $p : Y \rightarrow X$, we set

$$\Delta(Y) := \{(p(y), y) : y \in Y\} \subset X \times Y.$$

The following lemma characterizes differential operators by means of the distribution kernels K_T .

Lemma 2.3. *Let $p : Y \rightarrow X$ be a smooth map. A continuous operator $T : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ is a differential operator in the sense of Definition 2.1 if and only if $\text{Supp } K_T \subset \Delta(Y)$.*

Proof. Suppose $\text{Supp } K_T \subset \Delta(Y)$. Let $(x_o, y_o) \in \Delta(Y)$ and take a neighborhood U of $x_o = p(y_o)$ in X and a neighborhood U' of y_o in Y such that $U' \subset p^{-1}(U)$. We trivialize the bundles locally as $\mathcal{V}|_U \simeq U \times V$ and $\mathcal{W}|_{U'} \simeq U' \times W$. Let (x_1, \dots, x_m) be the coordinates in U . According to the structural theory of distributions supported on a submanifold $\Delta Y \subset X \times Y$ [S66, Chapter III, Théorème XXXVII], there exists a unique family $h_\alpha(y) \in \mathcal{D}'(U') \otimes W$ for a finite number of multi-indices $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{N}^m$, such that $\langle K_T, f \rangle \in \mathcal{D}'(U') \otimes \text{Hom}_{\mathbb{C}}(V, W)$ is locally given as a finite sum

$$(2.4) \quad \sum_{\alpha} h_{\alpha}(y) \frac{\partial^{|\alpha|} f}{\partial x^{\alpha}}(p(y)),$$

for every $f \in C^\infty(X, \mathcal{V})$. Hence $\langle K_T, f \rangle|_{U'} = 0$ if $f|_U = 0$. Thus T is a differential operator in the sense of Definition 2.1.

Conversely, take any $(x_o, y_o) \in \text{Supp } K_T$. By the definition of the distribution kernel K_T , for any neighborhood S of x_o in X there exists $f \in C^\infty(X, \mathcal{V})$ such that $\text{Supp } f \subset S$ and $(x_o, y_o) \in \text{Supp } f \times \text{Supp } Tf$. If T is a differential operator then by (2.1)

$$p(\text{Supp } Tf) \subset \text{Supp } f \subset S.$$

Since S is an arbitrary neighborhood of x_o , $p(y_o)$ must coincide with x_o . Hence $\text{Supp } K_T \subset \Delta(Y)$. \square

By (2.4), the terminology “differential operators” in Definition 2.1 is justified as follows:

Example 2.4. (1) *Let $p : Y \twoheadrightarrow X$ be a submersion. Choose an atlas of local coordinates $\{(x_i, z_j)\}$ on Y in such a way that $\{x_i\}$ form an atlas on X . Then, every $T \in \text{Diff}(\mathcal{V}_X, \mathcal{W}_Y)$ is locally of the form*

$$\sum_{\alpha \in \mathbb{N}^{\dim X}} h_\alpha(x, z) \frac{\partial^{|\alpha|}}{\partial x^\alpha} \quad (\text{finite sum}),$$

where $h_\alpha(x, z)$ are $\text{Hom}(V, W)$ -valued smooth functions on Y .

(2) *Let $i : Y \hookrightarrow X$ be an immersion. Choose an atlas of local coordinates $\{(y_i, z_j)\}$ on X in such a way that $\{y_i\}$ form an atlas on Y . Then, every $T \in \text{Diff}(\mathcal{V}_X, \mathcal{W}_Y)$ is locally of the form*

$$\sum_{(\alpha, \beta) \in \mathbb{N}^{\dim X}} g_{\alpha, \beta}(y) \frac{\partial^{|\alpha|+|\beta|}}{\partial y^\alpha \partial z^\beta} \quad (\text{finite sum}),$$

where $g_{\alpha, \beta}(y)$ are $\text{Hom}(V, W)$ -valued smooth functions on Y .

Next, suppose that the two vector bundles $\mathcal{V} \rightarrow X$ and $\mathcal{W} \rightarrow Y$ are equivariant with respect to a given Lie group G . Then we have natural actions of G on the Fréchet spaces $C^\infty(X, \mathcal{V})$ and $C^\infty(Y, \mathcal{W})$ by translations. Denote by $\text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(Y, \mathcal{W}))$ the space of continuous G -homomorphisms. We set

$$(2.5) \quad \text{Diff}_G(\mathcal{V}_X, \mathcal{W}_Y) := \text{Diff}(\mathcal{V}_X, \mathcal{W}_Y) \cap \text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(Y, \mathcal{W})).$$

Example 2.5. *Suppose X and Y are both Euclidean vector spaces with an injective linear map $p : Y \hookrightarrow X$. If G contains the subgroup of all translations of Y then $\text{Diff}_G(\mathcal{V}_X, \mathcal{W}_Y)$ is a subspace of the space of differential operators with constant coefficients.*

An analogous notion can be defined in the holomorphic setting. Let $\mathcal{V} \rightarrow X$ and $\mathcal{W} \rightarrow Y$ be two holomorphic vector bundles with a holomorphic map $p : Y \rightarrow X$ between the complex base manifolds X and Y . We say a differential operator $T : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ is *holomorphic* if

$$T_U(\mathcal{O}(U, \mathcal{V}|_U)) \subset \mathcal{O}(p^{-1}(U), \mathcal{W}|_{p^{-1}(U)})$$

for any open subset U of X . We denote by $\text{Diff}^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y)$ the vector space of holomorphic differential operators. When a Lie group G acts biholomorphically on the two holomorphic vector bundles $\mathcal{V} \rightarrow X$ and $\mathcal{W} \rightarrow Y$, we set

$$\text{Diff}_G^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y) := \text{Diff}^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y) \cap \text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(Y, \mathcal{W})).$$

2.2. Induced modules. Let \mathfrak{g} be a Lie algebra over \mathbb{C} , and $U(\mathfrak{g})$ its universal enveloping algebra. Let \mathfrak{h} be a Lie subalgebra of \mathfrak{g} .

Definition 2.6. For an \mathfrak{h} -module V , we define the induced $U(\mathfrak{g})$ -module $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V)$ as

$$\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V) := U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} V.$$

If \mathfrak{h} is a Borel subalgebra and $\dim V = 1$, then the \mathfrak{g} -module $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V)$ is the Verma module.

For later purposes we formulate the following statement in terms of the contragredient representation V^\vee . Let \mathfrak{h}' be another Lie subalgebra of \mathfrak{g} .

Proposition 2.7. *For a finite-dimensional \mathfrak{h}' -module W we have:*

- (1) $\text{Hom}_{\mathfrak{g}}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \simeq \text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$.
- (2) *If $\mathfrak{h}' \not\subseteq \mathfrak{h}$, then $\text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) = \{0\}$.*

Proof. The first statement is due to the functoriality of the tensor product.

For the second statement it suffices to treat the case where \mathfrak{h}' is one-dimensional. Then the assumption $\mathfrak{h}' \not\subseteq \mathfrak{h}$ implies that $\mathfrak{h}' \cap \mathfrak{h} = \{0\}$, and therefore there is a direct sum decomposition of vector spaces:

$$\mathfrak{g} = \mathfrak{h}' + \mathfrak{q} + \mathfrak{h},$$

for some subspace \mathfrak{q} in \mathfrak{g} . We fix a basis X_1, \dots, X_n of \mathfrak{q} , and define a subspace of $U(\mathfrak{g})$ by

$$U'(\mathfrak{q}) := \mathbb{C}\text{-span} \{X_1^{\alpha_1} \cdots X_n^{\alpha_n} : (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n\}.$$

Then, by the Poincaré–Birkhoff–Witt theorem we have an isomorphism of \mathfrak{h}' -modules:

$$\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \simeq U(\mathfrak{h}') \otimes_{\mathbb{C}} U'(\mathfrak{q}) \otimes_{\mathbb{C}} V^\vee.$$

In particular, $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ is a free $U(\mathfrak{h}')$ -module. Hence there does not exist a non-zero *finite-dimensional* \mathfrak{h}' -submodule in the \mathfrak{g} -module $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$. \square

Remark 2.8. We shall see in Theorem 2.9 that $\dim_{\mathbb{C}} \text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$ is equal to the dimension of the space of differential symmetry breaking operators from $C^\infty(X, \mathcal{V})$ to $C^\infty(Y, \mathcal{W})$ when H' is connected. In [KP15-2, Section 2], we give a family of sextuples $(\mathfrak{g}, \mathfrak{g}', \mathfrak{h}, \mathfrak{h}', V, W)$ such that this dimension is one.

2.3. Duality theorem for differential operators between two homogeneous spaces. Let G be a real Lie group, and $\mathfrak{g}(\mathbb{R}) := \text{Lie}(G)$. We denote by $U(\mathfrak{g})$ the universal enveloping algebra of the complexified Lie algebra $\mathfrak{g} := \mathfrak{g}(\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$. Analogous notations will be applied to other Lie groups.

Let H be a closed subgroup of G . Given a finite-dimensional representation $\lambda : H \rightarrow GL_{\mathbb{C}}(V)$ we define the homogeneous vector bundle $\mathcal{V}_X \equiv \mathcal{V} := G \times_H V$ over

$X := G/H$. As a G -module, the space $C^\infty(X, \mathcal{V})$ of smooth sections is identified with the following subspace of $C^\infty(G, V) \simeq C^\infty(G) \otimes V$:

$$\begin{aligned} C^\infty(G, V)^H &:= \{f \in C^\infty(G, V) : f(gh) = \lambda(h)^{-1}f(g) \text{ for any } g \in G, h \in H\} \\ &\simeq \{F \in C^\infty(G) \otimes V : \lambda(h)F(gh) = F(g) \text{ for any } g \in G, h \in H\}. \end{aligned}$$

In dealing with a representation V of a disconnected subgroup H (e.g. H is a parabolic subgroup of a real reductive Lie group G), we notice that the diagonal H -action on $U(\mathfrak{g}) \otimes_{\mathbb{C}} V^\vee$ defines a representation of H on $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) = U(\mathfrak{g}) \otimes_{\mathfrak{h}} V^\vee$ and thus $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ is endowed with a (\mathfrak{g}, H) -module structure.

Theorem 2.9 (Duality theorem). *Let $H' \subset H$ be (possibly disconnected) closed subgroups of a Lie group G with Lie algebras $\mathfrak{h}' \subset \mathfrak{h}$, respectively. Suppose V and W are finite-dimensional representations of H and H' , respectively. Let G' be any subgroup of G containing H' , and $\mathcal{V}_X := G \times_H V$ and $\mathcal{W}_Y := G' \times_{H'} W$ be the corresponding homogeneous vector bundles. Then, there is a natural linear isomorphism:*

$$(2.6) \quad D_{X \rightarrow Y} : \text{Hom}_{H'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y),$$

or equivalently,

$$D_{X \rightarrow Y} : \text{Hom}_{(\mathfrak{g}', H')}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y).$$

For $\varphi \in \text{Hom}_{H'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$ and $F \in C^\infty(X, \mathcal{V}) \simeq C^\infty(G, V)^H$, $D_{X \rightarrow Y}(\varphi)F \in C^\infty(Y, \mathcal{W}) \simeq C^\infty(G', W)^{H'}$ is given by the following formula:

$$(2.7) \quad \langle D_{X \rightarrow Y}(\varphi)F, w^\vee \rangle = \sum_j \langle dR(u_j)F, v_j^\vee \rangle|_{G'} \quad \text{for } w^\vee \in W^\vee,$$

where $\varphi(w^\vee) = \sum_j u_j v_j^\vee \in \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ ($u_j \in U(\mathfrak{g})$, $v_j^\vee \in V^\vee$).

When H' is connected, we can write the left-hand side of (2.6) by means of Lie algebras.

Corollary 2.10. *Suppose we are in the setting of Theorem 2.9. Assume that H' is connected. Then there is a natural linear isomorphism:*

$$(2.8) \quad D_{X \rightarrow Y} : \text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y),$$

or equivalently,

$$(2.8)' \quad D_{X \rightarrow Y} : \text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y).$$

The construction of $D_{X \rightarrow Y}$ and the fact that the formula (2.7) is well-defined will be explained in Section 2.4.

Remark 2.11. (1) Corollary 2.10 is known when $X = Y$, i.e. $G' = G$ and $H' = H$, especially in the setting of complex flag varieties, see e.g. [Kos74, HJ82].

- (2) When \mathfrak{g}' is a reductive subalgebra and \mathfrak{h}' is a parabolic subalgebra, the existence of an \mathfrak{h}' -module W for which the left-hand side of (2.8) is non-zero, is closely related to the “discretely decomposability” of the \mathfrak{g} -module $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ when restricted to the subalgebra \mathfrak{g}' ([K98b], [K12]). This relationship will be used in Section 5 in proving that any continuous symmetry breaking operator in a holomorphic setting is given by a differential operator (*localness theorem*).
- (3) Owing to Proposition 2.7, the left-hand side of (2.8)' is non-zero only when $\mathfrak{h}' \subset \mathfrak{h}$. Conversely, if $H' \subset H \cap G'$, then there is a natural morphism $Y = G'/H' \rightarrow X = G/H$ and therefore “differential operators” (in the sense of Definition 2.1) from $C^\infty(X, \mathcal{V})$ to $C^\infty(Y, \mathcal{W})$ are defined.
- (4) We shall consider the case where $H' = H \cap G'$ in later applications, however, Theorem 2.9 also covers the cases where the natural morphism $Y \rightarrow X$ is not injective, *i.e.* where $H' \not\subset H \cap G'$.

An analogous result to Theorem 2.9 holds in the holomorphic setting as well. To be precise, let $G_{\mathbb{C}}$ be a complex Lie group, $G'_{\mathbb{C}}$, $H_{\mathbb{C}}$ and $H'_{\mathbb{C}}$ be closed complex subgroups such that $H'_{\mathbb{C}} \subset H_{\mathbb{C}} \cap G'_{\mathbb{C}}$. We write \mathfrak{g} , \mathfrak{h} , \dots for the Lie algebras of the complex Lie groups $G_{\mathbb{C}}$, $H_{\mathbb{C}}$, \dots , respectively. Given finite-dimensional holomorphic representations V of $H_{\mathbb{C}}$ and W of $H'_{\mathbb{C}}$, we form holomorphic vector bundles $\mathcal{V} := G_{\mathbb{C}} \times_{H_{\mathbb{C}}} V$ over $X_{\mathbb{C}} = G_{\mathbb{C}}/H_{\mathbb{C}}$ and $\mathcal{W} := G'_{\mathbb{C}} \times_{H'_{\mathbb{C}}} W$ over $Y_{\mathbb{C}} = G'_{\mathbb{C}}/H'_{\mathbb{C}}$.

For simplicity, we assume that $H'_{\mathbb{C}}$ is connected. (This is always the case if $G'_{\mathbb{C}}$ is a connected complex reductive Lie group and $H'_{\mathbb{C}}$ is a parabolic subgroup of $G'_{\mathbb{C}}$.) Then we have:

Theorem 2.12 (Duality theorem in the holomorphic setting). *There is a canonical linear isomorphism:*

$$D_{X \rightarrow Y} : \text{Hom}_{\mathfrak{g}'} \left(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \right) \xrightarrow{\sim} \text{Diff}_{G'_{\mathbb{C}}}^{\text{hol}}(\mathcal{V}_{X_{\mathbb{C}}}, \mathcal{W}_{Y_{\mathbb{C}}}).$$

Suppose furthermore that G , G' , H and H' are real forms of the complex Lie groups $G_{\mathbb{C}}$, $G'_{\mathbb{C}}$, $H_{\mathbb{C}}$ and $H'_{\mathbb{C}}$, respectively. We regard V and W as H - and H' -modules by the restriction, and form vector bundles $\mathcal{V} = G \times_H V$ over $X = G/H$ and $\mathcal{W} = G' \times_{H'} W$ over $Y = G'/H'$.

We ask whether or not all symmetry breaking operators have holomorphic extensions. Here is a simple sufficient condition:

Corollary 2.13. *If H' is contained in the connected complexification $H'_{\mathbb{C}}$, then we have a natural bijection:*

$$\text{Diff}_{G'_{\mathbb{C}}}^{\text{hol}}(\mathcal{V}_{X_{\mathbb{C}}}, \mathcal{W}_{Y_{\mathbb{C}}}) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y).$$

Proof. Comparing Theorem 2.9 with Theorem 2.12, the proof of Corollary 2.13 reduces to the surjectivity of the inclusion

$$(2.9) \quad \text{Hom}_{(\mathfrak{g}', H')}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \hookrightarrow \text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)).$$

We note that $\text{Hom}_{(\mathfrak{g}', H'_\mathbb{C})}(\text{ind}_{\mathfrak{h}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$ is a subspace of the left-hand side of (2.9) because $H' \subset H'_\mathbb{C}$, whereas it coincides with the right-hand side of (2.9) if $H'_\mathbb{C}$ is connected. Hence (2.9) is surjective. Thus Corollary 2.13 is proved. \square

The rest of this section is devoted to the proof of Theorem 2.9. For Theorem 2.12, since the argument is parallel to that of Theorem 2.9, we omit the proof.

2.4. Construction of $D_{X \rightarrow Y}$. This subsection gives the definition of the linear map $D_{X \rightarrow Y}$ in Theorem 2.9.

Consider two actions dR and dL of the universal enveloping algebra $U(\mathfrak{g})$ on the space $C^\infty(G)$ of smooth complex-valued functions on G induced by the regular representation $L \times R$ of $G \times G$ on $C^\infty(G)$:

$$(2.10) \quad (dR(Z)f)(x) := \left. \frac{d}{dt} \right|_{t=0} f(xe^{tZ}) \quad \text{and} \quad (dL(Z)f)(x) := \left. \frac{d}{dt} \right|_{t=0} f(e^{-tZ}x),$$

for $Z \in \mathfrak{g}(\mathbb{R})$.

The right differentiation (2.10) defines a bilinear map

$$\Phi : C^\infty(G) \times U(\mathfrak{g}) \rightarrow C^\infty(G), \quad (F, u) \mapsto dR(u)F,$$

with the following properties

$$(2.11) \quad \Phi(L(g)F, u) = L(g)\Phi(F, u),$$

$$(2.12) \quad \Phi(F, u'u) = dR(u')\Phi(F, u),$$

for any $g \in G$ and $u, u' \in U(\mathfrak{g})$.

Combining Φ with the canonical pairing $V \times V^\vee \rightarrow \mathbb{C}$, we obtain a bilinear map

$$\Phi_V : C^\infty(G) \otimes V \times U(\mathfrak{g}) \otimes_{\mathbb{C}} V^\vee \rightarrow C^\infty(G).$$

Then we have the following:

Lemma 2.14. *The map Φ_V induces a well-defined diagram of maps:*

$$\begin{array}{ccc} C^\infty(G) \otimes V \times U(\mathfrak{g}) \otimes_{\mathbb{C}} V^\vee & \xrightarrow{\Phi_V} & C^\infty(G) \\ \uparrow & & \parallel \\ C^\infty(X, \mathcal{V}) \times \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) & \dashrightarrow & C^\infty(G). \end{array}$$

Proof. Denote by λ^\vee the contragredient representation of the representation (λ, V) of H , and by $d\lambda^\vee$ the infinitesimal representation of \mathfrak{h} . The kernel of the natural quotient map $U(\mathfrak{g}) \otimes_{\mathbb{C}} V^\vee \rightarrow \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ is generated by

$$-uY \otimes v^\vee + u \otimes d\lambda^\vee(Y)v^\vee$$

with $u \in U(\mathfrak{g})$, $Y \in \mathfrak{h}$ and $v^\vee \in V^\vee$. Hence it suffices to show

$$\Phi_V(f, -uY \otimes v^\vee + u \otimes d\lambda^\vee(Y)v^\vee) = 0$$

for any $f \in C^\infty(X, \mathcal{V}) \simeq C^\infty(G, V)^H$.

Since $f \in C^\infty(G, V)^H$ satisfies $dR(Y)f = -d\lambda(Y)f$ for $Y \in \mathfrak{h}$, we have

$$\begin{aligned} \Phi_V(f, uY \otimes v^\vee) &= \langle dR(u)dR(Y)f, v^\vee \rangle \\ &= \langle dR(u)f, d\lambda^\vee(Y)v^\vee \rangle \\ &= \Phi_V(f, u \otimes d\lambda^\vee(Y)v^\vee). \end{aligned}$$

Thus the lemma is proved. \square

Lemma 2.15. 1) *The bilinear map*

$$(2.13) \quad C^\infty(X, \mathcal{V}) \times \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \rightarrow \mathbb{C}, (f, m) \mapsto \Phi_V(f, m)(e)$$

is (\mathfrak{g}, H) -invariant.

2) *If $m \in \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ satisfies $\Phi_V(f, m)(e) = 0$ for all $f \in C^\infty(X, \mathcal{V})$ then $m = 0$.*

Proof. 1) Let $f \in C^\infty(X, \mathcal{V})$ and $m \in \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$. It follows from (2.11) and (2.12) that

$$\begin{aligned} \Phi_V(dL(Z)f, m) &= dL(Z)\Phi_V(f, m) \\ \Phi_V(f, Zm) &= dR(Z)\Phi_V(f, m) \end{aligned}$$

for any $Z \in \mathfrak{g}$. Since

$$(dL(Z) + dR(Z))F(e) = 0$$

for any $F \in C^\infty(G)$, we have shown the \mathfrak{g} -invariance of the bilinear map (2.13):

$$\Phi_V(dL(Z)f, m)(e) + \Phi_V(f, Zm)(e) = 0.$$

The proof for the H -invariance of (2.13) is similar.

2) Take a basis $\{v_1, \dots, v_k\}$ of V , and let $\{v_1^\vee, \dots, v_k^\vee\}$ be the dual basis in V^\vee . Choose a complementary subspace \mathfrak{q} of \mathfrak{h} in \mathfrak{g} , and fix a basis $\{X_1, \dots, X_n\}$ of \mathfrak{q} . Then by the Poincaré–Birkhoff–Witt theorem, we can write $m \in \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ as a finite sum:

$$m = \sum_{j=1}^k \sum_{\alpha=(\alpha_1, \dots, \alpha_n)} a_{\alpha, j} X_1^{\alpha_1} \dots X_n^{\alpha_n} v_j^\vee.$$

If m is non-zero, we can find a multi-index β and j_o ($1 \leq j_o \leq k$) such that $a_{\beta, j_o} \neq 0$ and that $a_{\alpha, j_o} = 0$ for any multi-index α satisfying $|\alpha| > |\beta|$ and for any j . Here $|\alpha| = \sum_{i=1}^n \alpha_i$ for $\alpha \in \mathbb{N}^n$. We take $f \in C^\infty(G, V)^H \simeq C^\infty(X, \mathcal{V})$ such that f is given in a right H -invariant neighborhood of H in G by

$$f\left(\exp\left(\sum_{i=1}^n x_i X_i\right)h\right) = x^\beta \lambda(h)^{-1} v_{j_o} \quad \text{for } x = (x_1, \dots, x_n) \in \mathbb{R}^n \text{ and } h \in H.$$

Then $\Phi_V(f, m)(e) = a_{\beta, j_o} \beta_1! \dots \beta_k! \neq 0$. The contraposition completes the proof. \square

We regard $C^\infty(G)$ as a $G \times \mathfrak{g}$ -module via the $(L \times dR)$ -action. Then the space $\text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(G))$ of continuous G -homomorphisms becomes a \mathfrak{g} -module by the remaining dR -action on the target space. By (2.11), (2.12) and Lemma 2.14, we get the following \mathfrak{g} -homomorphism:

$$(2.14) \quad \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \longrightarrow \text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(G)), \quad u \otimes v^\vee \mapsto (f \mapsto \langle dR(u)f, v^\vee \rangle).$$

Furthermore, it is actually a (\mathfrak{g}, H) -homomorphism, where the group H acts on $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) = U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} V^\vee$ diagonally and acts on $\text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(G))$ via the R -action on $C^\infty(G)$.

Let H' be a connected closed Lie subgroup of G . Given a finite-dimensional representation W of H' , we form a homogeneous vector bundle $\mathcal{W}_Z \equiv \mathcal{W} := G \times_{H'} W$ over $Z := G/H'$.

Taking the tensor product of the (\mathfrak{g}, H) -modules in (2.14) with the H' -module W , we get an $(H' \times (\mathfrak{g}, H))$ -homomorphism:

$$\text{Hom}_{\mathbb{C}}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \longrightarrow \text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(G, W)).$$

Let $\Delta(H')$ be a subgroup of $H' \times H$ defined by $\{(h, h) : h \in H'\}$. Taking $\Delta(H')$ -invariants, we obtain the following \mathbb{C} -linear map:

$$(2.15) \quad \text{Hom}_{H'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \longrightarrow \text{Hom}_G(C^\infty(X, \mathcal{V}), C^\infty(Z, \mathcal{W})), \quad \varphi \mapsto D_\varphi,$$

where D_φ satisfies

$$(2.16) \quad \langle D_\varphi f, w^\vee \rangle = \Phi_V(f, \varphi(w^\vee))$$

for any $f \in C^\infty(X, \mathcal{V})$ and any $w^\vee \in W^\vee$.

Remark 2.16. If H' is connected, then we can replace $\text{Hom}_{H'}$ by $\text{Hom}_{\mathfrak{h}'}$ in (2.15).

Lemma 2.17. *The map (2.15) is injective.*

Proof. By (2.16), Lemma 2.17 is derived from the second statement of the Lemma 2.15. \square

Take any subgroup G' of G containing H' and form a homogeneous vector bundle $\mathcal{W}_Y := G' \times_{H'} W$ over $Y = G'/H'$. Then, the vector bundle \mathcal{W}_Y is isomorphic to the restriction $\mathcal{W}_Z|_Y$ of the vector bundle \mathcal{W}_Z to the submanifold Y of the base space Z . Let

$$R_{Z \rightarrow Y} : C^\infty(Z, \mathcal{W}_Z) \rightarrow C^\infty(Y, \mathcal{W}_Y)$$

be the restriction map of sections. For $\varphi \in \text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$ we set

$$(2.17) \quad D_{X \rightarrow Y}(\varphi) := R_{Z \rightarrow Y} \circ D_\varphi.$$

Then $D_{X \rightarrow Y}(\varphi) : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ is a G' -equivariant differential operator, *i.e.* $D_{X \rightarrow Y}$ defines a linear map $\text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \rightarrow \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$. Theorem 2.9

describes explicitly the image $D_{X \rightarrow Y}$ when $H' \subset H \cap G'$, namely, when the following diagram exists:

$$\begin{array}{ccc} & Z = G/H' & \\ \swarrow & & \searrow \\ Y = G'/H' & \xrightarrow{\quad} & X = G/H \end{array}$$

Remark 2.18. The left-hand side of (2.8) does not depend on the choice of G' . This fact is reflected by the commutativity of the following diagram.

$$(2.18) \quad \begin{array}{ccc} \text{Hom}_{\mathfrak{h}'}(W^\vee, \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) & \xrightarrow{\sim} & \text{Diff}_G(\mathcal{V}_X, \mathcal{W}_Z) \\ & \searrow \sim & \swarrow \\ & \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) & \end{array}$$

$D_{X \rightarrow Y}$ $R_{Z \rightarrow Y}$

2.5. Proof of Theorem 2.9. We have already seen in Lemma 2.17 that $D_{X \rightarrow Y}$ is injective. In order to prove the surjectivity of the linear map $D_{X \rightarrow Y}$, we realize the induced $U(\mathfrak{g})$ -module $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ in the space of distributions.

We recall that $\mathcal{V}^* = \mathcal{V}^\vee \otimes \Omega_X$ is the dualizing bundle of a vector bundle \mathcal{V} over X . For a closed subset S and an open subset U in X containing S , we write $\mathcal{D}'_S(U, \mathcal{V}^*)$ for the space of \mathcal{V}^* -valued distributions on U with support in S . Obviously, $\mathcal{D}'_S(U, \mathcal{V}^*) = \mathcal{D}'_S(X, \mathcal{V}^*)$. If S is compact, then $\mathcal{D}'_S(U, \mathcal{V}^*)$ is contained in the space $\mathcal{E}'(U, \mathcal{V}^*)$ of distributions on U with compact support, and thus coincides with $\mathcal{E}'_S(U, \mathcal{V}^*) := \mathcal{D}'_S(U, \mathcal{V}^*) \cap \mathcal{E}'(U, \mathcal{V}^*)$.

We return to the setting of Theorem 2.9, where \mathcal{V} is a G -equivariant vector bundle over $X = G/H$. Then the Lie group G acts on $C^\infty(X, \mathcal{V})$ and $\mathcal{E}'(X, \mathcal{V}^*)$ by the pull-back of smooth sections and distributions, respectively. The infinitesimal action defines representations of the Lie algebra \mathfrak{g} on $C^\infty(U, \mathcal{V})$ and $\mathcal{E}'_S(U, \mathcal{V}^*)$.

The “integration map” (2.2)

$$(2.19) \quad \mathcal{E}'(X, \Omega_X) \rightarrow \mathbb{C}, \quad \omega \mapsto \int_X \omega$$

is G -invariant. Composing this with the G -invariant bilinear map (*contraction*):

$$C^\infty(X, \mathcal{V}) \times \mathcal{E}'(X, \mathcal{V}^*) \longrightarrow \mathcal{E}'(X, \Omega_X), \quad (f, h) \mapsto \langle f, h \rangle,$$

we obtain the following G -invariant bilinear form

$$(2.20) \quad C^\infty(X, \mathcal{V}) \times \mathcal{E}'(X, \mathcal{V}^*) \longrightarrow \mathbb{C}, \quad (f, h) \mapsto \int_X \langle f, h \rangle.$$

Similarly, we obtain the following local version:

Lemma 2.19. *Let S be a closed subset of X and U an open neighborhood of S in X . Then, we have the natural \mathfrak{g} -invariant bilinear form:*

$$C^\infty(U, \mathcal{V}) \times \mathcal{E}'_S(U, \mathcal{V}^*) \longrightarrow \mathbb{C}, \quad (f, h) \mapsto \int_U \langle f, h \rangle.$$

Moreover, if $S \subset U$ are both H -invariant subsets in X , then the bilinear form is also H -invariant.

We write $o = eH \in X$ for the origin. By Lemmas 2.15 and 2.19, we have obtained two (\mathfrak{g}, H) -invariant pairings:

$$\begin{aligned} C^\infty(X, \mathcal{V}) \times \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) &\longrightarrow \mathbb{C}, & (f, m) &\mapsto \Phi_V(f, m)(e), \\ C^\infty(X, \mathcal{V}) \times \mathcal{E}'_{\{o\}}(X, \mathcal{V}^*) &\longrightarrow \mathbb{C} & (f, h) &\mapsto \int_X \langle f, h \rangle. \end{aligned}$$

Let us show that there is a natural (\mathfrak{g}, H) -isomorphism between $\text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)$ and $\mathcal{E}'_{\{o\}}(X, \mathcal{V}^*)$. In fact, it follows from Lemma 2.15 that there exists an injective (\mathfrak{g}, H) -homomorphism

$$A : \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \rightarrow \mathcal{E}'_{\{o\}}(X, \mathcal{V}^*)$$

such that

$$\Phi_V(f, m)(e) = \int_X \langle f, A(m) \rangle \quad \text{for all } m \in \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \text{ and } f \in C^\infty(X, \mathcal{V}).$$

For a homogeneous vector bundle $\mathcal{V} = G \times_H V$ we define a vector-valued Dirac δ -function $\delta \otimes v^\vee \in \mathcal{E}'_{\{o\}}(X, \mathcal{V}^*)$, for $v \in V^\vee$ by

$$(2.21) \quad \langle f, \delta \otimes v^\vee \rangle := \langle f(e), v^\vee \rangle \quad \text{for } f \in C^\infty(X, \mathcal{V}) \simeq C^\infty(G, V)^H.$$

By the definition of Φ_V , we have

$$\Phi_V(f, 1 \otimes v^\vee)(e) = \langle f(e), v^\vee \rangle.$$

Hence $A(1 \otimes v^\vee) = \delta \otimes v$ by (2.21). Since A is a \mathfrak{g} -homomorphism, we have shown that

$$A(u \otimes v^\vee) = dL(u)(\delta \otimes v^\vee), \quad \text{for } u \in U(\mathfrak{g}), v \in V^\vee.$$

Lemma 2.20. *The (\mathfrak{g}, H) -homomorphism*

$$(2.22) \quad A : \text{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee) \longrightarrow \mathcal{E}'_{\{o\}}(X, \mathcal{V}^*), \quad u \otimes v^\vee \mapsto dL(u)(\delta \otimes v^\vee),$$

is bijective.

Proof. By Lemma 2.15 the map (2.22) is injective. Let us show that it is also surjective. By the structural theorem of (scalar-valued) distributions [S66, Chapter III, Théorème XXXVII], distributions supported on the singleton $\{o\}$ are obtained as a finite sum of derivatives of the Dirac's delta function. An analogous statement

holds for vector-bundle valued distributions supported on $\{o\}$, as we can see by trivializing the bundle near the origin o . Choose a complementary subspace $\mathfrak{q}(\mathbb{R})$ of $\mathfrak{h}(\mathbb{R}) = \text{Lie}(H)$ in $\mathfrak{g}(\mathbb{R}) = \text{Lie}(G)$. Since $dL(Z)$ ($Z \in \mathfrak{q}(\mathbb{R})$) spans the tangent space $T_o(G/H) \simeq \mathfrak{q}(\mathbb{R})$, any derivative of the vector-valued Dirac's delta function is given as a linear combination of elements of the form $dL(u)(\delta \otimes v^\vee)$ ($u \in U(\mathfrak{g}), v^\vee \in V^\vee$). Thus the map (2.22) is surjective. \square

Let $\mathbb{C}_{2\rho}$ denote the one-dimensional representation of H defined by

$$h \mapsto |\det(\text{Ad}_{G/H}(h) : \mathfrak{g}/\mathfrak{h} \rightarrow \mathfrak{g}/\mathfrak{h})|^{-1}.$$

If H is a parabolic subgroup of G with Langlands decomposition $P = MAN_+$ then the infinitesimal representation of $\mathbb{C}_{2\rho}$ is given by the sum of the roots for $\mathfrak{n}_+ = \text{Lie}(N_+)$. The bundle of densities $\Omega_{G/H}$ is given as a G -equivariant line bundle,

$$\Omega_{G/H} \simeq G \times_H |\det^{-1} \text{Ad}_{G/H}| \simeq G \times_H \mathbb{C}_{2\rho}.$$

For an H -module (λ, V) , we define a “twist” of the contragredient representation $\lambda_{2\rho}^\vee$ on the dual space V^\vee (or simply denoted by $V_{2\rho}^\vee$) by the formula

$$\lambda^* \equiv \lambda_{2\rho}^\vee := \lambda^\vee \otimes \mathbb{C}_{2\rho} = \lambda^\vee \otimes |\det^{-1} \text{Ad}_{G/H}|.$$

Then the dualizing bundle $\mathcal{V}^* = \mathcal{V}^\vee \otimes \Omega_{G/H}$ of the vector bundle $\mathcal{V} = G \times_H V$ is given, as a homogeneous vector bundle, by:

$$(2.23) \quad \mathcal{V}^* \equiv \mathcal{V}_{2\rho}^\vee \simeq G \times_H V_{2\rho}^\vee.$$

Then $\mathcal{D}'(X, \mathcal{V}^*)$ is identified with

$$(\mathcal{D}'(G) \otimes V_{2\rho}^\vee)^{\Delta(H)} = \{F \in \mathcal{D}'(G) \otimes V^\vee : \lambda_{2\rho}^\vee(h)F(\cdot h) = F(\cdot) \text{ for any } h \in H\}.$$

Now let us consider the setting of Theorem 2.9 where we have a G' -equivariant (but not necessarily injective) morphism from $Y = G'/H'$ to $X = G/H$.

Lemma 2.21. *Suppose that G' is a subgroup of G . Then the multiplication map*

$$m : G \times G' \rightarrow G, \quad (g, g') \mapsto (g')^{-1}g,$$

induces the isomorphism:

$$m^* : (\mathcal{D}'(X, \mathcal{V}^*) \otimes W)^{\Delta(H')} \xrightarrow{\sim} \mathcal{D}'(X \times Y, \mathcal{V}^* \boxtimes W)^{\Delta(G')}.$$

Proof. The image of the pull-back $m^* : \mathcal{D}'(G) \rightarrow \mathcal{D}'(G \times G')$ is $\mathcal{D}'(G \times G')^{\Delta(G')}$, where G' acts diagonally from the left. Thus, considering the remaining $G \times G'$ action from the right, we take $H \times H'$ -invariants with respect to the diagonal action in the $(G \times G' \times H \times H')$ -isomorphism:

$$m^* \otimes \text{id} \otimes \text{id} : \mathcal{D}'(G) \otimes V_{2\rho}^\vee \otimes W \xrightarrow{\sim} \mathcal{D}'(G \times G')^{\Delta(G')} \otimes V_{2\rho}^\vee \otimes W,$$

and therefore we get the lemma. \square

We recall from Section 2.1 that any continuous linear map $T : C^\infty(X, \mathcal{V}) \rightarrow C^\infty(Y, \mathcal{W})$ is given by a unique distribution kernel $K_T \in \mathcal{D}'(X \times Y, \mathcal{V}^* \boxtimes \mathcal{W})$. The following lemma gives a necessary and sufficient condition on the distribution K_T for the linear map T to be a G' -equivariant differential operator.

Lemma 2.22. *There is a natural linear isomorphism:*

$$(2.24) \quad \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) \xrightarrow{\sim} (\mathcal{D}'_{\{o\}}(X, \mathcal{V}^*) \otimes W)^{\Delta(H')}, \quad T \mapsto (m^*)^{-1}(K_T).$$

Proof. First, we show that the map (2.24) is well-defined. Suppose $T \in \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$. Since K_T is uniquely determined by T , the operator T is G' -equivariant, *i.e.* $L(g) \circ T \circ L(g^{-1}) = T$ for all $g \in G'$ if and only if $K_T \in \mathcal{D}'(X \times Y, \mathcal{V}^* \boxtimes \mathcal{W})^{\Delta(G')}$. By Lemma 2.3 the distribution kernel K_T is supported on the diagonal set $\Delta(Y) = \{(p(y), y) : y \in Y\} \subset X \times Y$. Via the bijection m^* given in Lemma 2.21 we thus have

$$\text{Supp}((m^*)^{-1}K_T) \subset \{o\}.$$

Hence the map (2.24) is well-defined. The injectivity of (2.24) is clear.

Conversely, take any element $k \in (\mathcal{D}'_{\{o\}}(X, \mathcal{V}^*) \otimes W)^{\Delta(H')}$. We set $K := m^*(k) \in \mathcal{D}'(X \times Y, \mathcal{V}^* \boxtimes \mathcal{W})^{\Delta(G')}$, and define a linear map

$$T : C^\infty(X, \mathcal{V}) \longrightarrow \mathcal{D}'(Y, \mathcal{W}), \quad f \mapsto \int_X f(x)K(x, \cdot).$$

Then T is G' -equivariant because K is $\Delta(G')$ -invariant.

Let us show that $Tf \in C^\infty(Y, \mathcal{W})$ for any $f \in C^\infty(X, \mathcal{V})$. To see this, we take neighborhoods U, U' and U'' of $x_o = p(y_o)$ in X , y_o in Y , and e in G' , respectively, such that $gU' \subset p^{-1}(U)$ for any $g \in U''$. Since the kernel K is supported on the diagonal set $\Delta(Y)$, $TF|_{U'}$ is locally of the form (2.4) as in the proof of Lemma 2.3.

Since T is G' -equivariant, we have

$$\sum_{\alpha} h_{\alpha}(y) \frac{\partial^{|\alpha|} f}{\partial x^{\alpha}}(gp(y)) = \sum_{\alpha} h_{\alpha}(gy) \frac{\partial^{|\alpha|} f}{\partial x^{\alpha}}(p(y)),$$

for any $y \in U', g \in U''$, and $f \in C^\infty(U) \otimes V$. By taking $f(x) = x^{\alpha} \otimes v$ ($\alpha \in \mathbb{N}^n$ and $v \in V$) as test functions, there are some $\varphi_{\alpha\beta} \in C^\infty(U'' \times U')$ for $|\beta| < |\alpha|$ such that

$$h_{\alpha}(gy) = h_{\alpha}(y) + \sum_{|\beta| < |\alpha|} \varphi_{\alpha\beta}(g, y) h_{\beta}(y).$$

Therefore we see inductively on $|\alpha|$ that $h_{\alpha}(y) \in C^\infty(U') \otimes \text{Hom}(V, W)$ for all α because G' acts transitively on Y . Hence $Tf|_{U'} \in C^\infty(U') \otimes W$. Thus we have shown that T maps $C^\infty(X, \mathcal{V})$ into $C^\infty(Y, \mathcal{W})$.

Finally, it follows from Lemma 2.3 that T is a differential operator because $\text{Supp}K \subset \Delta(Y)$. Now we have proved the lemma. \square

Proof of Theorem 2.9. Taking the tensor product of each term in (2.22) with the finite-dimensional representation W of H' , we get a bijection between the subspaces of \mathfrak{h}' -invariants:

$$\mathrm{Hom}_{\mathfrak{h}'}(W^\vee, \mathrm{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee)) \xrightarrow{\sim} (\mathcal{D}'_{\{o\}}(X, \mathcal{V}_X^*) \otimes W)^{\Delta(H')}.$$

Composing this with the bijection in Lemma 2.22, we obtain a bijection from $\mathrm{Hom}_{\mathfrak{h}'}(W^\vee, \mathrm{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$ to $\mathrm{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$, which is by construction nothing but $D_{X \rightarrow Y}$ in Theorem 2.9. \square

3. ALGEBRAIC FOURIER TRANSFORM FOR GENERALIZED VERMA MODULES

The duality theorem (Theorem 2.9) states that, to obtain a differential symmetry breaking operator $D \in \mathrm{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$, it suffices to find $\varphi \in \mathrm{Hom}_{H'}(W^\vee, \mathrm{ind}_{\mathfrak{h}}^{\mathfrak{g}}(V^\vee))$. In Section 4, we shall present a new method (*F-method*) which characterizes the “algebraic Fourier transform” of φ as a solution to a certain system of partial differential equations.

In this section we introduce and study the “algebraic Fourier transform” of generalized Verma modules. Proposition 3.11 is particularly important to the F-method.

3.1. Weyl algebra and algebraic Fourier transform. Let E be a vector space over \mathbb{C} . The Weyl algebra $\mathcal{D}(E)$ is the ring of holomorphic differential operators on E with polynomial coefficients.

Definition 3.1. We define the *algebraic Fourier transform* as an isomorphism of two Weyl algebras on E and its dual space E^\vee :

$$\mathcal{D}(E) \rightarrow \mathcal{D}(E^\vee), \quad T \mapsto \widehat{T},$$

induced by

$$(3.1) \quad \widehat{\frac{\partial}{\partial z_j}} := -\zeta_j, \quad \widehat{z_j} := \frac{\partial}{\partial \zeta_j}, \quad 1 \leq j \leq n = \dim E.$$

where (z_1, \dots, z_n) are coordinates on E and $(\zeta_1, \dots, \zeta_n)$ are the dual coordinates on E^\vee .

Example 3.2. Let $E_z := \sum_{j=1}^n z_j \frac{\partial}{\partial z_j}$ be the Euler operator on E . Then, by the commutation relations

$$(3.2) \quad \frac{\partial}{\partial \zeta_i} \zeta_j - \zeta_j \frac{\partial}{\partial \zeta_i} = \delta_{ij},$$

in the Weyl algebra $\mathcal{D}(E^\vee)$, where δ_{ij} is the Kronecker delta. Hence we have $\widehat{E_z} = -E_\zeta - n$.

The isomorphism $T \mapsto \widehat{T}$ in Definition 3.1 does not depend on the choice of coordinates. To see this, we consider the natural action of the general linear group $GL(E)$ on E , which yields automorphisms of the ring $\text{Pol}(E)$ of polynomials of E and the Weyl algebra $\mathcal{D}(E)$. For $A \in GL(E)$, we set

$$\begin{aligned} A_{\#} &: \text{Pol}(E) \longrightarrow \text{Pol}(E), & F &\mapsto F(A^{-1}\cdot), \\ A_{*} &: \mathcal{D}(E) \longrightarrow \mathcal{D}(E), & T &\mapsto A_{\#} \circ T \circ A_{\#}^{-1}. \end{aligned}$$

We denote by ${}^tA \in GL(E^{\vee})$ the dual map of A . Then we have

Lemma 3.3. *For any $A \in GL(E)$ and $T \in \mathcal{D}(E)$,*

$$\widehat{A_{*}T} = ({}^tA^{-1})_{*} \widehat{T}.$$

The proof is straightforward from the definition (3.1), and we omit it.

Next we consider the group homomorphism $GL(E) \longrightarrow GL(\text{Pol}(E))$, $A \mapsto A_{\#}$. Taking the differential, we get a Lie algebra homomorphism $\text{End}(E) \rightarrow \mathcal{D}(E)$. In the coordinates, we write $Z = {}^t(z_1, \dots, z_n)$ and $\partial_Z = {}^t\left(\frac{\partial}{\partial z_1}, \dots, \frac{\partial}{\partial z_n}\right)$. Then this homomorphism amounts to

$$(3.3) \quad \Psi_E : \text{End}(E) \rightarrow \mathcal{D}(E), \quad A \mapsto -{}^tZ {}^tA \partial_Z \equiv -\sum_{i,j} A_{ij} z_j \frac{\partial}{\partial z_i}.$$

Let $\sigma : \mathfrak{g} \rightarrow \text{End}(E)$ be a representation of a Lie algebra \mathfrak{g} on E , and $\sigma^{\vee} : \mathfrak{g} \rightarrow \text{End}(E^{\vee})$ the contragredient representation. Then the algebraic Fourier transform $T \mapsto \widehat{T}$ relates the two Lie algebra homomorphisms $\Psi_E \circ \sigma : \mathfrak{g} \rightarrow \mathcal{D}(E)$ and $\Psi_{E^{\vee}} \circ \sigma^{\vee} : \mathfrak{g} \rightarrow \mathcal{D}(E^{\vee})$ as follows:

Lemma 3.4.

$$\widehat{\Psi_E \circ \sigma} = \Psi_{E^{\vee}} \circ \sigma^{\vee} + (\text{Trace} \circ \sigma) \cdot \text{id}_{E^{\vee}}.$$

Proof. In the coordinates, we write $A := \sigma(Z) \in \text{End}(E) \simeq M(n, \mathbb{C})$ for $Z \in \mathfrak{g}$. Then,

$$\begin{aligned} \widehat{\Psi_E \circ \sigma}(Z) - \Psi_{E^{\vee}} \circ \sigma^{\vee}(Z) &= -{}^tZ {}^tA \partial_Z - {}^t\zeta A \partial_{\zeta} \\ &= {}^t\partial_{\zeta} {}^tA \zeta - {}^t\zeta A \partial_{\zeta} \\ &= (\text{Trace } A) \text{id}_{E^{\vee}}, \end{aligned}$$

where the last equality follows from the commutation relations (3.2). \square

For actual computations that will be undertaken in a subsequent paper [KP15-2], it is convenient to give another interpretation of the algebraic Fourier transform by using real forms of E .

Definition 3.5. Fix a real form $E(\mathbb{R})$ of the complex vector space E . Let $\mathcal{E}'_{\{0\}}(E(\mathbb{R}))$ be the space of distributions on the vector space $E(\mathbb{R})$ supported at the origin 0. We define a “Fourier transform” $\mathcal{F}_c : \mathcal{E}'_{\{0\}}(E(\mathbb{R})) \rightarrow \text{Pol}(E^\vee)$ by the following formula:

$$(3.4) \quad \mathcal{F}_c f(\zeta) := \langle f(\cdot), e^{\langle \cdot, \zeta \rangle} \rangle = \int_{E(\mathbb{R})} e^{\langle x, \zeta \rangle} f(x) \quad \text{for } \zeta \in E^\vee.$$

We have used the function $e^{\langle x, \zeta \rangle}$ in (3.4) rather than $e^{-\sqrt{-1}\langle x, \zeta \rangle}$ or $e^{-\langle x, \zeta \rangle}$ which are involved in the usual Fourier transform or the Laplace transform, respectively. This convention makes later computations simpler (see Remark 4.2).

Furthermore, with our convention

$$(3.5) \quad \mathcal{F}_c(f(A \cdot)) = (\mathcal{F}_c f)(\text{}^t A^{-1} \cdot),$$

for any $A \in GL_{\mathbb{R}}(E(\mathbb{R}))$.

The Fourier transform \mathcal{F}_c induces an algebra isomorphism

$$\mathcal{F}_c : \mathcal{E}'_{\{0\}}(E(\mathbb{R})) \xrightarrow{\sim} \text{Pol}(E^\vee)$$

between the polynomial algebra $\text{Pol}(E^\vee)$ with unit $\mathbf{1}$, the constant function on E^\vee , and the convolution algebra $\mathcal{E}'_{\{0\}}(E(\mathbb{R}))$ with unit δ , the Dirac delta function. We write $\mathcal{F}_c^{-1} : \text{Pol}(E^\vee) \xrightarrow{\sim} \mathcal{E}'_{\{0\}}(E(\mathbb{R}))$ for the inverse “Fourier transform”:

$$\mathcal{F}_c^{-1}(\mathbf{1}) = \delta.$$

Remark 3.6. The Weyl algebra $\mathcal{D}(E)$ acts naturally on the space of distributions on $E(\mathbb{R})$, and in particular, on $\mathcal{E}'_{\{0\}}(E(\mathbb{R}))$. The algebraic Fourier transform defined in Definition 3.1 satisfies

$$(3.6) \quad \widehat{T} = \mathcal{F}_c \circ T \circ \mathcal{F}_c^{-1} \quad \text{for } T \in \mathcal{D}(E),$$

and the formula (3.6) characterizes \widehat{T} . To see this, we take coordinates (x_1, \dots, x_n) on $E(\mathbb{R})$, and extend them to the complex coordinates (z_1, \dots, z_n) on E and the dual ones $(\zeta_1, \dots, \zeta_n)$ on E^\vee . Let $P(\zeta) = \zeta^\alpha \in \text{Pol}(E^\vee)$ and $T = \sum_{\beta, \gamma} a_{\beta, \gamma} z^\beta \frac{\partial^{|\gamma|}}{\partial z^\gamma} \in \mathcal{D}(E)$. Then we have

$$\widehat{T}P = \sum_{\beta, \gamma} (-1)^{|\gamma|} a_{\beta, \gamma} \frac{\partial^{|\beta|}}{\partial z^\beta} \zeta^{\alpha+\gamma},$$

and on the other hand,

$$\begin{aligned} \mathcal{F}_c \circ T \circ \mathcal{F}_c^{-1}P &= (-1)^{|\alpha|} \mathcal{F}_c \circ T(\delta^\alpha(x)) = (-1)^{|\alpha|} \mathcal{F}_c \left(\sum_{\beta, \gamma} a_{\beta, \gamma} x^\beta \delta^{\alpha+\gamma}(x) \right) \\ &= (-1)^{|\alpha|} \sum_{\beta, \gamma} (-1)^{|\alpha|+|\gamma|} a_{\beta, \gamma} \frac{\partial^{|\beta|}}{\partial z^\beta} \zeta^{\alpha+\gamma}. \end{aligned}$$

Hence the identity (3.6) holds on $\text{Pol}(E^\vee)$. Since the Weyl algebra $\mathcal{D}(E^\vee)$ acts faithfully on $\text{Pol}(E^\vee)$, we have shown (3.6). In particular, the composition $\mathcal{F}_c \circ T \circ \mathcal{F}_c^{-1}$ does not depend on the choice of a real form $E(\mathbb{R})$.

3.2. Holomorphic vector fields associated to the Gelfand–Naimark decomposition. It is convenient to prepare some notation in the *complex* reductive Lie algebras for later purpose.

Let \mathfrak{g} be a complex reductive Lie algebra, and $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$ a Levi decomposition of a parabolic subalgebra. Let $G_{\mathbb{C}}$ be a connected complex Lie group with Lie algebra \mathfrak{g} , and $P_{\mathbb{C}} = L_{\mathbb{C}} \exp \mathfrak{n}_+$ the parabolic subgroup with Lie algebra $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$. According to the Gelfand–Naimark decomposition $\mathfrak{g} = \mathfrak{n}_- + \mathfrak{l} + \mathfrak{n}_+$ of the Lie algebra \mathfrak{g} , we have a diffeomorphism

$$\mathfrak{n}_- \times L_{\mathbb{C}} \times \mathfrak{n}_+ \rightarrow G_{\mathbb{C}}, \quad (Z, \ell, Y) \mapsto (\exp Z)\ell(\exp Y),$$

into an open dense subset $G_{\mathbb{C}}^{\text{reg}}$ of $G_{\mathbb{C}}$. Let

$$p_{\pm} : G_{\mathbb{C}}^{\text{reg}} \longrightarrow \mathfrak{n}_{\pm}, \quad p_o : G_{\mathbb{C}}^{\text{reg}} \rightarrow L_{\mathbb{C}},$$

be the projections characterized by the identity

$$\exp(p_-(g))p_o(g)\exp(p_+(g)) = g.$$

We set

$$(3.7) \quad \alpha : \mathfrak{g} \times \mathfrak{n}_- \rightarrow \mathfrak{l}, \quad (Y, Z) \mapsto \left. \frac{d}{dt} \right|_{t=0} p_o(e^{tY}e^Z),$$

$$(3.8) \quad \beta : \mathfrak{g} \times \mathfrak{n}_- \rightarrow \mathfrak{n}_-, \quad (Y, Z) \mapsto \left. \frac{d}{dt} \right|_{t=0} p_-(e^{tY}e^Z).$$

$$(3.9) \quad \gamma : \mathfrak{g} \times \mathfrak{n}_- \rightarrow \mathfrak{l} + \mathfrak{n}_+, \quad (Y, Z) \mapsto \alpha(Y, Z) + \left. \frac{d}{dt} \right|_{t=0} p_+(e^{tY}e^Z).$$

We regard $\beta(Y, \cdot)$ as a holomorphic vector field on \mathfrak{n}_- through the following identification.

$$\mathfrak{n}_- \ni Z \mapsto \beta(Y, Z) \in \mathfrak{n}_- \simeq T_Z \mathfrak{n}_-.$$

Example 3.7. $G_{\mathbb{C}} = GL(p+q, \mathbb{C})$, $L_{\mathbb{C}} = GL(p, \mathbb{C}) \times GL(q, \mathbb{C})$, and $\mathfrak{n}_- \simeq M(p, q; \mathbb{C})$.

We note that \mathfrak{n}_- is realized as upper block matrices. Then for $g^{-1} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_{\mathbb{C}}$,

$Y = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in M(p+q; \mathbb{C})$ and $Z \in M(p, q; \mathbb{C})$ we have

$$\begin{aligned} p_-(g^{-1}) &= bd^{-1}, \\ p_o(g^{-1}) &= (a - bd^{-1}c, d) \in GL(p, \mathbb{C}) \times GL(q, \mathbb{C}), \\ \alpha(Y, Z) &= (A - ZC, CZ + D) \in \mathfrak{gl}_p(\mathbb{C}) \oplus \mathfrak{gl}_q(\mathbb{C}), \\ \beta(Y, Z) &= AZ + B - ZCZ - ZD. \end{aligned}$$

Then $\beta(Y, \cdot)$ is regarded as the following holomorphic vector field on $\mathfrak{n}_- \simeq M(p, q; \mathbb{C})$ given by

$$\begin{aligned} \text{Trace}(\beta(Y, Z) {}^t\partial_Z) &= \sum_{a=1}^p \sum_{b=1}^q \beta(Y, Z)_{ab} \frac{\partial}{\partial z_{ab}} \\ &= \sum_{a=1}^p \sum_{b=1}^q \left(\sum_{i=1}^p A_{ai} z_{ib} + B_{ab} - \sum_{i=1}^p \sum_{j=1}^q z_{aj} C_{ji} z_{ib} - \sum_{j=1}^q z_{aj} D_{jb} \right) \frac{\partial}{\partial z_{ab}}. \end{aligned}$$

A reductive Lie algebra \mathfrak{g} is said to be k -graded if it admits a direct sum decomposition $\mathfrak{g} = \bigoplus_{j=-k}^k \mathfrak{g}(j)$ such that $[\mathfrak{g}(i), \mathfrak{g}(j)] \subset \mathfrak{g}(i+j)$ for all i, j . Any parabolic subalgebra $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$ of \mathfrak{g} is given by $\mathfrak{l} = \mathfrak{g}(0)$ and $\mathfrak{n}_+ = \bigoplus_{j>0} \mathfrak{g}(j)$ for some k -gradation of \mathfrak{g} . We then have the following estimates of coefficients of holomorphic differential operators $d\pi_\mu(Y)$.

Lemma 3.8. *According to the direct sum decomposition $\mathfrak{g} = \bigoplus_{j=-k}^k \mathfrak{g}(j)$, we write $\gamma(Y, Z) = \sum_{\ell=0}^k \gamma_\ell$ and $\beta(Y, Z) = \sum_{\ell=-k}^{-1} \beta_\ell$, where $\gamma_\ell \in \mathfrak{g}(\ell)$ for $0 \leq \ell \leq k$ and $\beta_\ell \in \mathfrak{g}(\ell)$ for $-k \leq \ell \leq -1$. Then γ_ℓ and β_ℓ are polynomials in Z of degree at most $k - \ell$.*

Proof. Since the map $N_{-, \mathbb{C}} \times P_{\mathbb{C}} \xrightarrow{\sim} G_{\mathbb{C}}^{\text{reg}}$ is an analytic diffeomorphism, we have

$$(3.10) \quad e^{tY} e^Z = e^{Z+t\beta(Y,Z)+o(t)} e^{t\alpha(Y,Z)+o(t)}$$

for sufficiently small $t \in \mathbb{C}$, where we use the Landau symbol $o(t)$ for a \mathfrak{g} -valued function dominated by t when t tends to be zero. Multiplying (3.10) by e^{-Z} from the left, and taking the differential at $t = 0$, we get

$$\text{Ad}(e^{-Z})Y = \gamma(Y, Z) + \frac{e^{\text{ad}(Z)} - 1}{\text{ad}(Z)} \beta(Y, Z),$$

because $\left. \frac{d}{dt} \right|_{t=0} e^{-Z} e^{Z+tW} = \frac{e^{\text{ad}(Z)} - 1}{\text{ad}(Z)} W$. We note that $\text{ad}(Z)$ lowers the grading of \mathfrak{g} , namely $\text{ad}(Z)\mathfrak{g}(j) \subset \bigoplus_{i=-k}^{j-1} \mathfrak{g}(i)$ because $Z \in \mathfrak{n}_-$. In particular, we have

$$(3.11) \quad \sum_{j=0}^{2k} \frac{(-1)^j}{j!} \text{ad}(Z)^j Y = \gamma(Y, Z) + \sum_{i=0}^{k-1} \frac{\text{ad}(Z)^i}{(i+1)!} \beta(Y, Z).$$

Let $q_\ell : \mathfrak{g} \rightarrow \mathfrak{g}(\ell)$ be the projection according to the direct sum decomposition $\mathfrak{g} = \bigoplus_{j=-k}^k \mathfrak{g}(j)$. Suppose $\ell \geq 0$. Applying q_ℓ to (3.11), we have

$$\gamma_\ell = q_\ell \left(\sum_{j=0}^{k-\ell} \frac{(-1)^j \text{ad}(Z)^j}{j!} Y \right).$$

Hence γ_ℓ is a polynomial in Z of degree at most $k - \ell$.

Suppose $\ell < 0$. Applying q_ℓ to (3.11), we get

$$\beta_\ell = q_\ell \left(\sum_{j=0}^{k-\ell} \frac{(-1)^j \text{ad}(Z)^j}{j!} Y \right) - q_\ell \left(\sum_{i=0}^{j-\ell-1} \sum_{j=\ell}^{-1} \frac{\text{ad}(Z)^i}{(i+1)!} \beta_j \right).$$

By the downward induction on ℓ , we see that β_ℓ is a polynomial in Z of degree at most $k - \ell$ for $-k \leq \ell \leq -1$. \square

3.3. Fourier transform of principal series representations. Suppose \mathfrak{g} is a complex reductive Lie algebra, $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$ a parabolic subalgebra, and $\lambda : \mathfrak{p} \rightarrow \text{End}_{\mathbb{C}}(V)$ a finite-dimensional representation.

We use the letter μ to denote the representation of \mathfrak{p} on the dual space V^\vee given by

$$(3.12) \quad \mu := \lambda^* \equiv \lambda^\vee \otimes \text{Trace}(\text{ad}(\cdot) : \mathfrak{n}_+ \rightarrow \mathfrak{n}_+).$$

By applying the (algebraic) Fourier transform of the Weyl algebra, we define a Lie algebra homomorphism

$$\widehat{d\pi}_\mu : \mathfrak{g} \rightarrow \mathcal{D}(\mathfrak{n}_+) \otimes \text{End}_{\mathbb{C}}(V^\vee),$$

by using the complex flag variety $G_{\mathbb{C}}/P_{\mathbb{C}}$ in this subsection. In Section 3.4, we relate $\widehat{d\pi}_\mu$ with the ‘‘algebraic Fourier transform’’ of a generalized Verma module

$$F_c : \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \xrightarrow{\sim} \text{Pol}(\mathfrak{n}_+) \otimes V^\vee,$$

which is defined by using a real flag variety G/P , see (3.23).

Let $G_{\mathbb{C}}$ be a connected complex reductive Lie group with complex reductive Lie algebra \mathfrak{g} , and $P_{\mathbb{C}} = L_{\mathbb{C}}N_{+, \mathbb{C}}$ be the parabolic subgroup with Lie algebra \mathfrak{p} . Let $\Omega_{X_{\mathbb{C}}}$ be the canonical line bundle of the complex generalized flag variety $X_{\mathbb{C}} = G_{\mathbb{C}}/P_{\mathbb{C}}$.

Suppose λ lifts to a holomorphic representation of $P_{\mathbb{C}}$, then so does μ . We form a $G_{\mathbb{C}}$ -equivariant holomorphic vector bundle \mathcal{V} and $\mathcal{V}^{\vee} \otimes \Omega_{X_{\mathbb{C}}}$ over $X_{\mathbb{C}}$ associated to λ and μ , respectively.

We consider the regular representation π_{μ} of $G_{\mathbb{C}}$ on $C^{\infty}(G_{\mathbb{C}}/P_{\mathbb{C}}, \mathcal{V}^{\vee} \otimes \Omega_{X_{\mathbb{C}}})$. The infinitesimal action will be denoted by $d\pi_{\mu}$, which is defined on $C^{\infty}(U, \mathcal{V}^{\vee} \otimes \Omega_{X_{\mathbb{C}}|_U})$ for any open subset U of $G_{\mathbb{C}}/P_{\mathbb{C}}$. In particular, we take U to be the open Bruhat cell $\mathfrak{n}_{-} \hookrightarrow G_{\mathbb{C}}/P_{\mathbb{C}}$, $Z \mapsto \exp Z \cdot o$, where $o = eP_{\mathbb{C}} \in G_{\mathbb{C}}/P_{\mathbb{C}}$. By trivializing the holomorphic vector bundle $\mathcal{V}^{\vee} \otimes \Omega_{X_{\mathbb{C}}} \rightarrow G_{\mathbb{C}}/P_{\mathbb{C}}$ on it, we define a function $F \in C^{\infty}(\mathfrak{n}_{-}, V^{\vee})$ for a section $f \in C^{\infty}(G_{\mathbb{C}}/P_{\mathbb{C}}, \mathcal{V}^{\vee} \otimes \Omega_{X_{\mathbb{C}}})$ by

$$F(Z) := f(\exp Z) \quad \text{for } Z \in \mathfrak{n}_{-}.$$

Then the action of \mathfrak{g} on $C^{\infty}(\mathfrak{n}_{-}, V^{\vee})$ given by

$$\begin{aligned} (d\pi_{\mu}(Y)F)(Z) &= \left. \frac{d}{dt} \right|_{t=0} f(e^{-tY} e^Z) \\ (3.13) \quad &= \mu(\gamma(Y, Z))F(Z) - (\beta(Y, \cdot)F)(Z) \quad \text{for } Y \in \mathfrak{g}, \end{aligned}$$

where by a little abuse of notation μ stands for the infinitesimal action. The right-hand side of (3.13) defines a representation of Lie algebra \mathfrak{g} whenever μ (or λ) is a representation of the Lie algebra \mathfrak{p} without assuming that it lifts to a holomorphic representation of the complex reductive group $P_{\mathbb{C}}$.

It follows from (3.13) and Lemma 3.8 that we obtain a Lie algebra homomorphism

$$(3.14) \quad d\pi_{\mu} : \mathfrak{g} \rightarrow \mathcal{D}(\mathfrak{n}_{-}) \otimes \text{End}(V^{\vee}),$$

for any representation λ of the Lie algebra \mathfrak{p} . By taking the algebraic Fourier transform on the Weyl algebra $\mathcal{D}(\mathfrak{n}_{-})$ (see Definition 3.1), we get another Lie algebra homomorphism:

$$(3.15) \quad \widehat{d\pi_{\mu}} : \mathfrak{g} \rightarrow \mathcal{D}(\mathfrak{n}_{+}) \otimes \text{End}(V^{\vee}).$$

We use the same letter π_{μ} to denote the ‘‘action’’ of $G_{\mathbb{C}}$ on $C^{\infty}(\mathfrak{n}_{-}, V^{\vee})$ given as

$$(3.16) \quad (\pi_{\mu}(g)F)(Z) = \mu(p_o(g^{-1} \exp Z) \exp(p_+(g^{-1} \exp Z)))^{-1} F(p_-(g^{-1} \exp Z)).$$

This formula makes sense if F comes from $C^{\infty}(G_{\mathbb{C}}/P_{\mathbb{C}}, \mathcal{V} \otimes \Omega_{X_{\mathbb{C}}}^{\vee})$, or if $F \in C^{\infty}(\mathfrak{n}_{-}, V^{\vee})$ and $g \in G_{\mathbb{C}}$ and $Z \in \mathfrak{n}_{-}$ satisfy $g^{-1} \exp Z \in G_{\mathbb{C}}^{\text{reg}}$. In particular, if λ is trivial on the nilpotent radical \mathfrak{n}_{+} for $g = m \exp W$ with $m \in L_{\mathbb{C}}$ and $W \in \mathfrak{n}_{-}$, and if \mathfrak{n}_{+} is abelian we have

$$(3.17) \quad (\pi_{\mu}(g)F)(Z) = \mu(m)F(\text{Ad}(m)^{-1}Z - W).$$

Let us analyze $d\pi_{\mu}(Y)$ for $Y \in \mathfrak{l} + \mathfrak{n}_{+}$. We begin with the case $Y \in \mathfrak{l}$. We let the Levi subgroup $L_{\mathbb{C}}$ act on $\text{Pol}(\mathfrak{n}_{+})$ by

$$\text{Ad}_{\#}(l) : f(\cdot) \mapsto f(\text{Ad}(l^{-1})\cdot), \quad l \in L_{\mathbb{C}}.$$

Since this action is algebraic, the infinitesimal action defines a Lie algebra homomorphism into the Weyl algebra:

$$\mathrm{ad}_\# : \mathfrak{l} \rightarrow \mathcal{D}(\mathfrak{n}_+), \quad Y \mapsto \mathrm{ad}_\#(Y),$$

where $\mathrm{ad}_\#(Y)$ is a holomorphic vector field on \mathfrak{n}_+ given by $\mathrm{ad}_\#(Y)_x := \left. \frac{d}{dt} \right|_{t=0} \mathrm{Ad}(e^{-tY})x \in T_x(\mathfrak{n}_+)$ for $x \in \mathfrak{n}_+$.

Lemma 3.9. *Let λ be a representation of the parabolic Lie algebra $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$, and $\mu \equiv \lambda^*$ be as in (3.12). Then the following two representations of \mathfrak{l} on $\mathrm{Pol}(\mathfrak{n}_+) \otimes V^\vee$ are isomorphic:*

$$(3.18) \quad \widehat{d\pi_\mu}|_{\mathfrak{l}} \simeq \mathrm{ad}_\# \otimes \mathrm{id} + \mathrm{id} \otimes (\mu - \mathrm{Trace} \circ \mathrm{ad}|_{\mathfrak{n}_-}) = \mathrm{ad}_\#(Y) \otimes \mathrm{id} + \mathrm{id} \otimes (-\lambda).$$

In particular, if λ lifts to a holomorphic representation of $P_{\mathbb{C}}$ then the right-hand side is the infinitesimal action of $\mathrm{Ad}_\# \otimes \lambda^\vee$ of $L_{\mathbb{C}}$ on $\mathrm{Pol}(\mathfrak{n}_+) \otimes V^\vee$.

Proof. For $Y \in \mathfrak{l}$, $X \in \mathfrak{n}_-$ we have $\gamma(Y, X) = Y$, and the formula (3.13) reduces, in $\mathcal{D}(\mathfrak{n}_-) \otimes \mathrm{End}(V^\vee)$, to

$$d\pi_\mu(Y) = \mathrm{id} \otimes \mu(Y) - \beta(Y, \cdot) \otimes \mathrm{id}.$$

We apply Lemma 3.4 to the case where (σ, E) is the adjoint representation of \mathfrak{l} on \mathfrak{n}_- . Since $\beta(Y, \cdot) = -dL(Y)$ for $Y \in \mathfrak{l}$, we have $\Psi_{\mathfrak{n}_-} \circ \mathrm{ad} = -\beta$ on \mathfrak{l} , with the notation therein. Moreover, via the identification $\mathfrak{n}_-^\vee \simeq \mathfrak{n}_+$, the map $\Psi_{\mathfrak{n}_-^\vee} \circ \mathrm{ad}^\vee$ amounts to $\Psi_{\mathfrak{n}_+} \circ \mathrm{ad} = \mathrm{ad}_\#$. Therefore, we get

$$\begin{aligned} \widehat{d\pi_\mu}(Y) &= \mathrm{id} \otimes \mu(Y) + \Psi_{\mathfrak{n}_-^\vee} \circ \mathrm{ad}^\vee(Y) \otimes \mathrm{id} + \left(\mathrm{Trace} \circ \mathrm{ad}(Y) \Big|_{\mathfrak{n}_-} \right) \mathrm{id} \otimes \mathrm{id} \\ &= \mathrm{id} \otimes \mu(Y) + \mathrm{ad}_\#(Y) \otimes \mathrm{id} - \left(\mathrm{Trace} \circ \mathrm{ad}(Y) \Big|_{\mathfrak{n}_+} \right) \mathrm{id} \otimes \mathrm{id}. \end{aligned}$$

Thus, the lemma follows. \square

The differential operators $\widehat{d\pi_\mu}(Y)$ with $Y \in \mathfrak{n}_+$ play a central role in the F-method. If the parabolic subalgebra \mathfrak{p} is associated to a k -gradation of \mathfrak{g} , then these differential operators are at most of order $2k$ by Lemma 3.8. We describe their structure in the case where $k = 1$, namely \mathfrak{n}_+ is abelian.

Proposition 3.10. *Assume that \mathfrak{n}_+ is abelian. Let (λ, V) be a representation of \mathfrak{l} , extended trivially on \mathfrak{n}_+ , and $\mu = \lambda^*$ be as in (3.12). For every $Y \in \mathfrak{n}_+$ the operator $\widehat{d\pi_\mu}(Y)$ is of the form*

$$(3.19) \quad \sum a_i^{jk} \zeta^i \frac{\partial^2}{\partial \zeta^j \partial \zeta^k} + \sum b^j \frac{\partial}{\partial \zeta^j},$$

where a_i^{jk} and $b^j \in \mathrm{End}(V^\vee)$ are constants depending on Y .

Proof. Since \mathfrak{n}_+ is abelian, we can take a characteristic element H such that

$$\mathrm{Ad}(e^{sH})Y = e^s Y \quad \text{for any } Y \in \mathfrak{n}_+.$$

We set $m := e^{sH}$. Then ${}^t\mathrm{Ad}(m)^{-1} = e^{-s}\mathrm{id}$ on $\mathfrak{n}_- \simeq \mathfrak{n}_+^\vee$.

Taking the algebraic Fourier transform of the formula

$$d\pi_\mu(\mathrm{Ad}(m)Y) = \pi_\mu(m)d\pi_\mu(Y)\pi_\mu(m^{-1}),$$

where $(\pi_\mu(m)F)(Z) = \mu(m)F(\mathrm{Ad}(m^{-1})Z) = \mu(m)F(\mathrm{Ad}(m)_\#F)(Z)$ by (3.17), we get

$$\widehat{d\pi_\mu}(\mathrm{Ad}(m)Y) = ({}^t\mathrm{Ad}(m)^{-1})_* \widehat{d\pi_\mu}(Y)$$

by Lemma 3.3. Hence

$$(3.20) \quad e^s \widehat{d\pi_\mu}(Y) = (e^{-s}\mathrm{id})_* \widehat{d\pi_\mu}(Y)$$

If we write $\widehat{d\pi_\mu}(Y)$ in the form

$$\sum_{\alpha, \beta \in \mathbb{N}^n} C_{\alpha, \beta} \zeta^\alpha \frac{\partial^{|\beta|}}{\partial \zeta^\beta}$$

then (3.20) implies that $C_{\alpha, \beta} \neq 0$ only when $|\alpha| + |\beta| = -1$ because $(e^{-s}\mathrm{id})_* \frac{\partial}{\partial \zeta_j} = e^{-s} \frac{\partial}{\partial \zeta_j}$ and $(e^{-s}\mathrm{id})_* \zeta_j = e^s \zeta_j$ ($1 \leq j \leq n$). As $d\pi_\mu(Y)$ is a vector field there is no term for $|\alpha| > 1$. Hence we get the expression (3.19). \square

3.4. Fourier transform on the real flag varieties. In this subsection we define ‘‘algebraic Fourier transform’’ of generalized Verma modules, see (3.23):

$$F_c : \mathrm{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \xrightarrow{\sim} \mathrm{Pol}(\mathfrak{n}_+) \otimes V^\vee.$$

As we shall prove in Proposition 3.11, the Lie algebra homomorphism $\widehat{d\pi_{\lambda^*}} : U(\mathfrak{g}) \longrightarrow \mathcal{D}(\mathfrak{n}_+) \otimes \mathrm{End}(V^\vee)$ defined in (3.15) in the previous section can be reconstructed from F_c , namely, $\widehat{d\pi_{\lambda^*}}(u)$ ($u \in U(\mathfrak{g})$) is the operator S that is characterized by

$$SF_c(v) = F_c(u \cdot v) \quad \text{for any } v \in \mathrm{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee).$$

For later purpose, we work with a real form G of $G_{\mathbb{C}}$. From now on, let G be a real semisimple Lie group, P a parabolic subgroup of G with Levi decomposition $P = LN_+$, and V a finite-dimensional representation of P .

Let LN_- be the opposite parabolic subgroup of $P = LN_+$. We write $\mathfrak{n}_+(\mathbb{R})$ and $\mathfrak{n}_-(\mathbb{R})$ for the Lie algebras of N_+ and N_- , respectively, and set $\mathfrak{n}_+ = \mathfrak{n}_+(\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$. The open Bruhat cell is given as the image of the following embedding

$$\iota : \mathfrak{n}_-(\mathbb{R}) \hookrightarrow G/P, \quad X \mapsto \exp(X) \cdot o,$$

where $o = eP \in G/P$.

Let $\lambda : P \rightarrow GL_{\mathbb{C}}(V)$ be a finite-dimensional representation of P , and $\mathcal{V} = G \times_P V$ the G -equivariant vector bundle over the real flag variety G/P . The pullback of the

dualizing bundle $\mathcal{V}^* \equiv \mathcal{V}_{2\rho}^\vee \rightarrow G/P$ via ι is trivialized into the direct product bundle $\mathfrak{n}_-(\mathbb{R}) \times V^\vee \rightarrow \mathfrak{n}_-(\mathbb{R})$ and thus we have a linear isomorphism:

$$(3.21) \quad \iota^* : \mathcal{E}'_{\{o\}}(G/P, \mathcal{V}_{2\rho}^\vee) \xrightarrow{\sim} \mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee,$$

through which we induce the (\mathfrak{g}, P) -action on $\mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee$ from $\mathcal{E}'_{\{o\}}(G/P, \mathcal{V}_{2\rho}^\vee)$.

The Killing form of \mathfrak{g} identifies the dual space $\mathfrak{n}_-(\mathbb{R})^\vee$ with $\mathfrak{n}_+(\mathbb{R})$, and thus the Fourier transform \mathcal{F}_c in (3.4) gives rise to a linear isomorphism:

$$(3.22) \quad \mathcal{F}_c \otimes \text{id} : \mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee \xrightarrow{\sim} \text{Pol}(\mathfrak{n}_+) \otimes V^\vee,$$

through which we induce the (\mathfrak{g}, P) -action further on the right-hand side.

In summary we have the following (\mathfrak{g}, P) -isomorphisms:

$$(3.23) \quad F_c : \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \xrightarrow{(2.22)} \mathcal{E}'_{\{o\}}(G/P, \mathcal{V}_{2\rho}^\vee) \xrightarrow{(3.21)} \mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee \xrightarrow{\mathcal{F}_c \otimes \text{id}} \text{Pol}(\mathfrak{n}_+) \otimes V^\vee.$$

We say that F_c is the *algebraic Fourier transform of a generalized Verma module*.

The (\mathfrak{g}, P) -module structure of $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ is described by the following proposition.

Proposition 3.11. *Let (λ, V) be a finite-dimensional representation of P and define another representation of P on the dual space V^\vee by $\mu := \lambda^* \equiv \lambda^\vee \otimes \mathbb{C}_{2\rho}$. Then,*

- 1) *The \mathfrak{g} -action on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ induced by F_c in (3.23) coincides with the one given by $\widehat{d\pi_\mu}$ in (3.15).*
- 2) *The L action on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ induced by F_c in (3.23) coincides with the one given by $\text{Ad}_\# \otimes \lambda^\vee$.*

Proof. 1) Let $G_{\mathbb{C}}$ be a complexification of G and $P_{\mathbb{C}}$ the connected subgroup of $G_{\mathbb{C}}$ with Lie algebra $\mathfrak{p} = \text{Lie}(P) \otimes_{\mathbb{R}} \mathbb{C}$. First we assume that λ extends to a holomorphic representation of $P_{\mathbb{C}}$. Then the G -equivariant vector bundle $\mathcal{V}_{2\rho}^\vee$ over $X = G/P$ is the restriction of the $G_{\mathbb{C}}$ -equivariant holomorphic vector bundle $\mathcal{V}^\vee \otimes \Omega_{X_{\mathbb{C}}}$ over $X_{\mathbb{C}} = G_{\mathbb{C}}/P_{\mathbb{C}}$ that was introduced in the previous subsection. Therefore, the action of $Y \in \mathfrak{g}$ on $\mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee$ induced by ι^* in (3.21) is given by the restriction of the holomorphic differential operator $d\pi_\mu(Y)$.

In turn, the action of $Y \in \mathfrak{g}$ on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ induced by the isomorphism (3.22) is given by

$$(\mathcal{F}_c \otimes \text{id}) \circ d\pi_\mu(Y) \circ (\mathcal{F}_c^{-1} \otimes \text{id}),$$

which is equal to $\widehat{d\pi_\mu}(Y)$ by Remark 3.6.

To complete the proof in the general case we denote by $\text{Hom}(P_{\mathbb{C}}, GL_{\mathbb{C}}(V))$ the set of holomorphic representations of $P_{\mathbb{C}}$ on V and by $\text{Hom}(\mathfrak{p}, \text{End}(V))$ the set of Lie algebra representations of \mathfrak{p} . Since the former is Zariski dense in the latter, the

two \mathfrak{g} -actions on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ coincide for all λ because both depend algebraically (actually affinely) on $\lambda \in \text{Hom}(\mathfrak{p}, \text{End}(V))$.

2) This statement is the analogue of Lemma 3.9 for the Lie group L . Indeed, since the group L normalizes $\mathfrak{n}_-(\mathbb{R})$ and fixes the origin 0, the isomorphism ι^* in (3.21) respects the L -action when L acts diagonally on $\mathcal{E}'_{\{0\}}(\mathfrak{n}_-(\mathbb{R})) \otimes V^\vee$. To conclude the proof we use (3.5). \square

The map F_c does not depend on the choice of a real form G of $G_{\mathbb{C}}$ that appears in the two middle terms of (3.23). Moreover, the isomorphism $F_c : \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \xrightarrow{\sim} \text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ depends only on the infinitesimal action of P on V . In fact, the following corollary follows immediately from the statement 1) of Proposition 3.11.

Corollary 3.12. *The algebraic Fourier transform of generalized Verma modules (see (3.23))*

$$F_c : \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \xrightarrow{\sim} \text{Pol}(\mathfrak{n}_+) \otimes V^\vee$$

is given by

$$(u \otimes v^\vee) \mapsto \widehat{d\pi_{\lambda^*}(u)}(1 \otimes v^\vee), \quad u \in U(\mathfrak{g}), v^\vee \in V^\vee.$$

4. F-METHOD

In Section 2 we have established a one-to-one correspondence between differential symmetry breaking operators for vector bundles and certain Lie algebra homomorphisms (Theorem 2.9). Using this framework our aim is to find explicit formulæ for such operators, in particular, when such operators are a priori known to be unique up to scalar. For this purpose we propose a new method, which we call the *F-method*. Its theoretical foundation is summarized in Theorem 4.1. This method becomes particularly simple when \mathfrak{h} is a parabolic subalgebra with abelian nilradical. In this case we develop the F-method in more details, and give its *recipe* in Section 4.4. Some useful lemmas for actual computations for vector-valued differential operators are collected in Section 4.5.

4.1. Construction of equivariant differential operators by algebraic Fourier transform. Let E be a finite-dimensional vector spaces over \mathbb{C} and E^\vee its dual space. Let $\text{Diff}^{\text{const}}(E)$ denote the ring of holomorphic differential operators on E with constant coefficients. We define the symbol map

$$\text{Symb} : \text{Diff}^{\text{const}}(E) \xrightarrow{\sim} \text{Pol}(E^\vee), \quad D_z \mapsto Q(\zeta)$$

by the following characterization

$$D_z e^{(z, \zeta)} = Q(\zeta) e^{(z, \zeta)}.$$

Then Symb is an algebra isomorphism. The differential operator on E with symbol $Q(\zeta)$ will be denoted by ∂Q_z .

By the definition of the algebraic Fourier transform (Definition 3.1) one has

$$(4.1) \quad \widehat{\partial P_z} = (-1)^\ell P(\zeta), \quad \widehat{Q(z)} = \partial Q_\zeta$$

for any homogeneous polynomial P on E^\vee of degree ℓ and any polynomial Q on E seen as a multiplication operator.

We recall from Corollary 3.12 that $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ is a (\mathfrak{g}, P) -module if V is a P -module. Note that the action of $\exp(\mathfrak{n}_+)$ ($\subset P$) on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ is not geometric, namely, it is not given by the pull-back of polynomials via the action on the base space \mathfrak{n}_+ .

The key tool for the F-method that we explain in Section 4.4 is the following assertion. We note that the two approaches (the canonical invariant pairing (2.20)) and the algebraic Fourier transform (3.23)) give rise to the same differential operators, provided that \mathfrak{n}_+ is abelian:

Theorem 4.1. *Suppose that \mathfrak{p} is a parabolic subalgebra \mathfrak{g} and that $P = L \exp(\mathfrak{n}_+)$ is its Levi decomposition. Let P' be a closed subgroup of P such that P' has a decomposition $P' = L' \exp(\mathfrak{n}'_+)$ with $L' \subset L$ and $\mathfrak{n}'_+ \subset \mathfrak{n}_+$. Let G' be an arbitrary subgroup of G containing P' . For a representation (λ, V) of P and a representation (ν, W) of P' , we form a G -equivariant vector bundle $\mathcal{V} = G \times_P V$ over $X = G/P$ and a G' -equivariant vector bundle $\mathcal{W} = G' \times_{P'} W$ over $Y = G'/P'$, respectively. Let $\mu := \lambda^*$ be as in (3.12).*

(1) *There is a natural isomorphism*

$$(4.2) \quad \begin{aligned} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) &\simeq (\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W))^{L', \widehat{d\pi}_\mu(\mathfrak{n}'_+)} \\ &\simeq (\text{Hom}_{L'}(V \otimes \text{Pol}(\mathfrak{n}_+), W))^{\widehat{d\pi}_\mu(\mathfrak{n}'_+)}. \end{aligned}$$

Here the right-hand side of (4.2) consists of $\psi \in \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W)$ satisfying

$$(4.3) \quad \nu(\ell) \circ \text{Ad}_\#(\ell)\psi \circ \lambda(\ell^{-1}) = \psi \quad \text{for all } \ell \in L',$$

$$(4.4) \quad (\widehat{d\pi}_\mu(C) \otimes \text{id}_W + \text{id} \otimes \nu(C))\psi = 0 \quad \text{for all } C \in \mathfrak{n}'_+.$$

(2) *Assume that the nilradical \mathfrak{n}_+ is abelian. Then the following diagram commutes:*

$$\begin{array}{ccc} \text{Hom}_{\mathbb{C}}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) & \xrightarrow{\mathcal{F}_c \otimes \text{id}} & \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W) & \xleftarrow{\text{Symb} \otimes \text{id}} & \text{Diff}^{\text{const}}(\mathfrak{n}_-) \otimes \text{Hom}_{\mathbb{C}}(V, W) \\ \cup & & \circlearrowright & & \cup \\ \text{Hom}_{P'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) & & \xrightarrow{D_{X \rightarrow Y}} & & \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y). \end{array}$$

Remark 4.2. The convention on the Fourier transform \mathcal{F}_c in Definition 3.5 makes the diagram in Theorem 4.1 commutative without additional powers of $\sqrt{-1}$.

Theorem 4.1 may be regarded as a construction of symmetry breaking operators by using the Fourier transform of generalized Verma modules.

Corollary 4.3. *Assume that \mathfrak{n}_+ is abelian and that $P' = L' \exp(\mathfrak{n}'_+)$ with $L' \subset L$ and $\mathfrak{n}'_+ \subset \mathfrak{n}_+$. Then the following diagram of three isomorphisms commutes.*

$$\begin{array}{ccc}
 & \text{Hom}_{L'}(V \otimes \text{Pol}(\mathfrak{n}_+), W)^{\widehat{d\pi_\mu}(\mathfrak{n}'_+)} & \\
 F_c \otimes \text{id} \nearrow & & \nwarrow \text{Symb} \otimes \text{id} \\
 \text{Hom}_{P'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{q}}(V^\vee)) & \xrightarrow[\sim]{D_{X \rightarrow Y}} & \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)
 \end{array}$$

In the above corollary, $\text{Hom}_{L'}(V \otimes \text{Pol}(\mathfrak{n}_+), W)^{\widehat{d\pi_\mu}(\mathfrak{n}'_+)}$ consists of L' -equivariant, $\text{Hom}_{\mathbb{C}}(V, W)$ -valued polynomial solutions ψ on \mathfrak{n}_+ to a system of partial differential equations of second order, see Sections 3.3 and 4.4. Corollary 4.3 implies that, once we find such a polynomial solution ψ , we obtain a P' -submodule W^\vee in $\text{ind}_{\mathfrak{p}}^{\mathfrak{q}}(V^\vee)$ (sometimes referred to as *singular vectors*) by $(F_c \otimes \text{id})^{-1}(\psi)$, and a differential symmetry breaking operator by $(\text{Symb} \otimes \text{id})^{-1}(\psi)$.

We first give proofs for the first statement of Theorem 4.1 here. The proof of the second statement is postponed until the next subsection.

Proof of Theorem 4.1 (1). Combining the duality theorem (Theorem 2.9) with the algebraic Fourier transform (Corollary 3.12) we have an isomorphism

$$\text{Hom}_{P'}(W^\vee, \text{Pol}(\mathfrak{n}_+) \otimes V^\vee) \xrightarrow{\sim} \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$$

where the P' -action on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ is defined via the algebraic Fourier transform F_c , namely, the left-hand side consists of $\psi \in \text{Hom}_{\mathbb{C}}(W^\vee, \text{Pol}(\mathfrak{n}_+) \otimes V^\vee) \simeq \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W)$ satisfying

$$(\widehat{d\pi_\mu}(C) \otimes \text{id}_W + \text{id} \otimes \nu(C))\psi = 0 \quad \text{for all } C \in \mathfrak{l}' + \mathfrak{n}_+,$$

provided L' is connected. Owing to Lemma 3.9, the condition for $C \in \mathfrak{l}'$ is equivalent to that $\psi \in (\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W))^{\mathfrak{l}'}$, where \mathfrak{l}' acts on $\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W)$ by $\text{ad}_{\#} \otimes \text{id} + \text{id} \otimes (\lambda^\vee \otimes \text{id} + \text{id} \otimes \nu)$.

In a more general setting where we allow L' to be disconnected, by the same argument as in the proof of Lemma 3.9, we see that the P' -action on $\text{Pol}(\mathfrak{n}_+) \otimes V^\vee$ via the algebraic Fourier transform F_c of generalized Verma modules (Corollary 3.12) coincides with the tensor product representation $\text{Ad}_{\#} \otimes \lambda^\vee$ when restricted to the Levi subgroup L . Thus the isomorphism (4.2) is proved. \square

4.2. Symbol map and reversing signatures. The purpose of this section is to carefully and clearly set up relations involving various signatures in connection with the algebraic Fourier transform in a coordinates-free fashion.

Denote by $\gamma : S(E) \xrightarrow{\sim} \text{Pol}(E^\vee)$ the canonical isomorphism, and define another algebra isomorphism

$$\gamma_{sgn} : S(E) \xrightarrow{\sim} \text{Pol}(E^\vee),$$

by $\gamma \circ a$, where $a : S(E) \rightarrow S(E)$ denotes the automorphism of the symmetric algebra $S(E)$ induced by the linear map $X \mapsto -X$ for $X \in E$.

Now we regard E as an abelian Lie algebra over \mathbb{C} , and identify its enveloping algebra $U(E)$ with the symmetric algebra $S(E)$. Then, the right and left-infinitesimal actions induce two isomorphisms:

$$dR : S(E) \xrightarrow{\sim} \text{Diff}^{\text{const}}(E), \quad dL : S(E) \xrightarrow{\sim} \text{Diff}^{\text{const}}(E).$$

By the definition of the symbol map, we get,

$$\text{Symb} \circ dR = \gamma, \quad \text{Symb} \circ dL = \gamma_{sgn}.$$

On the other hand, it follows from (4.1) that

$$\widehat{dL(u)} = \gamma(u), \quad \widehat{dR(u)} = \gamma_{sgn}(u),$$

for every $u \in S(E) \simeq U(E)$, where polynomials are regarded as multiplication operators. Hence we have proved

Lemma 4.4. *Let E be an abelian Lie algebra over \mathbb{C} . For any $u \in U(E)$,*

$$\text{Symb} \circ dR(u) = \widehat{dL(u)}, \quad \text{Symb} \circ dL(u) = \widehat{dR(u)}.$$

4.3. Proof of Theorem 4.1 (2). We are ready to complete the proof of Theorem 4.1 (and Corollary 4.3).

Proof. Take an arbitrary $\varphi \in \text{Hom}_{\mathbb{C}}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee))$, which may be written as a finite sum

$$\varphi = \sum_j u_j \otimes \psi_j \in U(\mathfrak{n}_-) \otimes \text{Hom}_{\mathbb{C}}(V, W)$$

by the Poincaré–Birkhoff–Witt theorem $U(\mathfrak{g}) \simeq U(\mathfrak{n}_-) \otimes U(\mathfrak{p})$. Then it follows from (2.22) and (3.23) that

$$F_c \varphi = \sum_j \mathcal{F}_c(dL(u_j)\delta) \otimes \psi_j \in \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W).$$

Since $\delta = \mathcal{F}_c^{-1}(\mathbf{1})$, we get

$$F_c \varphi = \sum_j \widehat{dL(u_j)} \otimes \psi_j.$$

On the other hand, by the construction (2.18),

$$D_{X \rightarrow Y}(\varphi) = \sum_j dR(u_j) \otimes \psi_j.$$

Now we use the assumption that \mathfrak{n}_+ or equivalently \mathfrak{n}_- is abelian. Then, in the coordinates $\mathfrak{n}_-(\mathbb{R}) \hookrightarrow G/P$ the operator $dR(u_j)$ for $u_j \in U(\mathfrak{n}_-)$ defines a constant

coefficient differential operator on \mathfrak{n}_- . Thus $D_{X \rightarrow Y}(\varphi)$ can be regarded as an element of $\text{Diff}^{\text{const}}(\mathfrak{n}_-(\mathbb{R})) \otimes \text{Hom}_{\mathbb{C}}(V, W)$.

Applying the symbol map we have

$$(\text{Symb} \otimes \text{id}) \circ D_{X \rightarrow Y}(\varphi) = \sum_j \text{Symb} \circ dR(u_j) \otimes \psi_j = \sum_j \widehat{dL}(u_j) \otimes \psi_j,$$

where the last equation follows from Lemma 4.4. Thus we have proved that

$$(F_c \otimes \text{id})\varphi = (\text{Symb} \otimes \text{id}) \circ D_{X \rightarrow Y}(\varphi),$$

whence the second statement of Theorem 4.1. \square

4.4. Recipe of the F-method for abelian nilradical \mathfrak{n}_+ . Our goal is to find an explicit form of a differential symmetry breaking operator from \mathcal{V}_X to \mathcal{W}_Y . Equivalently, what we call F-method provides a way to find an explicit element in the space $\text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) \simeq \text{Hom}_{\mathfrak{p}'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee))$.

A semisimple element Z in \mathfrak{g} is called *hyperbolic* if all the eigenvalues of $\text{ad}(Z)$ are real. A hyperbolic element Z defines a parabolic subalgebra $\mathfrak{p}(Z) = \mathfrak{l}(Z) + \mathfrak{n}_+(Z)$, where $\mathfrak{l}(Z)$ and $\mathfrak{n}_+(Z)$ are the sum of eigenspaces of $\text{ad}(Z)$ with zero and positive eigenvalues, respectively.

Let \mathfrak{g}' be a reductive subalgebra in \mathfrak{g} , in the sense that \mathfrak{g}' itself is reductive and the adjoint representation of \mathfrak{g}' on \mathfrak{g} is completely reducible.

Definition 4.5. A parabolic subalgebra \mathfrak{p} is said to be *\mathfrak{g}' -compatible* if there exists a hyperbolic element $Z \in \mathfrak{g}'$ such that $\mathfrak{p} = \mathfrak{p}(Z)$.

If $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$ is \mathfrak{g}' -compatible, then $\mathfrak{p}' := \mathfrak{p} \cap \mathfrak{g}'$ becomes a parabolic subalgebra of \mathfrak{g}' with the following Levi decomposition:

$$\mathfrak{p}' = \mathfrak{l}' + \mathfrak{n}'_+ := (\mathfrak{l} \cap \mathfrak{g}') + (\mathfrak{n}_+ \cap \mathfrak{g}'),$$

which satisfies the assumptions of Theorem 4.1 2).

In this case the space $\text{Diff}_{\mathfrak{g}'}(\mathcal{V}_X, \mathcal{W}_Y)$ of differential symmetry breaking operators is always finite-dimensional owing to Corollary 2.10 because:

$$\dim_{\mathbb{C}} \text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) < \infty$$

for any finite-dimensional representations V and W of \mathfrak{p} and \mathfrak{p}' , respectively [K14, Proposition 2.8].

Our assumption here is that $\mathfrak{p} = \mathfrak{l} + \mathfrak{n}_+$ is a \mathfrak{g}' -compatible parabolic subalgebra of \mathfrak{g} with abelian nilradical \mathfrak{n}_+ . Based on the following diagram (see Corollary 4.3),

$$(4.5) \quad \begin{array}{ccc} & (\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W))^{L', \widehat{d\pi}_\mu(\mathfrak{n}_+)} & \\ & \nearrow^{F_c \otimes \text{id}} \quad \quad \quad \nwarrow^{\text{Symb} \otimes \text{id}} & \\ \text{Hom}_{P'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) & \xrightarrow[\sim]{D_{X \rightarrow Y}} & \text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) \end{array}$$

we develop a method as follows:

- Step 0. Fix a finite-dimensional representation (λ, V) of the parabolic subgroup P . It defines a G -equivariant vector bundle $\mathcal{V}_X = G \times_P V$ over $X = G/P$.
- Step 1. Let $\mu := \lambda^\vee \otimes \mathbb{C}_{2\rho}$ and compute (see (3.14) and (3.15)),

$$\begin{aligned} d\pi_\mu : \mathfrak{g} &\rightarrow \mathcal{D}(\mathfrak{n}_-) \otimes \text{End}(V^\vee), \\ \widehat{d\pi}_\mu : \mathfrak{g} &\rightarrow \mathcal{D}(\mathfrak{n}_+) \otimes \text{End}(V^\vee). \end{aligned}$$

According to (3.13), $\widehat{d\pi}_\mu$ only depends on the infinitesimal representation λ of the parabolic subalgebra \mathfrak{p} .

- Step 2. Find a finite-dimensional representation (ν, W) of the Lie group P' such that

$$\text{Hom}_{P'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)) \neq \{0\}.$$

It defines a G' -equivariant vector bundle $\mathcal{W}_Y = G' \times_{P'} W$ over $Y = G'/P'$ such that $\text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$ is non-trivial.

- Step 3. Consider $\psi \in \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W)$ satisfying (4.3) and (4.4). Note that the system of partial differential equations (4.4) is of second order (see Proposition 3.10).
- Step 4. Take a slice S for generic L'_C -orbits on \mathfrak{n}_+ . Use invariant theory for (4.3) and consider the system of differential equations on S induced from (4.4). Find polynomials $\psi \in \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}(V, W)$ satisfying (4.3) and (4.4) by solving those equations on S .
- Step 5. Let ψ be a polynomial solution to (4.3) and (4.4) obtained in Step 4. In the diagram (4.5), $(\text{Symb} \otimes \text{id})^{-1}(\psi)$ gives the desired differential symmetry breaking operator in the coordinates \mathfrak{n}_- of X by Theorem 4.1. In the same diagram, $(F_c \otimes \text{id})^{-1}(\psi)$ gives an explicit element in $\text{Hom}_{P'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee))$ ($\simeq \text{Hom}_{\mathfrak{g}'}(\text{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee), \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee))$), which is sometimes referred to as a *singular vector*.

This method gives all non-trivial differential symmetry breaking operators for given data $(Y \leftrightarrow X, \mathcal{V}_X)$ by providing G' -equivariant vector bundles \mathcal{W}_Y and explicit elements in $\text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$. In fact, Step 2 based on Theorem 2.9 gives a necessary and sufficient condition for a P' -module W to ensure that $\text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$ is non-zero. Steps 1, 3 and 5 based on Theorem 4.1 show that any differential symmetry breaking operator is of the form $(F_c \otimes \text{id})^{-1}(\psi)$ where ψ is a polynomial solution to (4.3) and (4.4).

Actual applications of the F-method include the following cases:

1. Holomorphic discrete series representations.
2. Principal series representations of real reductive groups (Corollary 2.13).

The latter is related to questions in conformal geometry (more generally parabolic geometry), see [J09, KØSS13]. The former case includes the classical Rankin–Cohen bidifferential operators as a prototype, and it is the main object of the second part of this work [KP15-2]. The connection between these two is discussed in [KKP15]. We note that the normalization and formulation in [KØSS13] is different from this paper, see [K13].

Here we give some comments on the actual applications of the F-method when X and Y are Hermitian symmetric spaces. In Theorem 5.3 we prove that all continuous symmetry breaking operators in this case are given by holomorphic differential operators that extend to the complex flag varieties, so that the F-method for a parabolic subalgebra with abelian nilradical applies.

Furthermore, if (G, G') is a reductive symmetric pair, we know a priori that $\text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y)$ is one-dimensional for line bundles \mathcal{V}_X with generic parameter [K14, Theorem 2.7]. Thus, it is natural to look for explicit formulæ for such canonical operators. In Step 2 we can use explicit branching laws (see [KP15-2, Fact 4.2]) to find all W such that $\text{Hom}_{\mathfrak{p}'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee))$ is non-zero. Conversely, the differential equations in Step 3 are useful in certain cases to get a finer structure of branching laws, *e.g.*, to find the Jordan–Hölder series of the restriction for exceptional parameters λ (see [KØSS13]).

The Rankin–Cohen operators as well as Juhl’s conformally covariant differential operators are recovered by the F-method as a special case where generic L'_C -orbits on \mathfrak{n}_+ are of codimension one. The induced system of (4.4) reduces to ordinary differential equations on the one dimensional complex manifold S . In the second part of this work [KP15-2] we shall treat all the six geometries with a one-dimensional slice S .

4.5. F-method – supplement for vector valued cases. In order to deal with the general case where the target \mathcal{W}_Y is no longer a line bundle but a vector bundle, *i.e.*, where W is an arbitrary finite-dimensional irreducible \mathfrak{l}' -module, we may find the condition (4.3) somewhat complicated in practice, even though it is a system

of differential equations of first order. In this section we give two useful lemmas to simplify Step 3 in the recipe by reducing (4.3) to a simpler algebraic question on polynomial rings, so that we can focus on the crucial part consisting of a system of differential equations of second order (4.4). The results here will be used in [KP15-2, Sections 7 and 8].

We fix a Borel subalgebra $\mathfrak{b}(\mathfrak{l}')$ of \mathfrak{l}' . Let $\chi : \mathfrak{b}(\mathfrak{l}') \rightarrow \mathbb{C}$ be a character. For an \mathfrak{l}' -module U , we set

$$U_\chi := \{u \in U : Zu = \chi(Z)u \text{ for any } Z \in \mathfrak{b}(\mathfrak{l}')\}.$$

Suppose that W is an irreducible representation of \mathfrak{l}' with lowest weight $-\chi$. Then the contragredient representation W^\vee has a highest weight χ . We fix a non-zero highest weight vector $w^\vee \in (W^\vee)_\chi$. Then the contraction map

$$U \otimes W \rightarrow U, \quad \psi \mapsto \langle \psi, w^\vee \rangle,$$

induces a bijection between the following two subspaces:

$$(4.6) \quad (U \otimes W)^{\mathfrak{l}'} \xrightarrow{\sim} U_\chi,$$

if U is completely reducible as an \mathfrak{l}' -module. By using the isomorphism (4.6), we reformulate Step 3 of the recipe for the F-method as follows:

Lemma 4.6. *Suppose we are in the setting of Section 4.4. Assume that W is an irreducible representation of the parabolic subalgebra \mathfrak{p}' . Let $-\chi$ be the lowest weight of W as an \mathfrak{l}' -module. Then we have a natural injective homomorphism*

$$\text{Diff}_{G'}(\mathcal{V}_X, \mathcal{W}_Y) \hookrightarrow \{Q \in (\text{Pol}(\mathfrak{n}_+) \otimes V^\vee)_\chi : \widehat{d\pi}_\mu(C)Q = 0 \text{ for all } C \in \mathfrak{n}'_+\},$$

which is bijective if L' is connected.

Proof. Applying (4.6) to the \mathfrak{l}' -module $U := \text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}_{\mathbb{C}}(V, W)$, we get an isomorphism:

$$(4.7) \quad (\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}(V, W))^{\mathfrak{l}'} \xrightarrow{\sim} (\text{Pol}(\mathfrak{n}_+) \otimes V^\vee)_\chi.$$

Since W is an irreducible \mathfrak{p}' -module, the Lie subalgebra \mathfrak{n}' acts trivially on W and \mathfrak{l}' acts irreducibly. In particular, the condition (4.4) amounts to

$$(\widehat{d\pi}_\mu(C) \otimes \text{id}_W) \psi = 0 \text{ for all } C \in \mathfrak{n}'_+.$$

Therefore, the isomorphism (4.7) induces a bijection

$$\begin{aligned} & \left\{ \psi \in (\text{Pol}(\mathfrak{n}_+) \otimes \text{Hom}(V, W))^{\mathfrak{l}'} : \psi \text{ satisfies (4.4)} \right\} \\ & \xrightarrow{\sim} \left\{ Q \in (\text{Pol}(\mathfrak{n}_+) \otimes V^\vee)_\chi : \widehat{d\pi}_\mu(C)Q = 0 \text{ for all } C \in \mathfrak{n}'_+ \right\}. \end{aligned}$$

Now Lemma follows from Theorem 4.1. □

Since any non-zero vector in W^\vee is cyclic, the next lemma explains how to recover $D_{X \rightarrow Y}(\varphi)$ from Q given in Lemma 4.6.

We assume, for simplicity, that the \mathfrak{l} -module (λ, V) lifts to $L_{\mathbb{C}}$, the \mathfrak{l}' -module (ν, W) lifts to $L'_{\mathbb{C}}$, and use the same letters to denote their liftings.

Lemma 4.7. *For any $\varphi \in \text{Hom}_{\mathfrak{p}'}(W^\vee, \text{ind}_{\mathfrak{p}}^{\mathfrak{q}}(V^\vee))$, $\ell \in L'_{\mathbb{C}}$ and $w^\vee \in W^\vee$,*

$$(4.8) \quad \langle D_{X \rightarrow Y}(\varphi), \nu^\vee(\ell)w^\vee \rangle = (\text{Ad}(\ell) \otimes \lambda^\vee(\ell)) \langle D_{X \rightarrow Y}(\varphi), w^\vee \rangle.$$

Proof. We write $\varphi = \sum_j u_j \otimes \psi_j \in U(\mathfrak{n}_-) \otimes \text{Hom}_{\mathbb{C}}(V, W)$. Since φ is \mathfrak{p}' -invariant, we have the identity:

$$\sum_j u_j \otimes \psi_j = \sum_j \text{Ad}(\ell)u_j \otimes \nu(\ell) \circ \psi_j \circ \lambda(\ell^{-1}) \quad \text{for } \mathfrak{l} \in L'_{\mathbb{C}}.$$

In turn, we have

$$\begin{aligned} \langle D_{X \rightarrow Y}(\varphi), \nu^\vee(\ell)w^\vee \rangle &= \sum_j dR(\text{Ad}(\ell)u_j) \otimes \langle \psi_j, w^\vee \rangle \circ \lambda(\ell^{-1}) \\ &= ((\text{Ad}(\ell) \otimes \lambda^\vee(\ell)) \langle D_{X \rightarrow Y}(\varphi), w^\vee \rangle). \end{aligned}$$

Thus, we have proved Lemma. \square

We notice that the right-hand side of (4.8) can be computed by using the identity in $\text{Diff}^{\text{const}}(\mathfrak{n}_-) \otimes V^\vee$:

$$\langle D_{X \rightarrow Y}(\varphi), w^\vee \rangle = (\text{Symb}^{-1} \otimes \text{id}_{V^\vee})(Q),$$

once we know the polynomial $Q = \langle \psi, w^\vee \rangle$ with $\psi = (F_c \otimes \text{id})(\varphi)$ (see Theorem 4.1). In [KP15-2, Sections 7 and 8], we find explicit formulæ for vector-bundle valued equivariant differential operators by solving equations for the polynomials Q .

5. LOCALNESS AND EXTENSION THEOREM FOR SYMMETRY BREAKING OPERATORS

Let $G \supset G'$ be a pair of real reductive Lie groups. In general, continuous symmetry breaking operators between two principal series representations of G and G' are not always given by differential operators. Actually, generic ones are supposed to be given by integral transforms and their meromorphic continuation, as one can see from a classification result [KS14]. In this section, however, we formulate and prove a quite remarkable phenomenon (*localness theorem*) that any continuous G' -intertwining operator between two representation spaces consisting of holomorphic sections over Hermitian symmetric spaces is given by differential operators, see Theorem 5.3. In particular, the covariant holomorphic differential operators which we shall obtain explicitly in the second part [KP15-2] of this work exhaust all continuous symmetry breaking operators.

5.1. Formulation of the localness theorem. Let G be a connected reductive Lie group, θ a Cartan involution, and G/K the associated Riemannian symmetric space. We write $\mathfrak{c}(\mathfrak{k})$ for the center of the complexified Lie algebra $\mathfrak{k} := \text{Lie}(K) \otimes_{\mathbb{R}} \mathbb{C} \cong \mathfrak{k}(\mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$. In order to formulate a localness theorem, we suppose that G/K is a Hermitian symmetric space. This means that there exists a characteristic element $Z \in \mathfrak{c}(\mathfrak{k})$ such that the eigenvalues of $\text{ad}(Z) \in \text{End}(\mathfrak{g})$ is 0 or ± 1 and that we have an eigenspace decomposition

$$\mathfrak{g} = \mathfrak{k} + \mathfrak{n}_+ + \mathfrak{n}_-$$

of $\text{ad}(Z)$ with eigenvalues 0, 1, and -1 , respectively. We note that $\mathfrak{c}(\mathfrak{k})$ is one-dimensional if G is simple. With the notation of the previous sections, the complex Lie algebra \mathfrak{k} plays the role of the Levi subalgebra \mathfrak{l} .

Let $G_{\mathbb{C}}$ be a complex reductive Lie group with Lie algebra \mathfrak{g} , and $P_{\mathbb{C}}$ the maximal parabolic subgroup with Lie algebra $\mathfrak{p} := \mathfrak{k} + \mathfrak{n}_+$, with abelian nilradical \mathfrak{n}_+ . The complex structure of the homogeneous G/K is induced from the open embedding

$$G/K \subset G_{\mathbb{C}}/K_{\mathbb{C}} \exp \mathfrak{n}_+ = G_{\mathbb{C}}/P_{\mathbb{C}}.$$

Let G' be a connected reductive subgroup of G . Without loss of generality we may and do assume that G' is θ -stable. We set $K' := K \cap G'$. Our crucial assumption throughout this section is

$$(5.1) \quad Z \in \mathfrak{k}'.$$

Lemma 5.1. *If (5.1) holds, then the parabolic subalgebra \mathfrak{p} is \mathfrak{g}' -compatible (see Definition 4.5), and the homogeneous space G'/K' is a Hermitian sub-symmetric space of G/K such that the embedding $G'/K' \hookrightarrow G/K$ is holomorphic.*

Proof. Let $G'_{\mathbb{C}}$ be the connected complex subgroup of $G_{\mathbb{C}}$ with Lie algebra $\mathfrak{g}' := \text{Lie}(G') \otimes_{\mathbb{R}} \mathbb{C}$. Then $\mathfrak{p}' := \mathfrak{k}' + \mathfrak{n}'_+ \cong (\mathfrak{k} \cap \mathfrak{g}') + (\mathfrak{n}_+ \cap \mathfrak{g}')$ is the sum of the eigenspaces of $\text{ad}(Z)$ in \mathfrak{g}' with 0 and $+1$ eigenvalues, respectively, and therefore is a parabolic subalgebra of \mathfrak{g}' . We set $P'_{\mathbb{C}} := P_{\mathbb{C}} \cap G'_{\mathbb{C}}$. Then, the Riemannian symmetric space G'/K' becomes a Hermitian symmetric space, for which the complex structure is induced from the open embedding in the complex flag variety $Y_{\mathbb{C}} := G'_{\mathbb{C}}/P'_{\mathbb{C}}$:

$$\begin{array}{ccc} G'/K' & \hookrightarrow & G/K \\ \text{open } \cap & & \cap \text{ open} \\ Y_{\mathbb{C}} = G'_{\mathbb{C}}/P'_{\mathbb{C}} & \hookrightarrow & G_{\mathbb{C}}/P_{\mathbb{C}} = X_{\mathbb{C}}. \end{array}$$

Since $Y_{\mathbb{C}}$ is a complex submanifold of $X_{\mathbb{C}} = G_{\mathbb{C}}/P_{\mathbb{C}}$, the embedding $G'/K' \hookrightarrow G/K$ is holomorphic. \square

Notice that in the setting of Lemma 5.1 the complexified Lie algebra of K' is a Levi subalgebra of the parabolic subalgebra \mathfrak{p}' .

- Example 5.2.** (1) *Let G' be a connected simple Lie group such that the associated Riemannian symmetric space G'/K' is a Hermitian symmetric space. We take a characteristic element $Z' \in \mathfrak{c}(\mathfrak{k}')$. Let $G := G' \times G'$, and we realize G' as the diagonal subgroup $\Delta(G') := \{(g, g) : g \in G'\}$ of G . Then $Z := (Z', Z') \in \mathfrak{c}(\mathfrak{k})$ satisfies (5.1), yielding a holomorphic embedding $\Delta : G'/K' \hookrightarrow G/K = G'/K' \times G'/K'$.*
- (2) *Let G be a connected simple Lie group such that the associated symmetric space G/K is a Hermitian symmetric space with Z a characteristic element in $\mathfrak{c}(\mathfrak{k})$. Suppose τ is an automorphism of G of finite order such that $\tau(Z) = Z$. Let G' be the identity component of the subgroup $G^\tau := \{g \in G : \tau(g) = g\}$, and $K' := G' \cap K$. Then the assumption (5.1) is satisfied, and G'/K' is a Hermitian sub-symmetric space of G/K . We shall focus on the case where (G, G^τ) is a symmetric pair, namely, τ is of order two in [KP15-2] for detailed analysis.*

Consider a finite-dimensional representation of K on a complex vector space V . We extend it to a holomorphic representation of $P_{\mathbb{C}}$ by letting the unipotent subgroup $\exp(\mathfrak{n}_+)$ act trivially, and form a holomorphic vector bundle $\mathcal{V}_{X_{\mathbb{C}}} = G_{\mathbb{C}} \times_{P_{\mathbb{C}}} V$ over $X_{\mathbb{C}} = G_{\mathbb{C}}/P_{\mathbb{C}}$. The restriction to the open set G/K defines a G -equivariant holomorphic vector bundle $\mathcal{V} := G \times_K V$. We then have a natural representation of G on the vector space $\mathcal{O}(G/K, \mathcal{V})$ of global holomorphic sections endowed with the Fréchet topology of uniform convergence on compact sets.

Likewise, given a finite-dimensional representation W of K' , we form the G' -equivariant holomorphic vector bundle $\mathcal{W} = G' \times_{K'} W$ and consider the representation of G' on $\mathcal{O}(G'/K', \mathcal{W})$. By definition, it is clear that

$$(5.2) \quad \text{Diff}_{G'}^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y) \subset \text{Hom}_{G'}(\mathcal{O}(G/K, \mathcal{V}), \mathcal{O}(G'/K', \mathcal{W})).$$

Theorem 5.3 below shows that the two spaces do coincide.

Theorem 5.3. *Let G' be a reductive subgroup of G satisfying (5.1). Let V and W be any finite-dimensional representations of K and K' , respectively. Then,*

- (1) (localness theorem) *any continuous G' -homomorphism from $\mathcal{O}(G/K, \mathcal{V})$ to $\mathcal{O}(G'/K', \mathcal{W})$ is given by a holomorphic differential operator, in the sense of Definition 2.1, with respect to a holomorphic map between the Hermitian symmetric spaces $G'/K' \hookrightarrow G/K$, that is,*

$$\text{Diff}_{G'}^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y) = \text{Hom}_{G'}(\mathcal{O}(G/K, \mathcal{V}), \mathcal{O}(G'/K', \mathcal{W}));$$

- (2) (extension theorem) *any such a differential operator (or equivalently, any continuous G' -homomorphism) extends to a $G'_{\mathbb{C}}$ -equivariant holomorphic differential operator with respect to a holomorphic map between the flag varieties*

$$(5.3) \quad \begin{aligned} Y_{\mathbb{C}} = G'_{\mathbb{C}}/P'_{\mathbb{C}} &\hookrightarrow X_{\mathbb{C}} = G_{\mathbb{C}}/P_{\mathbb{C}}, \text{ namely, the injection} \\ \text{Diff}_{G'_{\mathbb{C}}}^{\text{hol}}(\mathcal{V}_{X_{\mathbb{C}}}, \mathcal{W}_{Y_{\mathbb{C}}}) &\hookrightarrow \text{Diff}_{G'}^{\text{hol}}(\mathcal{V}_X, \mathcal{W}_Y) \end{aligned}$$

is bijective.

Remark 5.4. More generally, we may ask whether an analogous statement to Theorem 5.3 (1) holds or not if we replace $\mathcal{O}(G/K, \mathcal{V})$ and $\mathcal{O}(G'/K', \mathcal{W})$ by some other topological vector spaces having the same underlying (\mathfrak{g}, K) -module and (\mathfrak{g}', K') -module, respectively (e.g. the Casselman–Wallach globalization, Hilbert space globalization, etc.). This question was raised by D. Vogan in May 2014. It turns out that this generalization is also true, as we shall show in the proof of Theorem 5.3, that the natural injection

$$(5.4) \quad \text{Diff}_{G'_{\mathbb{C}}}^{\text{hol}}(\mathcal{V}_{X_{\mathbb{C}}}, \mathcal{W}_{Y_{\mathbb{C}}}) \hookrightarrow \text{Hom}_{(\mathfrak{g}', K')}(\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}, \mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}})$$

is surjective if the assumption (5.1) is satisfied.

Remark 5.5. An analogous statement for real parabolic subgroups is not true. For instance, for the pair $(G, G') = (O(n+1, 1), O(n, 1))$ there always exists a non-zero continuous G' -equivariant map from the spherical principal series representations $C^\infty(G/P, \mathcal{L}_\lambda)$ of G to the one $C^\infty(G'/P', \mathcal{L}_\nu)$ of G' for any $(\lambda, \nu) \in \mathbb{C}^2$, however, non-zero G' -equivariant *differential operators* exist if and only if $\nu - \lambda \in 2\mathbb{N}$ [KS14].

Remark 5.6. Suppose that V is a generic character of K and (G, G') is a symmetric pair. Then owing to Theorems 2.12 and 5.3 (2), $\text{Diff}_{G'}^{\text{hol}}(\mathcal{V}_{X_{\mathbb{C}}}, \mathcal{W}_{Y_{\mathbb{C}}})$ is at most one-dimensional for any irreducible K' -module W , and [KP15-2, Fact 4.2] tells us precisely when it is non-zero.

In [KP15-2] we describe explicit formulæ of such differential operators by using the F-method (Theorem 4.1) for the six complex geometries arising from symmetric pairs of split rank one.

5.2. Proof of the localness theorem. Theorem 5.3 is a reflection of the theory of *discretely decomposable restrictions* (see [K94, K98a, K98b]). The proof is based on a careful analysis of the following three objects:

$$(\mathfrak{g}, K)\text{-modules, } (\mathfrak{g}, K')\text{-modules, and } (\mathfrak{g}', K')\text{-modules.}$$

We say that a K' -module Z is *K' -admissible* if the multiplicity

$$[M : F] := \dim \text{Hom}_{K'}(F, M|_{K'})$$

is finite for any $F \in \widehat{K'}$. Then, K' -admissibility is preserved by taking the tensor product with finite-dimensional representations.

We write $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}$ for the space of K -finite vectors of $\mathcal{O}(G/K, \mathcal{V})$, which becomes naturally a (\mathfrak{g}, K) -module.

Lemma 5.7. *The (\mathfrak{g}, K) -module $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}$ is K' -admissible if $Z \in \mathfrak{g}'$.*

Proof. As a K -module, we have the following isomorphism

$$\begin{aligned} \mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}} &\simeq S(\mathfrak{n}_+) \otimes V \\ &\simeq \left(\bigoplus_{a \geq 0} S^a(\mathfrak{n}_+) \right) \otimes V, \end{aligned}$$

where $S^a(\mathfrak{n}_+)$ denotes the space of symmetric tensors of homogeneous degree a . Since $\exp(\mathbb{R}\sqrt{-1}Z)$ acts on $S^a(\mathfrak{n}_+)$ as the scalar $e^{\sqrt{-1}at}$ ($t \in \mathbb{R}$), the whole $S(\mathfrak{n}_+)$ is admissible as a module of the one-dimensional subgroup $\exp(\mathbb{R}\sqrt{-1}Z)$, and so is $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}$. Hence it is also admissible as a K' -module by [K94, Theorem 1.2]. Alternatively, the lemma follows as a special case of the general result [K94, Theorem 2.7] or [K98a, Theorem 4.1]. \square

Given a (\mathfrak{g}, K') -module M , we consider the contragredient representation on the dual space $M^\vee := \text{Hom}_{\mathbb{C}}(M, \mathbb{C})$. Collecting K' -finite vectors in M^\vee , we get a (\mathfrak{g}, K') -module $(M^\vee)_{K'\text{-finite}}$.

Lemma 5.8. *Let M be a K' -admissible (\mathfrak{g}, K') -module. Then,*

- (1) *M is discretely decomposable as a (\mathfrak{g}', K') -module.*
- (2) *The (\mathfrak{g}, K') -module $(M^\vee)_{K'\text{-finite}}$ is K' -admissible and one has the following K' -isomorphism*

$$(M^\vee)_{K'\text{-finite}} \simeq \bigoplus_{F \in \widehat{K'}} [M : F] F^\vee.$$

For the proof we refer to [K98b, Proposition 1.6].

Lemma 5.9. *Let M be a K' -admissible (\mathfrak{g}, K) -module. Then,*

$$(M^\vee)_{K\text{-finite}} = (M^\vee)_{K'\text{-finite}}.$$

Proof. There is an obvious inclusion $(M^\vee)_{K\text{-finite}} \subset (M^\vee)_{K'\text{-finite}}$. We shall prove that the multiplicities in $(M^\vee)_{K\text{-finite}}$ and $(M^\vee)_{K'\text{-finite}}$ are both finite and are the same. Indeed, M being K' -admissible, one has

$$[M : F] = \bigoplus_{E \in \widehat{K}} [M : E][E : F] < \infty.$$

Conversely, $(M^\vee)_{K'\text{-finite}} \simeq \bigoplus_{F \in \widehat{K'}} [M : F] F^\vee$ and thus,

$$(M^\vee)_{K\text{-finite}} \simeq \bigoplus_{E \in \widehat{K}} [M : E] E^\vee \simeq \bigoplus_{F \in \widehat{K'}} \left(\bigoplus_{E \in \widehat{K}} [M : E][E : F] \right) F^\vee,$$

which concludes the proof. \square

The next lemma is known to experts, but for the sake of completeness, we give a proof.

Lemma 5.10. *There is a natural (\mathfrak{g}, K) -isomorphism:*

$$\left((\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}})^\vee \right)_{K\text{-finite}} \simeq \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee).$$

Proof. As in Lemma 2.15, there is a natural non-degenerate \mathfrak{g} -invariant bilinear form

$$\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}} \times \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \rightarrow \mathbb{C}.$$

Hence, we have an injective (\mathfrak{g}, K) -homomorphism $\text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \subset (\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}})^\vee$. Taking K -finite vectors we get the following commutative diagram of K -modules isomorphisms:

$$\begin{array}{ccc} \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) & \subset & \left((\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}})^\vee \right)_{K\text{-finite}} \\ \Big\| & & \Big\| \\ S(\mathfrak{n}_-) \otimes V^\vee & \simeq & ((\text{Pol}(\mathfrak{n}_-) \otimes V)^\vee)_{K\text{-finite}}. \end{array}$$

Hence the first row is also bijective. □

Combining Lemmas 5.7, 5.9 and 5.10 we have shown the following key result:

Proposition 5.11. *There is a natural (\mathfrak{g}, K) -isomorphism:*

$$\left((\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}})^\vee \right)_{K'\text{-finite}} \simeq \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee).$$

Proof of Theorem 5.3. Let $T : \mathcal{O}(G/K, \mathcal{V}) \rightarrow \mathcal{O}(G'/K', \mathcal{W})$ be a continuous G' -intertwining operator. It induces a (\mathfrak{g}', K') -homomorphism

$$(5.5) \quad T_K : \mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}} \rightarrow \mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}}.$$

We shall prove that any such (\mathfrak{g}', K') -homomorphism T_K comes from a $G'_\mathbb{C}$ -equivariant differential operator on the flag variety.

To see this, we take the dual map (5.5), and apply Lemma 5.10 and Proposition 5.11. Then there is a (\mathfrak{g}', K') -homomorphism $\psi : \text{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee) \rightarrow \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)$ such that the following diagram commutes:

$$\begin{array}{ccc} \left(\mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}}^\vee \right)_{K'\text{-finite}} & \xrightarrow{T_K^\vee} & \left(\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}^\vee \right)_{K'\text{-finite}} \\ \Big\| \text{Lemma 5.10} & & \Big\| \text{Proposition 5.11} \\ \text{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee) & \xrightarrow{\psi} & \text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee) \end{array}$$

The correspondence $T_K \mapsto \psi$ is one-to-one, and thus we have obtained the following natural injective map

$$\mathrm{Hom}_{(\mathfrak{g}', K')}(\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}, \mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}}) \hookrightarrow \mathrm{Hom}_{\mathfrak{g}'}(\mathrm{ind}_{\mathfrak{p}'}^{\mathfrak{g}'}(W^\vee), \mathrm{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)).$$

According to Theorems 2.12 and 4.1 the latter space is isomorphic to $\mathrm{Diff}_{G'}^{\mathrm{hol}}(\mathcal{V}_X, \mathcal{W}_Y)$. This shows that (5.4) is surjective.

Since the injective map (5.4) factors the two injective maps (5.2) and (5.3), both (5.2) and (5.3) are bijective. \square

5.3. Automatic continuity theorem in the unitary case. Any unitary highest weight module is realized as a subrepresentation of $\mathcal{O}(G/K, \mathcal{V})$ for some G -equivariant holomorphic vector bundle \mathcal{V} over G/K . In this subsection, we prove that any continuous homomorphism between Fréchet modules $\mathcal{O}(G/K, \mathcal{V})$ and $\mathcal{O}(G'/K', \mathcal{W})$ induces a continuous homomorphism between their unitary submodules.

Definition 5.12. For a Fréchet G -module \mathcal{F} , we say a G -submodule \mathcal{H} is a *unitary submodule* if \mathcal{H} is a Hilbert space such that the inclusion map $\mathcal{H} \hookrightarrow \mathcal{F}$ is continuous and that G acts unitarily on \mathcal{H} .

If V is an irreducible K -module, then there exists at most one non-zero unitary submodule (up to a scaling of the inner product) of $\mathcal{O}(G/K, \mathcal{V})$. We denote by \mathcal{H}_V^G the unitary submodule of $\mathcal{O}(G/K, \mathcal{V})$. The classification of irreducible K -modules V for which $\mathcal{H}_V^G \neq \{0\}$ was accomplished in [EHW83]. We shall prove that any G' -equivariant differential operator in Theorem 5.3 preserves the unitary submodules in the following sense:

Theorem 5.13. *Let G' be a reductive subgroup of G satisfying (5.1). Let V and W be any irreducible finite-dimensional representations of K and K' , respectively. Suppose that $T : \mathcal{O}(G/K, \mathcal{V}) \rightarrow \mathcal{O}(G'/K', \mathcal{W})$ is a G' -equivariant differential operator such that $T|_{\mathcal{H}_V^G} \neq 0$. Then $\mathcal{H}_W^{G'} \neq \{0\}$ and T induces a continuous G' -equivariant linear map from the Hilbert space \mathcal{H}_V^G onto the Hilbert space $\mathcal{H}_W^{G'}$.*

Applying Theorems 5.3 and 5.13 to the setting of Example 5.2 (1), we have:

Example 5.14. *Any symmetry breaking operator for the tensor product of two holomorphic discrete series representations is given by a holomorphic differential operator if those representations are realized in the space of holomorphic sections for G -equivariant holomorphic vector bundles over the Hermitian symmetric space G/K . The Rankin–Cohen bidifferential operators are such operators for $G = SL(2, \mathbb{R})$.*

Remark 5.15. As we shall see in the proof, the unitary representation \mathcal{H}_V^G decomposes discretely when restricted to G' if the condition (5.1) is satisfied. The unitary

submodule $\mathcal{H}_W^{G'}$ occurs as a discrete summand of the restriction of the unitary representation \mathcal{H}_V^G of G to the subgroup G' .

Let V be an irreducible representation of K as before. Then, there exists a unique K -submodule of $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}} \simeq S(\mathfrak{n}_+) \otimes V$ isomorphic to V , namely $S^0(\mathfrak{n}_+) \otimes V \simeq V$.

Lemma 5.16. *Let M be a non-zero (\mathfrak{g}, K) -submodule of $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}}$. Then,*

- 1) *The module M contains V .*
- 2) *If M is unitarizable, then its Hilbert completion can be realized in $\mathcal{O}(G/K, \mathcal{V})$ and $M = (\mathcal{H}_V^G)_{K\text{-finite}}$.*

Proof. 1) Since any non-zero quotient of the (generalized) Verma module $\text{ind}_{\mathfrak{p}}^{\mathfrak{g}}(V^\vee)$ contains V^\vee , the first statement follows from Lemma 5.10. Alternatively, since the infinitesimal action of \mathfrak{n}_- on $\mathcal{O}(G/K, \mathcal{V})_{K\text{-finite}} \simeq \text{Pol}(\mathfrak{n}_-) \otimes V$ is given by directional derivatives, iterated operators of \mathfrak{n}_- yield non-zero elements in V .

2) Denote by (π, \overline{M}) the unitary representation of G obtained as an (abstract) Hilbert completion of the (\mathfrak{g}, K) -module M . We regard V as a K -submodule of M , and also of \overline{M} . Then the map

$$G \times \overline{M} \times V \longrightarrow \mathbb{C}, \quad (g, w, v) \mapsto (w, \pi(g)v)_{\overline{M}},$$

induces an injective G -homomorphism $\iota : \overline{M} \longrightarrow \mathcal{O}(G/K, \mathcal{V})$. Since \mathcal{H}_V^G is the unique non-zero unitary submodule, ι is an isomorphism onto \mathcal{H}_V^G . \square

Proof of Theorem 5.13. By Lemma 5.7, the module $(\mathcal{H}_V^G)_{K\text{-finite}}$ is K' -admissible. Therefore, the unitary representation \mathcal{H}_V^G decomposes into a Hilbert direct sum of irreducible unitary representations $\{U_j\}$ of G' :

$$(5.6) \quad \mathcal{H}_V^G \simeq \sum_j^{\oplus} m_j U_j,$$

with $m_j < \infty$ for all j ([K94, Theorem 1.1]) and the underlying (\mathfrak{g}, K) -module $(\mathcal{H}_V^G)_{K\text{-finite}}$ is isomorphic to an algebraic direct sum of irreducible and unitarizable (\mathfrak{g}', K') -modules

$$(5.7) \quad (\mathcal{H}_V^G)_{K\text{-finite}} = (\mathcal{H}_V^G)_{K'\text{-finite}} \simeq \bigoplus_j m_j (U_j)_{K'\text{-finite}},$$

with the same multiplicities [K98b]. (We remark that an analogous statement fails for the restriction $\pi|_{G'}$ of an irreducible unitary representation π of G if the branching law contains continuous spectrum).

As we saw in the proof of Theorem 5.3, the G' -equivariant differential operators T induces a (\mathfrak{g}', K') -homomorphism

$$T_K : (\mathcal{H}_V^G)_{K\text{-finite}} \longrightarrow \mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}}.$$

By (5.7), $M := T_K((\mathcal{H}_V^G)_{K\text{-finite}})$ is an algebraic direct sum of some irreducible unitarizable (\mathfrak{g}', K') -modules. Since $\mathcal{O}(G'/K', \mathcal{W})_{K'\text{-finite}}$ contains at most one irreducible unitarizable (\mathfrak{g}', K') -module, M is irreducible as a (\mathfrak{g}', K') -module, and we can realize its Hilbert completion as $\mathcal{H}_W^{G'}$ by Lemma 5.16.

In view of (5.6) and (5.7), there exists a continuous G' -homomorphism between Hilbert spaces:

$$\tilde{T} : \mathcal{H}_V^G \longrightarrow \mathcal{H}_W^{G'}$$

such that $\tilde{T}|_{(\mathcal{H}_V^G)_{K\text{-finite}}} = T|_{(\mathcal{H}_V^G)_{K\text{-finite}}}$. Since the inclusion map $\mathcal{H}_V^G \hookrightarrow \mathcal{O}(G/K, \mathcal{V})$ and the differential operator $T : \mathcal{O}(G/K, \mathcal{V}) \longrightarrow \mathcal{O}(G'/K', \mathcal{W})$ are both continuous, we get $\tilde{T} = T$ on \mathcal{H}_V^G . Hence Theorem is proved. \square

5.4. Orthogonal projectors. If V is one-dimensional and (G, G') is a reductive symmetric pair satisfying (5.1), then all the multiplicities m_j in (5.6) are equal to one (see [K08]) and it becomes meaningful to describe the projector from \mathcal{H}_V^G to each G' -irreducible summand. We explain briefly the relationship between the projector for the unitary representation and the symmetry breaking operator.

For this, suppose $T : \mathcal{O}(G/K, \mathcal{V}) \rightarrow \mathcal{O}(G'/K', \mathcal{W})$ is a G' -equivariant differential operator such that $T|_{\mathcal{H}_V^G} \neq 0$. By Theorem 5.13, T induces a continuous map $T : \mathcal{H}_V^G \rightarrow \mathcal{H}_W^{G'}$. Let $T^* : \mathcal{H}_W^{G'} \rightarrow \mathcal{H}_V^G$ be its the adjoint operator. Then the composition $T^*T : \mathcal{H}_V^G \rightarrow \mathcal{H}_V^G$ is a G' -intertwining operator onto the G' -irreducible summand which is isomorphic to $\mathcal{H}_W^{G'}$. Since T vanishes on the orthogonal complement to $T^*(\mathcal{H}_W^{G'})$, it is (up to scaling) the orthogonal projector onto $\mathcal{H}_W^{G'}$.

Explicit description of such differential operators T will be the main concern of the second part [KP15-2] of this work.

Acknowledgements. T. Kobayashi was partially supported by Institut des Hautes Études Scientifiques, France and Grant-in-Aid for Scientific Research (B) (22340026) and (A) (25247006), Japan Society for the Promotion of Science. Both authors were partially supported by Max Planck Institute for Mathematics (Bonn) where a large part of this work was done.

REFERENCES

- [DP07] G. van Dijk, M. Pevzner, Ring structures for holomorphic discrete series and Rankin–Cohen brackets. *J. Lie Theory*, **17**, (2007), pp. 283–305.
- [EHW83] Th. Enright, R. Howe, N. Wallach, A classification of unitary highest weight modules. *Representation theory of reductive groups* (Park City, Utah, 1982), pp. 97–143, Progress in Mathematics, **40**, Birkhäuser, Boston, MA, 1983.
- [HJ82] M. Harris, H. P. Jakobsen, Singular holomorphic representations and singular modular forms. *Math. Ann.* **259** (1982), pp.227–244.

- [J09] A. Juhl, *Families of conformally covariant differential operators, Q-curvature and holography*. Progress in Mathematics, **275**. Birkhäuser, Basel, 2009.
- [KnSt71] A. W. Knap, E. M. Stein, Intertwining operators for semisimple groups. *Ann. of Math.* **93** (1971), pp. 489–578.
- [K94] T. Kobayashi, Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups and its applications, *Invent. Math.* **117** (1994), pp. 181–205.
- [K98a] T. Kobayashi, Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups II—micro-local analysis and asymptotic K -support, *Ann. of Math.* (2) **147** (1998), pp. 709–729.
- [K98b] T. Kobayashi, Discrete decomposability of the restriction of $A_q(\lambda)$ with respect to reductive subgroups III—restriction of Harish-Chandra modules and associated varieties, *Invent. Math.* **131** (1998), pp. 229–256.
- [K08] T. Kobayashi, Multiplicity-free theorems of the restrictions of unitary highest weight modules with respect to reductive symmetric pairs. Representation theory and automorphic forms, Progr. Math., **255**, Birkhäuser, Boston, (2008), pp. 45–109.
- [K12] T. Kobayashi, Restrictions of generalized Verma modules to symmetric pairs, *Transformation Groups* **17**, (2012), pp. 523–546.
- [K13] T. Kobayashi. F-method for constructing equivariant differential operators. *Contemporary Mathematics*, **598** (2013), pp. 141–148. Amer. Math. Soc.
- [K14] T. Kobayashi, F-method for symmetry breaking operators, *Diff. Geometry and its Appl.* **33**, (2014), pp. 272–289.
- [KØSS13] T. Kobayashi, B. Ørsted, P. Somberg, V. Souček, Branching laws for Verma modules and applications in parabolic geometry. Part I. Preprint. ArXiv:1305.6040.
- [KKP15] T. Kobayashi, T. Kubo, M. Pevzner, Vector-valued covariant differential operators for the Möbius transformation. In Lie theory and its applications in Physics. V. Dobrev (Ed.) *Springer Proceedings in Mathematics and Statistics*. **111** (2015) pp. 67–86.
- [KP15-2] T. Kobayashi, M. Pevzner, Differential symmetry breaking operators. II. Rankin–Cohen operators for symmetric pairs, to appear in *Selecta Mathematica* (the same issue), arXiv:1301.2111.
- [KS14] T. Kobayashi, B. Speh, Intertwining operators and the restriction of representations of rank one orthogonal groups, *C. R. Acad. Sci. Paris, Sér. I.* **352**, (2014), pp. 89–94.
- [Kos74] B. Kostant, Verma modules and the existence of quasi-invariant differential operators, *Lecture Notes in Math.* **466**, Springer-Verlag, (1974), pp. 101–129.
- [Pee59] J. Peetre, Une caractérisation abstraite des opérateurs différentiels, *Math. Scand.* **7** (1959), pp. 211–218.
- [S66] L. Schwartz, *Théorie des distributions*. Hermann, Paris, 1966.

T. Kobayashi. Kavli IPMU and Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo, 153-8914 Japan; toshi@ms.u-tokyo.ac.jp.

M. Pevzner. Laboratoire de Mathématiques de Reims, Université de Reims-Champagne-Ardenne, FR 3399 CNRS, F-51687, Reims, France; pevzner@univ-reims.fr.