

## Multiplicity in restricting small representations

By Toshiyuki KOBAYASHI

Graduate School of Mathematical Sciences and Kavli IPMU (WPI), The University of Tokyo.

**Abstract:** We give a geometric criterion for the bounded multiplicity property of “small” infinite-dimensional representations of real reductive Lie groups in both induction and restrictions.

In particular, for a reductive symmetric pair  $(G, H)$ , we determine the reductive subgroups  $G'$  having the property that any irreducible  $H$ -distinguished admissible representations of  $G$  are of bounded multiplicity when restricted to  $G'$ .

**Key words:** branching law; multiplicity; reductive group; symmetric pair; visible action; spherical variety.

**1. Introduction** By branching problems in representation theory, we mean the broad problem of understanding how irreducible representations of a group behave when restricted to a subgroup. As viewed in [12], we may divide the branching problems into the following three stages:

**Stage A.** Abstract features of the restriction;

**Stage B.** Branching law;

**Stage C.** Construction of symmetry breaking operators.

The role of Stage A is to develop a theory on the restriction of representations as generally as possible. In turn, we may expect a detailed study of the restriction in Stages B (decomposition of representations) and C (decomposition of vectors) in the “promising” settings that are suggested by the general theory in Stage A.

This article concerns a question in Stage A about “multiplicity” in branching problems.

Let  $G$  be a real reductive Lie group,  $\mathcal{M}(G)$  the category of finitely generated, smooth admissible representations of  $G$  of moderate growth [31, Chap. 11], and  $\text{Irr}(G)$  the set of irreducible objects in  $\mathcal{M}(G)$ . We shall use the uppercase letter  $\Pi$  for representations of the group  $G$ , and the lowercase letter  $\pi$  for those of a reductive subgroup  $G'$ .

For Stage A, we may formulate an abstract feature of the restrictions as a property for

- the pair  $(G, G')$ ,
- the triple  $(G, G', \Pi)$ , or
- the quadruple  $(G, G', \Pi, \pi)$ .

The formulation for *the triple*  $(G, G', \Pi)$  was adopted in the study of  *$G'$ -admissible restriction* of  $\Pi$ , namely, the restriction  $\Pi|_{G'}$  of  $\Pi \in \text{Irr}(G)$  being discretely decomposable with finite multiplicity, see [5, 6, 7] for the general theory, and [17] for some classification theory of the triples  $(G, G', \Pi)$ .

On the other hand, Fact 2.1 below is formulated as a property for *the pair*  $(G, G')$ . This is the study of “multiplicity” of the restrictions, see [11, 16] for the general theory, and [15] for the classification of the pairs  $(G, G')$ . In this article, we discuss its refinement in a *formulation for the triple*  $(G, G', \Omega)$  or for *the quadruple*  $(G, G', \Omega, \Omega')$  where  $\Omega \subset \mathcal{M}(G)$  and  $\Omega' \subset \mathcal{M}(G')$  are families of “small” infinite-dimensional representations, see Problems 2.3 and 4.1. This refinement reveals the underlying geometric structures of some concrete examples, *e.g.*, [1, 13, 24], and yields much broader settings that seem to be promising for analysis of branching problems in Stage C.

Detail proofs of the theorems in this article will appear in [14].

### 2. Bounded multiplicity in restriction

Throughout this article, we shall assume that  $G \supset G'$  are real forms of complex reductive algebraic Lie groups  $G_{\mathbb{C}} \supset G'_{\mathbb{C}}$ , respectively. Their compact real forms will be denoted by  $G_U \supset G'_U$ . The Lie algebras will be denoted by the corresponding lowercase German letters  $\mathfrak{g}, \mathfrak{g}_{\mathbb{C}}, \mathfrak{g}_U, \mathfrak{g}'$ , etc.

For  $\Pi \in \mathcal{M}(G)$  and  $\pi \in \mathcal{M}(G')$ , we define the **multiplicity** of the restriction  $\Pi|_{G'}$  in the category  $\mathcal{M}$  by

$$[\Pi|_{G'} : \pi] := \dim_{\mathbb{C}} \text{Hom}_{G'}(\Pi|_{G'}, \pi) \in \mathbb{N} \cup \{\infty\},$$

where  $\text{Hom}_{G'}(\cdot, \cdot)$  denotes the space of continuous  $G'$ -homomorphisms between the Fréchet representations.

In [16, Thms. C and D] we proved the following geometric criteria:

**Fact 2.1.** Let  $G \supset G'$  be a pair of algebraic real reductive Lie groups.

(1) **Bounded multiplicity** for a pair  $(G, G')$ :

$$(2.1) \quad \sup_{\Pi \in \text{Irr}(G)} \sup_{\pi \in \text{Irr}(G')} [\Pi|_{G'} : \pi] < \infty$$

if and only if  $(G_{\mathbb{C}} \times G'_{\mathbb{C}})/\text{diag } G'_{\mathbb{C}}$  is spherical.

(2) **Finite multiplicity** for a pair  $(G, G')$ :

$$(2.2) \quad [\Pi|_{G'} : \pi] < \infty, \quad \forall \Pi \in \text{Irr}(G), \forall \pi \in \text{Irr}(G')$$

if and only if  $(G \times G')/\text{diag } G'$  is real spherical.

Here we recall that a complex  $G_{\mathbb{C}}$ -manifold  $X$  is called *spherical* if a Borel subgroup of  $G_{\mathbb{C}}$  has an open orbit in  $X$ , and that a  $G$ -manifold  $Y$  is called *real spherical* if a minimal parabolic subgroup of  $G$  has an open orbit in  $Y$ .

A remarkable feature of Fact 2.1 (1) is that the bounded multiplicity property (2.1) is determined only by the complexifications of  $G$  and  $G'$ , hence the classification of such pairs  $(G, G')$  is reduced to a classical result [20]: the pair  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  is the direct sum of the following ones up to abelian ideals:

$$(2.3) \quad (\mathfrak{sl}_n, \mathfrak{gl}_{n-1}), (\mathfrak{so}_n, \mathfrak{so}_{n-1}), \text{ or } (\mathfrak{so}_8, \mathfrak{spin}_7).$$

On the other hand, the finite multiplicity property (2.2) depends on real forms. It is fulfilled for any Riemannian symmetric pair by Harish-Chandra's admissibility theorem, whereas it is not the case for some reductive symmetric pairs such as  $(G, G') = (SL(p+q, \mathbb{R}), SO(p, q))$ . A complete classification of the symmetric pairs  $(G, G')$  satisfying the finite multiplicity property (2.2) was accomplished in [15].

**Example 2.2.** Let  $p_1 + p_2 = p$ ,  $q_1 + q_2 = q$ , and  $(G, G') = (O(p, q), O(p_1, q_1) \times O(p_2, q_2))$ . Suppose  $p + q \geq 5$ . The criteria in Fact 2.1 give the equivalences:

$$(2.1) \iff p_1 + q_1 = 1 \text{ or } p_2 + q_2 = 1.$$

$$(2.2) \iff p_1 + q_1 = 1, p_2 + q_2 = 1, p = 1, \text{ or } q = 1.$$

This means that for general  $p_1, q_1, p_2, q_2$ , there exist  $\Pi \in \text{Irr}(G)$  and  $\pi \in \text{Irr}(G')$  such that  $[\Pi|_{G'} : \pi] = \infty$ . Nevertheless, a multiplicity-free theorem holds for the restriction  $\Pi|_{G'}$  for any  $p_1, p_2, q_1, q_2$ , and for any discrete series representation  $\Pi$  for the symmetric space  $G/H$  with  $H = O(p-1, q)$ , see [13] for a

precise statement.

This example suggests us to work with the *triple*  $(G, G', \Pi)$  rather than the *pair*  $(G, G')$  for the finer study of multiplicity estimates as mentioned in Introduction.

Take  $\Pi \in \mathcal{M}(G)$ . We say the restriction  $\Pi|_{G'}$  has the *finite multiplicity property* if  $[\Pi|_{G'} : \pi] < \infty$  for all  $\pi \in \text{Irr}(G')$ , and has the *bounded multiplicity property* if  $m(\Pi|_{G'}) < \infty$ , where we set

$$(2.4) \quad m(\Pi|_{G'}) := \sup_{\pi \in \text{Irr}(G')} [\Pi|_{G'} : \pi] \in \mathbb{N} \cup \{\infty\}.$$

In search for broader settings in which we could expect a detailed study of the restriction  $\Pi|_{G'}$  in Stages B and C, we address the following:

**Problem 2.3.** Given a pair  $G \supset G'$ , find a subset  $\Omega$  of  $\mathcal{M}(G)$  such that  $\sup_{\Pi \in \Omega} m(\Pi|_{G'}) < \infty$ .

We bear in mind that branching problems often arise for a family of representations  $\Pi$ . For a better understanding of Problem 2.3, we first examine two opposite extremal choices of  $\Omega$ . When  $\Omega$  is a singleton, Problem 2.3 concerns the triple  $(G, G', \Pi)$  having the bounded multiplicity property. When  $\Omega$  is the whole set  $\text{Irr}(G)$ , Problem 2.3 asks the condition (2.1), and is solved by the geometric criterion for the pair  $(G, G')$ , as seen in Fact 2.1 (1). Second, we note that Problem 2.3 is nontrivial even when  $G$  is a compact Lie group where  $m(\Pi|_{G'})$  is individually finite. In this article we discuss Problem 2.3 with focus on the following two cases:

- (1)  $\Omega = \text{Irr}(G)_H$ , the set of  $H$ -distinguished irreducible representations of  $G$  (Theorem 3.2);
- (2)  $\Omega = \Omega_P, \Omega_{P, q}$ : families of degenerate principal series representations (Theorems 4.2 and 4.3).

**Remark 2.4.** One may wonder why we did not use  $[\pi : \Pi|_{G'}] := \dim_{\mathbb{C}} \text{Hom}_{G'}(\pi, \Pi|_{G'})$  instead of  $[\Pi|_{G'} : \pi]$ . The reason is that the space  $\text{Hom}_{G'}(\pi, \Pi|_{G'})$  may be too small to capture the whole picture of the restriction  $\Pi|_{G'}$  in the category  $\mathcal{M}$ . This feature is akin to the fact in the category of Harish-Chandra modules that  $\text{Hom}_{\mathfrak{g}', K'}(\pi_{K'}, \Pi_K|_{\mathfrak{g}'})$  vanishes unless  $\Pi_K$  is “discretely decomposable” as a  $(\mathfrak{g}', K')$ -module [7].

**3.  $H$ -distinguished representations of  $G$**   
For  $\Pi \in \text{Irr}(G)$ , we denote by  $\Pi^{-\infty}$  the representation on the space of distribution vectors, that is, the topological dual of  $\Pi$ . For a closed subgroup  $H$  of  $G$ , we set

$$(3.1) \quad \text{Irr}(G)_H := \{\Pi \in \text{Irr}(G) : (\Pi^{-\infty})^H \neq \{0\}\}.$$

The Frobenius reciprocity tells  $\Pi \in \text{Irr}(G)_H$  if and only if  $\text{Hom}_G(\Pi^\vee, C^\infty(G/H)) \neq \{0\}$ , where  $\Pi^\vee$  is the contragredient representation in the category  $\mathcal{M}(G)$ . Elements  $\Pi$  in  $\text{Irr}(G)_H$  (or  $\Pi^\vee$ ) are sometimes referred to as *H-distinguished*, or having *nonzero H-periods*.

For a reductive symmetric pair  $(G, H)$ , the set  $\text{Irr}(G)_H$  is described by the Cartan–Helgason theorem when  $H$  is compact, whereas the full classification is far from being achieved in the general setting where  $H$  is not compact, although one has still some useful information about  $\text{Irr}(G)_H$ , see e.g., Theorem 6.2 below.

The following notions are a key in answering Problem 2.3 for  $\Omega = \text{Irr}(G)_H$ .

**Definition 3.1.** Let  $G/H$  be a reductive symmetric space defined by an involution  $\sigma$  of  $G$ . We take  $G_U \subset G_{\mathbb{C}}$  such that  $G_U \cap H$  is a maximal compact subgroup of  $H$ .

- (1) We say a complex parabolic subalgebra  $\mathfrak{q}$  of  $\mathfrak{g}_{\mathbb{C}}$  is a *Borel subalgebra* for  $G/H$  if  $\mathfrak{q}$  is defined by a generic element in  $\sqrt{-1}\mathfrak{g}_U^{-\sigma}$ .
- (2) We say a real parabolic subalgebra  $\mathfrak{p}$  of  $\mathfrak{g}$  is a *minimal parabolic subalgebra* for  $G/H$  if  $\mathfrak{p}$  is defined by a generic element in  $\mathfrak{g} \cap \sqrt{-1}\mathfrak{g}_U^{-\sigma}$ .

Borel subalgebras for the symmetric space  $G/H$  are unique up to inner automorphisms of  $\mathfrak{g}_{\mathbb{C}}$ . Likewise, minimal parabolic subalgebras for  $G/H$  are unique up to inner automorphisms of  $\mathfrak{g}$ . We shall write  $B_{G/H} \subset G_{\mathbb{C}}$  and  $P_{G/H} \subset G$  for the corresponding parabolic subgroups, referred to as a *Borel subgroup* and a *minimal parabolic subgroup* for the symmetric space  $G/H$ , respectively. We note that the Borel subalgebra  $\mathfrak{b}_{G/H}$  for  $G/H$  is not necessarily solvable, and that it is determined only by the complexification  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ .

Here is an answer to Problem 2.3 for  $\Omega = \text{Irr}(G)_H$  when  $(G, H)$  is a reductive symmetric pair.

**Theorem 3.2.** Let  $B_{G/H}$  be a Borel subgroup for  $G/H$ . Suppose  $G'$  is an algebraic reductive subgroup of  $G$ . Then the following three conditions on the triple  $(G, H, G')$  are equivalent:

- (i)  $\sup_{\Pi \in \text{Irr}(G)_H} m(\Pi|_{G'}) < \infty$ .
- (ii)  $G_{\mathbb{C}}/B_{G/H}$  is  $G'_U$ -strongly visible.
- (iii)  $G_{\mathbb{C}}/B_{G/H}$  is  $G'_{\mathbb{C}}$ -spherical.

See [8, Def. 3.3.1] for the definition of strongly visible actions on complex manifolds, and [29] for the equivalence (ii)  $\iff$  (iii).

The list of the triples  $(G, H, G')$  is given in The-

orem 5.1 below in the setting that  $(G, G')$  is a symmetric pair and that  $\mathfrak{g}_{\mathbb{C}}$  is simple.

We also discuss the following finite multiplicity property **(FM)** for the restriction  $\Pi|_{G'}$ , weaker than the bounded multiplicity property (i) in Theorem 3.2:

**(FM)**  $[\Pi|_{G'}; \pi] < \infty, \forall \Pi \in \text{Irr}(G)_H, \forall \pi \in \text{Irr}(G')$ .

**Proposition 3.3.** Let  $P_{G/H}$  be a minimal parabolic subgroup for a reductive symmetric space  $G/H$ . Let  $G'$  be an algebraic reductive subgroup of  $G$ , and  $P'$  a minimal parabolic subgroup of  $G'$ .

- (1) If  $\#(P'_{\mathbb{C}} \backslash G_{\mathbb{C}} / (P_{G/H})_{\mathbb{C}}) < \infty$ , then **(FM)** holds.
- (2) If **(FM)** holds,  $G/P_{G/H}$  is  $G'$ -real spherical.

Proposition 3.3 (2) was proved in [11]. The converse statement of Proposition 3.3 (2) holds in the group manifold case, namely, if  $G/H$  is of the form  $(G \times G) / \text{diag } G$  and if  $G'$  is of the form  $G'_1 \times G'_2$ , see Fact 2.1 (2).

**4. Degenerate principal series representations** Let  $P$  be a parabolic subgroup of  $G$ . We write  $\text{Irr}(P)_f$  for the set of equivalence classes of irreducible finite-dimensional representations of  $P$ . Let  $\text{Ind}_P^G(\xi)$  be the degenerate principal series representation of  $G$  obtained as a smooth induction from  $\xi \in \text{Irr}(P)_f$ . Then  $\text{Ind}_P^G(\xi) \in \mathcal{M}(G)$ .

Suppose that  $P'$  is a parabolic subgroup of a real reductive algebraic subgroup  $G'$  of  $G$ . Degenerate principal series representations  $\text{Ind}_{P'}^{G'}(\eta)$  of  $G'$  are defined similarly for  $\eta \in \text{Irr}(P')_f$ . This section studies the multiplicity  $[\text{Ind}_P^G(\xi)|_{G'} : \text{Ind}_{P'}^{G'}(\eta)]$ , namely, the dimension of the space  $\text{Hom}_{G'}(\text{Ind}_P^G(\xi)|_{G'}, \text{Ind}_{P'}^{G'}(\eta))$  of “symmetry breaking operators”.

In the case  $(G, G') = (O(n+1, 1), O(n, 1))$ , this is the space of conformally covariant symmetry breaking operators for the totally geodesic embedding  $S^{n-1} \hookrightarrow S^n$ . All such operators have been constructed and classified recently, see [18] for the scalar case, and [19] for differential forms. In this case, the multiplicity takes the values in  $\{0, 1, 2\}$ .

For a finer estimate of the multiplicity  $[\text{Ind}_P^G(\xi)|_{G'} : \text{Ind}_{P'}^{G'}(\eta)]$  in the general setting, we implement yet other parabolic subgroups  $Q \subset P_{\mathbb{C}}$  and  $Q' \subset P'_{\mathbb{C}}$ . What we call a “QP estimate” of the multiplicity will play a key role in the proof of Theorem 3.2 for  $H$ -distinguished representations.

Let  $Q$  be a complex parabolic subgroup of  $G_{\mathbb{C}}$  with  $\mathfrak{q} \subset \mathfrak{p}_{\mathbb{C}}$ . We do not require  $\mathfrak{q}$  to be defined over  $\mathbb{R}$ . For  $\xi \in \text{Irr}(P)_f$ , we define  $d_{\mathfrak{q}}(\xi)$  to be the min-

imum of the dimensions of non-zero  $\mathfrak{q}$ -submodules in  $\eta$ , and denote by  $\text{Irr}(P; \mathfrak{q})_f$  the subset of  $\text{Irr}(P)_f$  with  $d_{\mathfrak{q}}(\xi) = 1$ .

We define subsets of  $\mathcal{M}(G)$  by

$$(4.1) \quad \Omega_P := \{\text{Ind}_P^G(\xi) : \xi \text{ is a character of } P\},$$

$$(4.2) \quad \Omega_{P, \mathfrak{q}} := \{\text{Ind}_P^G(\xi) : \xi \in \text{Irr}(P; \mathfrak{q})_f\}.$$

Obviously, one has  $\Omega_P \subset \Omega_{P, \mathfrak{q}}$ . Moreover,  $\Omega_{P, \mathfrak{q}}$  is the whole set  $\{\text{Ind}_P^G(\xi) : \xi \in \text{Irr}(P)_f\}$  if  $\mathfrak{q}$  is a Borel subalgebra of  $\mathfrak{g}_{\mathbb{C}}$ .

We consider the following refinement of Problem 2.3:

**Problem 4.1.** Given a pair  $G \supset G'$ , find subsets  $\Omega \subset \mathcal{M}(G)$  and  $\Omega' \subset \mathcal{M}(G')$  such that

$$\sup_{\Pi \in \Omega} \sup_{\pi \in \Omega'} [\Pi|_{G'} : \pi] < \infty.$$

One observes that Problem 2.3 corresponds to the case where  $\Omega' = \text{Irr}(G')$ .

**Theorem 4.2** (“ $QP$  estimate” for restriction). *Suppose that  $Q$  and  $Q'$  are complex parabolic subgroups of  $G_{\mathbb{C}}$  and  $G'_{\mathbb{C}}$ , respectively, such that  $\mathfrak{q} \subset \mathfrak{p}_{\mathbb{C}}$ ,  $\mathfrak{q}' \subset \mathfrak{p}'_{\mathbb{C}}$ , and  $\#(Q'_{\text{opp}} \backslash G_{\mathbb{C}}/Q) < \infty$ . Here  $Q'_{\text{opp}}$  stands for the opposite parabolic subgroup of  $Q'$  in  $P'_{\mathbb{C}}$ . Then there exists  $C > 0$  such that*

$$(4.3) \quad [\text{Ind}_P^G(\xi)|_{G'} : \text{Ind}_{P'}^{G'}(\eta)] \leq C d_{\mathfrak{q}}(\xi) d_{\mathfrak{q}'}(\eta)$$

for any  $\xi \in \text{Irr}(P)_f$  and any  $\eta \in \text{Irr}(P')_f$ . In particular, one has

$$\sup_{\xi \in \text{Irr}(P; \mathfrak{q})_f} \sup_{\eta \in \text{Irr}(P'; \mathfrak{q}')_f} [\text{Ind}_P^G(\xi)|_{G'} : \text{Ind}_{P'}^{G'}(\eta)] \leq C.$$

When  $Q'$  is a Borel subgroup of  $G'_{\mathbb{C}}$ , one obtains the converse statement of Theorem 4.2 as follows.

**Theorem 4.3.** *Let  $G \supset G'$  be a pair of real reductive algebraic Lie groups,  $P$  a parabolic subgroup of  $G$ , and  $Q$  a complex parabolic subgroup of  $G_{\mathbb{C}}$  such that  $\mathfrak{q} \subset \mathfrak{p}_{\mathbb{C}}$ . Then the following four conditions on  $(G, G'; P, Q)$  are equivalent:*

- (i)  $\sup_{\Pi \in \Omega_{P, \mathfrak{q}}} m(\Pi|_{G'}) < \infty$ .
- (ii) There exists  $C > 0$  such that

$$m(\text{Ind}_P^G(\xi)|_{G'}) \leq C d_{\mathfrak{q}}(\xi) \quad \text{for all } \xi \in \text{Irr}(P)_f.$$

- (iii)  $G_{\mathbb{C}}/Q$  is  $G'_U$ -strongly visible.
- (iv)  $G_{\mathbb{C}}/Q$  is  $G'_{\mathbb{C}}$ -spherical.

The parabolic subgroups  $Q$  in (iv) are classified in [2] in the setting where  $(G_{\mathbb{C}}, G'_{\mathbb{C}})$  is a symmetric pair. Theorem 4.3 with  $Q = P_{\mathbb{C}}$  shows:

**Corollary 4.4.** *Let  $P$  be a parabolic subgroup of  $G$ , and  $G'$  an algebraic subgroup of  $G$ . Then one*

*has the equivalence on the triple  $(G, G'; P)$ :*

$$G_{\mathbb{C}}/P_{\mathbb{C}} \text{ is } G'_{\mathbb{C}}\text{-spherical} \iff \sup_{\Pi \in \Omega_P} m(\Pi|_{G'}) < \infty.$$

**Example 4.5.** If the unipotent radical of  $P$  is abelian, then Corollary 4.4 applies for any symmetric pair  $(G, G')$  by [8, Cor. 15].

Theorem 4.2 also implies the following.

**Theorem 4.6** (Invariant trilinear forms). *Let  $G$  be a real reductive algebraic Lie group, and  $P_j$  ( $j = 1, 2, 3$ ) parabolic subgroups of  $G$ . Suppose that  $Q_j$  ( $j = 1, 2, 3$ ) are complex parabolic subgroups of  $G_{\mathbb{C}}$  such that  $Q_j \subset (P_j)_{\mathbb{C}}$  ( $1 \leq j \leq 3$ ) and  $\#(\text{diag}(G_{\mathbb{C}}) \backslash (G_{\mathbb{C}} \times G_{\mathbb{C}} \times G_{\mathbb{C}}) / (Q_1 \times Q_2 \times Q_3)) < \infty$ . Then there exists  $C > 0$  such that*

$$\dim_{\mathbb{C}} \text{Hom}_G(\bigotimes_{j=1}^3 \text{Ind}_{P_j}^G(\xi_j), \mathbb{C}) \leq C \prod_{j=1}^3 d_{\mathfrak{q}_j}(\xi_j)$$

for all  $\xi_j \in \text{Irr}(P_j)_f$  ( $j = 1, 2, 3$ ).

See [22, 23] for a classification of  $(Q_1, Q_2, Q_3)$  with the above geometric property for some classical groups  $G_{\mathbb{C}}$ .

For  $\Pi_1, \Pi_2 \in \mathcal{M}(G)$ , we consider the tensor product representation  $\Pi_1 \otimes \Pi_2$ , and set

$$m(\Pi_1 \otimes \Pi_2) := \sup_{\Pi \in \text{Irr}(G)} \dim_{\mathbb{C}} \text{Hom}_G(\Pi_1 \otimes \Pi_2, \Pi).$$

A special case of Theorem 4.6 implies (v)  $\Rightarrow$  (i) of the theorem below.

**Theorem 4.7.** *Let  $G$  be a real reductive algebraic Lie group, and  $P_j$  ( $j = 1, 2$ ) parabolic subgroups. Then the following five conditions on the triple  $(G, P_1, P_2)$  are equivalent:*

- (i) There exists  $C > 0$  such that

$$m(\text{Ind}_{P_1}^G(\xi_1) \otimes \text{Ind}_{P_2}^G(\xi_2)) \leq C \dim \xi_1 \dim \xi_2$$

for all  $\xi_j \in \text{Irr}(P_j)_f$  ( $j = 1, 2$ ).

- (ii) There exists  $C > 0$  such that

$$m(\text{Ind}_{P_1}^G(\xi_1) \otimes \text{Ind}_{P_2}^G(\xi_2)) \leq C$$

for all characters  $\xi_j$  of  $P_j$  ( $j = 1, 2$ ).

(iii)  $\mathcal{O}(G_{\mathbb{C}}/P_{1\mathbb{C}}, \mathcal{L}_1) \otimes \mathcal{O}(G_{\mathbb{C}}/P_{2\mathbb{C}}, \mathcal{L}_2)$  is a multiplicity free  $G_{\mathbb{C}}$ -module for any  $G_{\mathbb{C}}$ -equivariant holomorphic line bundles  $\mathcal{L}_j$  on  $G_{\mathbb{C}}/P_{j\mathbb{C}}$  ( $j = 1, 2$ ).

(iv)  $G_{\mathbb{C}}/P_{1\mathbb{C}} \times G_{\mathbb{C}}/P_{2\mathbb{C}}$  is  $\text{diag}(G_U)$ -strongly visible.

(v)  $G_{\mathbb{C}}/P_{1\mathbb{C}} \times G_{\mathbb{C}}/P_{2\mathbb{C}}$  is  $\text{diag}(G_{\mathbb{C}})$ -spherical.

The classification of such pairs  $(P_{1\mathbb{C}}, P_{2\mathbb{C}})$  appeared in different contexts. For instance, one may read from [26] for the multiplicity-free results on finite-dimensional representations (iii). The classification theory of visible actions also gives a complete

$\mathfrak{g}_{\mathbb{C}}$	$\mathfrak{h}_{\mathbb{C}}$	$\mathfrak{g}'_{\mathbb{C}}$
$\mathfrak{sl}_n$	$\mathfrak{gl}_{n-1}$	$\mathfrak{sl}_p \oplus \mathfrak{sl}_q \oplus \mathbb{C}$
$\mathfrak{sl}_{2m}$	$\mathfrak{gl}_{2m-1}$	$\mathfrak{sp}_m$
$\mathfrak{sl}_6$	$\mathfrak{sp}_3$	$\mathfrak{sl}_4 \oplus \mathfrak{sl}_2 \oplus \mathbb{C}$
$\mathfrak{so}_n$	$\mathfrak{so}_{n-1}$	$\mathfrak{so}_p \oplus \mathfrak{so}_q$
$\mathfrak{so}_{2m}$	$\mathfrak{so}_{2m-1}$	$\mathfrak{gl}_m$
$\mathfrak{so}_{2m}$	$\mathfrak{so}_{2m-2} \oplus \mathbb{C}$	$\mathfrak{gl}_m$
$\mathfrak{sp}_n$	$\mathfrak{sp}_{n-1} \oplus \mathfrak{sp}_1$	$\mathfrak{sp}_p \oplus \mathfrak{sp}_q$
$\mathfrak{sp}_n$	$\mathfrak{sp}_{n-2} \oplus \mathfrak{sp}_2$	$\mathfrak{sp}_{n-1} \oplus \mathfrak{sp}_1$
$\mathfrak{e}_6$	$\mathfrak{f}_4$	$\mathfrak{so}_{10} \oplus \mathbb{C}$
$\mathfrak{f}_4$	$\mathfrak{so}_9$	$\mathfrak{so}_9$

Table 5.1. Triples  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  with  $\mathfrak{g}_{\mathbb{C}}$  simple in Theorem 5.1

list of the pairs  $(P_{1\mathbb{C}}, P_{2\mathbb{C}})$  satisfying (iv), see [10] for type A, and [28] for the other cases. See also [21] for the list satisfying (v) when  $P_{j\mathbb{C}}$  are maximal.

**Example 4.8.** Let  $G$  be a real reductive Lie group, and  $P_1, P_2$  parabolic subgroups with abelian unipotent radical. The double flag variety  $G_{\mathbb{C}}/P_{1\mathbb{C}} \times G_{\mathbb{C}}/P_{2\mathbb{C}}$  is strongly visible via the diagonal  $G_U$ -action [9, Thm. 1.7], hence Theorem 4.7 applies. In particular, by taking  $P_2$  to be the opposite parabolic subgroup of  $P_1$ , one sees from Theorem 4.7 the uniform bounded multiplicity property in the Plancherel formula for any para-Hermitian symmetric space.

### 5. Classification of triples $(G, H, G')$

In this section, we present the classification of the triples  $(G, H, G')$  satisfying

$$(5.1) \quad \sup_{\Pi \in \text{Irr}(G)_H} m(\Pi|_{G'}) < \infty$$

on the level of Lie algebras up to outer automorphisms in the following setting:

- both  $(G, H)$  and  $(G, G')$  are symmetric pairs,
- $\mathfrak{g}_{\mathbb{C}}$  is simple.

**Theorem 5.1.** *Suppose that  $\mathfrak{g}_{\mathbb{C}}$  is simple and that  $(G, H)$  and  $(G, G')$  are symmetric pairs. Then the triple  $(G, H, G')$  satisfies the bounded multiplicity property (5.1) if and only if the triple  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  of the complexified Lie algebras is in Table 5.1 or the pair  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  is in (2.3). In the table,  $p, q$  are arbitrary subject to  $n = p + q$ .*

**Example 5.2.** The triple  $(G, H, G')$  in Example 2.2 is a real form of the triple  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  in the fourth row of Table 5.1, hence Theorem 5.1 guarantees the bounded multiplicity property of the restriction  $\Pi|_{G'}$  for all  $\Pi \in \text{Irr}(G)_H$ , see [13, 24].

**Remark 5.3.** When the pair  $(\mathfrak{g}_{\mathbb{C}}, \mathfrak{g}'_{\mathbb{C}})$  is in the list (2.3), the supremum of the multiplicity (2.1)

is equal to one for many of the real forms such as  $(SO(p, q), SO(p-1, q))$ , see [27].

**6. Sketch of the proof for our main results** We give two ingredients that are used in the proof of our main results.

In the classical harmonic analysis on the Riemannian symmetric space  $G/K$ , building blocks of representations in  $C^\infty(G/K)$  are constructed by the twisted Poisson transform, an integral  $G$ -intertwining operator from the spherical principal series representation to  $C^\infty(G/K)$ . More generally, for a closed subgroup  $H$  in  $G$ , we consider the space  $\text{Hom}_G(\text{Ind}_P^G(\xi), \text{Ind}_H^G(\tau))$  of generalized Poisson transforms, where  $P$  is a parabolic subgroup of  $G$ ,  $\xi \in \text{Irr}(P)_f$ , and  $\tau \in \text{Irr}(H)_f$ . We give a “ $QP$  estimate” of the dimension of this space. Along the same line as in [11, 16], the “ $QP$  estimate” for restriction (e.g., the implication (iv)  $\Rightarrow$  (i) in Theorem 4.3) is deduced from the following “ $QP$  estimates for induction” applied to  $(G \times G')/\text{diag } G'$ . Theorem 6.1 (1) below is a generalization of some results in [16] relying on the “boundary valued maps” and in Tauchi [30] relying on the theory of holonomic  $\mathcal{D}$ -modules [3, 4].

**Theorem 6.1** (“ $QP$  estimate” for induction). *Let  $G$  be a real reductive algebraic Lie group,  $H$  an algebraic subgroup,  $P$  a parabolic subgroup of  $G$ , and  $Q$  a complex parabolic subgroup of  $G_{\mathbb{C}}$  with  $Q \subset P_{\mathbb{C}}$ .*

- (1) *If  $\#(Q \backslash G_{\mathbb{C}}/H_{\mathbb{C}}) < \infty$ , then there exists  $C > 0$  such that for all  $\eta \in \text{Irr}(P)_f$  and all  $\tau \in \text{Irr}(H)_f$*

$$\dim_{\mathbb{C}} \text{Hom}_G(\text{Ind}_P^G(\eta), \text{Ind}_H^G(\tau)) \leq C d_q(\eta) \dim \tau.$$

- (2) *Conversely, if the conclusion in (1) holds, then  $Q$  has an open orbit in  $G_{\mathbb{C}}/H_{\mathbb{C}}$ .*

For the proof of Theorem 3.2, we also use the following reformulation [14] of Casselman–Oshima’s subrepresentation theorem [25, 31].

**Theorem 6.2** (Quotient representation theorem). *Let  $G/H$  be a reductive symmetric space, and  $P_{G/H}$  and  $\mathfrak{b}_{G/H}$  a minimal parabolic subgroup and a Borel subalgebra for  $G/H$ , respectively, with  $\mathfrak{b}_{G/H} \subset (\mathfrak{p}_{G/H})_{\mathbb{C}}$ . Then for any  $\Pi \in \text{Irr}(G)_H$ , there exists  $\xi \in \text{Irr}(P_{G/H}; \mathfrak{b}_{G/H})_f$  such that  $\Pi$  is a quotient of the degenerate principal series representation  $\text{Ind}_{P_{G/H}}^G(\xi)$ .*

**(Acknowledgements)** This work was partially supported by Grant-in-Aid for Scientific Research (A) (18H03669), Japan Society for the Promotion of Science.

## References

- [ 1 ] J.-L. Clerc, T. Kobayashi, B. Ørsted, and M. Pevzner, *Generalized Bernstein–Reznikov integrals*, *Math. Ann.*, **349** (2011), pp. 395–431.
- [ 2 ] X. He, K. Nishiyama, H. Ochiai, Y. Oshima, On orbits in double flag varieties for symmetric pairs, *Transform. Groups*, **18**, (2013), 1091–1136.
- [ 3 ] M. Kashiwara, *Systems of Microdifferential Equations*, *Progr. Math.*, **34**, Birkhäuser, Boston, 1983, xv+159pp.
- [ 4 ] M. Kashiwara, T. Kawai, On holonomic systems of microdifferential equations. III. Systems with regular singularities, *Publ. Res. Inst. Math. Sci.* **17**, (1981), 813–979.
- [ 5 ] T. Kobayashi, *Discrete decomposability of the restriction of  $A_q(\lambda)$  with respect to reductive subgroups and its applications*, *Invent. Math.*, **117** (1994), 181–205.
- [ 6 ] T. Kobayashi, *Discrete decomposability of the restriction of  $A_q(\lambda)$  with respect to reductive subgroups II—micro-local analysis and asymptotic  $K$ -support*, *Ann. of Math.*, **147** (1998), pp. 709–729.
- [ 7 ] T. Kobayashi, *Discrete decomposability of the restriction of  $A_q(\lambda)$  with respect to reductive subgroups III—restriction of Harish-Chandra modules and associated varieties*, *Invent. Math.*, **131** (1998), 229–256.
- [ 8 ] T. Kobayashi, *Multiplicity-free representations and visible actions on complex manifolds*, *Publ. Res. Inst. Math. Sci.* **41** (2005), pp. 497–549, special issue commemorating the fortieth anniversary of the founding of RIMS.
- [ 9 ] T. Kobayashi, Visible actions on symmetric spaces, *Transf. Groups*, **12**, (2007), 671–694.
- [10] T. Kobayashi, *A generalized Cartan decomposition for the double coset space  $(U(n_1) \times U(n_2) \times U(n_3)) \backslash U(n) / (U(p) \times U(q))$* , *J. Math. Soc. Japan* **59** (2007), pp. 669–691.
- [11] T. Kobayashi, *Shintani functions, real spherical manifolds, and symmetry breaking operators*, *Dev. Math.*, **37**, (2014), 127–159. Springer.
- [12] T. Kobayashi, *A program for branching problems in the representation theory of real reductive groups*, In: *Representations of Reductive Groups—In Honor of the 60th Birthday of David A. Vogan, Jr.*, *Progr. Math.*, **312**, Birkhäuser, 2015, 277–322.
- [13] T. Kobayashi, Branching laws of unitary representations associated to minimal elliptic orbits for indefinite orthogonal group  $O(p, q)$ , *Adv. Math.*, **388**, (2021), Paper No. 107862, 38pp.
- [14] T. Kobayashi, Bounded multiplicity theorems for induction and restriction, *J. Lie Theory* **32**, (2022), no. 1, 197–238.
- [15] T. Kobayashi, T. Matsuki, *Classification of finite-multiplicity symmetric pairs*, *Transform. Groups*, **19** (2014), 457–493, Special issue in honor of Dynkin for his 90th birthday.
- [16] T. Kobayashi, T. Oshima, *Finite multiplicity theorems for induction and restriction*, *Adv. Math.*, **248** (2013), 921–944.
- [17] T. Kobayashi, Y. Oshima, *Classification of symmetric pairs with discretely decomposable restrictions of  $(\mathfrak{g}, K)$ -modules*, *J. Reine Angew. Math.* **703**, (2015), 201–223.
- [18] T. Kobayashi, B. Speh, Symmetry Breaking for Representations of Rank One Orthogonal Groups, (2015), *Mem. Amer. Math. Soc.* **238** no.1126, 118 pages.
- [19] T. Kobayashi, B. Speh, Symmetry Breaking for Representations of Rank One Orthogonal Groups, Part II, *Lecture Notes in Math.*, **2234** Springer, 2018. xv+342 pages.
- [20] M. Krämer, *Multiplicity free subgroups of compact connected Lie groups*, *Arch. Math. (Basel)* **27**, (1976), 28–36.
- [21] P. Littelmann, On spherical double cones, *J. Algebra* **166**, (1994), 142–157.
- [22] P. Magyoar, J. Weyman, A. Zelevinsky, Multiple flag varieties of finite type, *Adv. Math.*, **141**, (1999), 97–118.
- [23] T. Matsuki, Orthogonal multiple flag varieties of finite type I: Odd degree case, *J. Algebra*, **425**, (2015), 450–523.
- [24] B. Ørsted, B. Speh, Branching laws for discrete series of some affine symmetric spaces, preprint, arXiv:1907.07544.
- [25] T. Oshima, A realization of semisimple symmetric spaces and construction of boundary value maps. *Advanced Studies in Pure Mathematics*, **14**, (1988), 603–650.
- [26] J. R. Stembridge, Multiplicity-free products and restrictions of Weyl characters, *Represent. Theory* **7**, (2003), 404–439.
- [27] B. Sun, C.-B. Zhu, Multiplicity one theorems: the Archimedean case, *Ann. of Math.*, **175**, (2012), pp. 23–44.
- [28] Y. Tanaka, Classification of visible actions on flag varieties, *Proc. Japan Acad. Ser. A Math. Sci.*, **88**, (2012), 91–96.
- [29] Y. Tanaka, Visible actions of compact Lie groups on complex spherical varieties, to appear in *J. Diff. Geom.*
- [30] T. Tauchi, A generalization of the Kobayashi–Oshima uniformly bounded multiplicity theorem, *Internat. J. Math.*, **32**, (2021), no. 14, Paper No. 2150107, 25pp.
- [31] N. R. Wallach, *Real reductive groups. I, II*, *Pure Appl. Math.* **132** Academic Press, Inc., Boston, MA, 1988; **132-II**, *ibid*, 1992.